9ο Πανελλήνιο Συνέδριο Γεωμετρίας

29-31 Μαΐου, 2009, Ακαδημαϊκό Χωριό Ανωγείων

Περιλήψεις των διαλέξεων

Τμήμα Μαθηματικών του Πανεπιστημίου Κρήτης

9ο Πανελλήνιο Συνέδριο Γεωμετρίας

29-31 Μαΐου, 2009, Ακαδημαϊκό Χωριό Ανωγείων

Ιστοσελίδα: http://www.math.uoc.gr/∼geomconf09
e-mail: geomconf09@math.uoc.gr

Οργανωτική Επιτροπή:

Κ. Αθανασόπουλος
Χ. Κουρουνιώτης
Μ. Λυδάκης
Ι. Πλατής

Herbert Abels

Fakultät für Mathematik, Universität Bielefeld

Affine crystallographic and properly discontinuous groups

There is a long standing conjecture of Auslander (1964) which states that every affine crystallographic group has a solvable subgroup of finite index. Milnor (1977) asked if, more generally, every properly discontinuous affine group has a solvable subgroup of finite index. This more general conjecture was disproved by Margulis (1983). The Auslander conjecture has been proved in many cases, but is wide open in dimension at least 7.

I will explain the geometric and algebraic relevance of the notions and conjectures and will give an idea of the methods involved in the constructions and proofs in our joint work with Margulis and Soifer. We use dynamics of linear and affine maps. The connection with the Tits alternative and proximal maps will be pointed out.

Κωνσταντίνος Αθανασόπουλος

Τμήμα Μαθηματικών, Πανεπιστήμιο Κρήτης

An algorithm for the equipartition of curves

In 1940 J.F. Pal posed the following problem: Let (X, ρ) be a metric space and let $c : [0, 1] \to X$ be a continuous curve such that $c(0) \neq c(1)$. Is there a partition $0 < t_1 < \cdots < t_n < 1$ such that $\rho(c(t_{i-1}), c(t_i)) = \rho(c(t_i), c(t_{i+1}))$, where $t_0 = 0$ and $t_{n+1} = 1$? K. Urbanik (1954) and M.A. Lopez and S. Reisner (2007) have given proofs of the existence of a solution using the Brouwer fixed point theorem. Our approach is based on the topological study of the zero level set of a certain continuous function defined on the n-dimensional simplex and leads to an approximate algorithm for finding all solutions with high accuracy. This is joint work with C. Panagiotakis and G. Tziritas (Computer Science Department).

Ιάκωβος Ανδρουλιδάκης

Centro de Matematica da Universidade do Porto

Pseudodifferential calculus for singular foliations

We discuss the definition of pseudodifferential operators (bounded and unbounded) along the leaves of a singular foliation. The analytic index of such operators takes values in the K-theory of the C^*-algebra of a singular foliation, which is constructed by the method of bi-submersions. This is joint work with G. Skandalis.

Ανδρέας Αρβανιτογεώργος

Τμήμα Μαθηματικών, Πανεπιστήμιο Πατρών

Einstein metrics on compact Lie groups which are not naturally reductive

A Riemannian manifold (M, g) is called Einstein if the Ricci tensor $Ric(g)$ of the metric g satisfies $Ric(g) = eg$ for some constant e. For the case of compact Lie groups, left-invariant Einstein metrics have not been studied extensively. Even in low dimensional examples such as $SU(3)$ and $SU(2) \times SU(2)$, the number of left-invariant Einstein metrics is still unknown. The only complete work is the one by J.E. D'Atri and W. Ziller Naturally reductive metrics and Einstein metrics on compact Lie groups, (Mem. Amer. Math. Soc. 18 (215) (1979)), in which they obtained a large number of left-invariant Einstein metrics which are naturally reductive.

The problem of finding non-naturally reductive left-invariant Einstein metrics on compact Lie groups seems to be harder, and in fact this is stressed by the above authors. In 1996 K. Mori obtained non-naturally reductive Einstein metrics on the Lie group $SU(n)$ for $n > 5$, by using a method of Riemannian submersions (Left invariant Einstein metrics on $SU(N)$ that are not naturally reductive, Master Thesis (in Japanese) Osaka University 1994, English translation Osaka University RPM 96 – 10 (preprint series) 1996).

In the present work we prove existence of non-naturally reductive Einstein metrics on the compact simple Lie groups $SO(n)$ $(n > 10)$, $Sp(n)$ $(n > 2)$, $E6$, $E7$, and $E8$.

Γεώργιος Καϊμακάμης

Στρατιωτική Σχολή Ευελπίδων

Lorentz hypersurfaces in E_1^4 satisfying $\Delta H = aH$

A hypersurface M_1^3 in the four-dimensional pseudo-Euclidean space E_1^4 is called a Lorentz hypersurface if its normal vector is space-like. We show that if the mean curvature vector field of M_1^3 satisfies the equation $\Delta H = aH$ (a constant), then M_1^3 has constant mean curvature. This equation is a natural generalization of the biharmonic submanifold equation $\Delta H = 0$.

Θέμης Κουφογιώργος

Τμήμα Μαθηματικών, Πανεπιστήμιο Ιωαννίνων

3-dimensional contact metric manifolds with $Trl = c$ (const.)

In contact geometry, the Jacobi operator $l = R(., x)x$ plays a fundamental role. The class of contact metric manifolds with $Trl = c$ (const.) is particularly large, since, among others it includes the well known classes of Sasakian manifolds and (κ, μ)-contact manifolds. In the present work, a complete local description of 3-dimensional contact metric manifolds with $Trl = c$, is given. As a consequence of it, several new examples of such manifolds are obtained.

Μιχαήλ Μάρκελλος και Βασίλης Παπαντωνίου

Τμήμα Μαθηματικών, Πανεπιστήμιο Πατρών

Ricci solitons and contact manifolds

In [3], the authors introduced the notion of $(\kappa; \mu, \nu)$-contact metric manifolds in terms of a specifc curvature condition, where κ, μ, ν are smooth functions. Especially, they proved that a contact metric 3-manifold $[M; (\eta; \xi; \phi; g)]$ is a $(\kappa; \mu, \nu)$-contact metric manifold on an everywhere open and dense subset of M if and only if the characteristic vector field ξ is an harmonic vector field. Recently, Cho ([1]) proved that an odd-dimensional sphere does not admit a Ricci soliton whose potential vector field is the Reeb vector field. More generally, in [2] the authors proved that if the metric of a non-Sasakian $(\kappa; \mu)$-contact manifold $[M^{2n+1}; (\eta; \xi; \phi; g)]$ is a gradient Ricci soliton, then M is locally flat in dimension three, and locally isometric to $E^{n+1} \times S^n(4)$ in higher dimensions. Extending the above results, we prove that every 3-dimensional $(\kappa; \mu, \nu)$-contact metric manifold $[M; (\eta; \xi; \phi; g)]$ with κ non-constant smooth function on M, $\kappa < 1$ everywhere on M and the function ν or μ constant on M, does not admit a gradient Ricci soliton.

In the sequel, we concentrate on the unit tangent sphere bundle T_1M of a complete Riemannian manifold $(M^n; g)$ $(n \geq 2)$ equipped with the standard contact metric structure $(\eta; \xi; \phi; \bar{g})$, where the metric $\bar{g}$ is homothetic to the Sasaki metric g_S and ξ is the geodesic flow of M^n. More precisely, we prove that the metric $\bar{g}$ is not a Ricci soliton with potential vector field the geodesic flow ξ. Moreover, we construct examples of potential vector fields with respect to the metric $\bar{g}$ of $T_1\mathbb{R}^3$.

References

1. Cho J.T. Ricci solitons and odd-dimensional spheres, *Monatsh. Math.(to appear).*

2. Ghosh Amalendu, Sharma Ramesh, and Cho Jong Taek. Contact metric manifolds with η-parallel torsion tensor, *Ann. Glob. Anal. Geom.*, 34(3):287-299, 2008.

3. Koufogiorgos Themis, Markellos Michael, and Papantoniou Vassilis J. The harmonicity of the Reeb vector field on contact metric 3-manifolds, *Pacific J.Math.*, 234(2):325-344 , 2008.

Ευαγγελία Μουτάφη

Γυμνάσιο Κορυφής Νομού Ημαθίας και Τμήμα Μαθηματικών του Α.Π.Θ.

Some Classes of 3-dimensional Pseudo-symmetric Contact Metric Manifolds

We classify
a) the class of 3-dimensional pseudo-symmetric contact metric manifolds of constant type satisfying $\nabla_\xi \tau = 2a\tau\phi$, where a is a smooth function,
b) the class of 3-dimensional pseudo-symmetric contact metric manifolds satisfying $Q\xi = \rho\xi$, where ρ is a smooth function constant in the characteristic direction ξ,
c) the class of pseudo-symmetric complete (κ, μ, ν)-contact metric 3-manifolds of type constant in the characteristic direction ξ and
d) the class of 3-dimensional pseudo-symmetric contact metric manifolds with harmonic curvature and Trl constant in the direction ξ.

Ben McReynolds

Department of Mathematics, University of Chicago.

Controlling manifold covers

I will discuss a generalization of Selberg's lemma and some geometric applications. Most notable is the resolution of Nimershiem's conjecture on the geometry of cusp cross sections of finite volume, complete, hyperbolic n-manifolds.

Ιωάννης Παπαδοπεράκης

Γενικό Τμήμα, Γεωπονικό Πανεπιστήμιο

Υπερβολικοί χώροι και το θεώρημα μοναδικότητας του Mostow

Μελετάμε τις πλήρεις υπερβολικές 3-διάστατες πολλαπλότητες χωρίς σύνορο που κατασκευάζονται με την συγκόλληση υπερβολικών ιδεωδών τετραέδρων. Η συγκόλληση γίνεται πάνω στις πλευρές των τετραέδρων. Οι πολλαπλότητες που δημιουργούνται έχουν παντού καμπυλότητα -1 εκτώς ίσως από κάποιους άξονες που η γωνία γύρω από αυτούς είναι διαφορετική από 2π, άξονες δε του χώρου λέμε τις γεωδαισιακές που προέρχοναι από τις ακμές των τετραέδρων. Οι πολλαπλότητες αυτές είναι εν γένει υπερβολικές κατά Gromov. Σταθεροποιώντας τον τρόπο συγκόλλησης και μεταβάλοντας τις διέδρες γωνίες των τετραέδρων μελετάμε, κάτω από ισοτοπία, τον χώρο των δομών. Σ' αυτόν τον χώρο Teichmüller εξετάζουμε τη δυνατότητα παραμέτρισής του από τις γωνίες των αξόνων και την συσχέτισή του με το θεώρημα μοναδικότητας του Mostow. Δίδονται χαρακτηριστικά παραδείγματα.

Δέσποινα Παπαδοπούλου

Τμήμα Μαθηματικών, Α.Π.Θ.

Ύπαρξη σμηνών ευθειών με κοινές αναλλοιώτους.

Στον χώρο E^3 μελετούμε προβλήματα ύπαρξης και καθορισμού σμηνών ευθειών, που έχουν κοινές αναλλοίωτες ποσότητες του σμήνους, όπως η ορική απόσταση, η καμπυλότητα ή η μέση καμπυλότητα αυτού.

Κωνσταντίνος Πετούμενος

Τμήμα Μαθηματικών, Πανεπιστήμιο Πατρών

Biharmonic hypersurfaces of type M_2^3 in E_2^4.

A submanifold M_r^n of a pseudo-Euclidean space E_s^m is said to have harmonic mean curvature vector field if $\triangle\overrightarrow{H} = \overrightarrow{0}$, where $\overrightarrow{H}$ denotes the mean curvature vector field and $\triangle$ the Laplacian of M_r^n with respect to the induced pseudo-Riemannian metric. Submanifolds with harmonic mean curvature vector field are also known as biharmonic submanifolds. In the present talk, by using all possible canonical forms of the shape operator of the pseudo-Euclidean hypersurface M_2^3 in E_2^4 we prove that every nondegenerate biharmonic hypersurface M_2^3 in E_2^4, is minimal.

Ιωάννης Πλατής

Τμήμα Μαθηματικών, Πανεπιστήμιο Κρήτης

The $PU(2,1)$ configuration space of four points in S^3 and the cross-ratio variety

In the classical case, the configuration space of four points in S^2 is biholomorphic to $\mathbb{C} \setminus \{0,1\}$. Things are much more complicated in the configuration space of four points in S^3. We present the latest results on the subject and pose some open questions.

Gregory Soifer

Department of Mathematics and Statistics, Bar-Ilan University

Affine crystallographic semigroups acting on a hyperbolic space

Let $G_n = \mathrm{Aff}(\mathbb{R}^n)$ be the group of affine transformations of $\mathbb{R}^n$. The group G_n is the semidirect product $GL_n(\mathbb{R}) \ltimes \mathbb{R}^n$ where $\mathbb{R}^n$ is identified with its group of translations. A semigroup S of G_n is said to act *properly discontinuous* on $\mathbb{R}^n$ if for every compact subset K of $\mathbb{R}^n$ the set $\{g \in S : gK \cap K \neq \emptyset\}$ is finite. If a discrete group consists of isometries, then it acts properly on $\mathbb{R}^n$. But this is not true for an arbitrary discrete subgroup of G_n, e.g. for an infinite discrete subgroup of $GL_n(\mathbb{R})$. A semigroup S of G_n is called *crystallographic* if S acts properly discontinuous on $\mathbb{R}^n$ and there exists a compact subset K_0 of $\mathbb{R}^n$ such that $\bigcup_{s\in S} sK_0 = \mathbb{R}^n$.

If the signature of a non degenerate quadratic form B on $\mathbb{R}^n$ is $(n-1,1)$, then the form B is called *hyperbolic*.

The motivating question here is the following:

Question. (H.Abels, G.Margulis, G.Soifer). *Is the Zariski closure of a crystallographic affine semigroup leaving a hyperbolic form invariant a virtually solvable group ?*

Our interest in these questions came from our work on the Auslander conjecture. The goal of my talk is to explain a main ideas of our proof of the following

Main Theorem. *Let S be a crystallographic semigroup in Aff$(\mathbb{R}^n)$, $n \leq 3$. Then the Zariski closure of S is a virtually solvable group.*

Γεώργιος Τσαπόγας

Τμήμα Μαθηματικών, Πανεπιστήμιο του Αιγαίου

On the Mapping class group of a handlebody

For a surface Σ, the complex of curves introduced by Harvey is a simplicial complex on which the mapping class group $MCG(\Sigma)$ acts simplicially. This action is widely used for the study of $MCG(\Sigma)$. For a handlebody H, we introduce a complex of incompressible surfaces (in H) which naturally contains the complex of curves for the boundary surface of H. A bijection between the $MCG(H)$ and the group of automorphisms of the complex of incompressible surfaces is established by induction, where the important initial case of a genus 2 handlebody is studied by analyzing the links of the vertices from the graph theoretical point of view.

Χαράλαμπος Τσιχλιάς

Τμήμα Μαθηματικών, Πανεπιστήμιο Ιωαννίνων

3-dimensional contact metric manifolds with $\nabla_\xi \tau = 0$

In contact geometry, the tensor $\tau = \mathcal{L}_\xi g$ plays a fundamental role. In the present work, a complete local description of 3-dimensional contact metric manifolds with $\nabla_\xi \tau = 0$ is given. Moreover, we obtain several new examples of this class of manifolds.

Ανέστης Φωτιάδης

Τμήμα Μαθηματικών και Στατιστικής, Πανεπιστήμιο Κύπρου

Το πρόβλημα του Dirichlet για αρμονικές συναρτήσεις συμμετρικών χώρων

Θα γενικεύσουμε τον ορισμό των αρμονικών συναρτήσεων σε ορισμό που αναφέρεται σε απεικονίσεις μεταξύ πολλαπλότήτων και θα ασχοληθούμε με το σχετικό πρόβλημα του Dirichlet σε συμμετρικούς χώρους.

Ιωάννης Χρυσικός

Τμήμα Μαθηματικών, Πανεπιστήμιο Πατρών

Homogeneous Einstein metrics on generalized flag manifolds

A Riemannian manifold (M, g) is called Einstein if the Ricci tensor Ric_g is a constant multiple of the metric g, i.e. $Ric_g = \Lambda \cdot g$, for some $\Lambda \in \mathbb{R}$. If M is compact, Einstein metrics of volume one can be characterized variationally as the critical points of the scalar curvature functional $T(g) = \int_M S_g dvol_g$ on the space $\mathcal{M}_1$ of Riemannian metrics of volume one. At the present time no general existence results of Einstein metrics are known. Important progress has been made when M is a homogenous space [3]. In this case we are mainly interested in G-invariant Einstein metrics on M. Such a metric is precisely a critical point of T restricted on $\mathcal{M}_1^G$, the set of G-invariant metrics of volume one [1].

An important class of compact homogeneous spaces consists of generalized flag manifolds. These are the adjoint orbits of a compact connected semisimple Lie group G and are complex homogeneous spaces of the form G/K, where K is the centralizer of a torus in G. In [2], the Einstein equation explicity solved for all flag manifolds with three isotropy summands. We solve explicity the Einstein equation for all flag manifolds with two and four isotropy summands. First we classify these spaces by using the notion of t-roots. Next we compute the Ricci tensor for a G-invariant metric on $M = G/K$ and we present new G-invariant Einstein metrics on M. For the case which corresponds to two isotropy summands we also exam the nature of the new Einstein metrics, as critical points of the restriction $T|_{\mathcal{M}_1^G}$.

References

1. A. L. Besse: Einstein Manifolds, Springer-Verlag, Berlin, 1986.

2. M. Kimura: Homogeneous Einstein metrics on certain Kähler C-spaces, Adv. Stud. Pure. Math., 18-I (1990) 303-320.

3. M. Wang and W. Ziller: Existence and non-existence of Homogeneous Einstein metrics, Inventiones Math., 84 (1986) 177-194.