

The Effective Dynamics of the Volume Preserving Mean Curvature Flow

Ilias Chenn¹ · G. Fournodavlos² · I. M. Sigal¹ 

Received: 7 December 2017 / Accepted: 13 April 2018
© Springer Science+Business Media, LLC, part of Springer Nature 2018

Abstract We consider the dynamics of small closed submanifolds (‘bubbles’) under the volume preserving mean curvature flow. We construct a map from $(n + 1)$ -dimensional Euclidean space into a given $(n + 1)$ -dimensional Riemannian manifold which characterizes the existence, stability and dynamics of constant mean curvature submanifolds. This is done in terms of a reduced area function on the Euclidean space, which is given constructively and can be computed perturbatively. This allows us to derive adiabatic and effective dynamics of the bubbles. The results can be mapped by rescaling to the dynamics of fixed size bubbles in almost Euclidean Riemannian manifolds.

Keywords Volume preserving mean curvature flow · Constant mean curvature surface · Submanifold · Mean curvature flow · Adiabatic dynamic · Effective dynamic

1 Background and Main Results

The goal of this note is to introduce the Lyapunov–Schmidt map extending the Feshbach–Schur one ([7], see also [8, 19])¹ to a nonlinear setting and to use it to derive the adiabatic (and effective) dynamics of submanifolds under the volume preserving mean curvature flow.

¹ The Feshbach–Schur map was introduced in the context of quantum electrodynamics and was used in statistical mechanics and random Schrödinger operators in [25–27] and [10], respectively (see also [14, 15]).

To Jürg, Tom and Herbert with friendship and admiration.

✉ I. M. Sigal
im.sigal@utoronto.ca
Ilias Chenn
il.chen@mail.utoronto.c
G. Fournodavlos
gf313@dpmms.cam.ac.uk

¹ Department of Mathematics, University of Toronto, Toronto, Canada

² DPMMS, Centre for Mathematical Sciences, Wilberforce Road, Cambridge CB3 0WB, UK

Let (M, g) be an $n + 1$ -dim oriented Riemannian manifold. Recall that a family of immersions $\psi(t, \cdot) : \Sigma \rightarrow M, \psi(t, \Sigma) := S_t, t \geq 0$, of compact, connected, orientable manifolds² is called the volume preserving mean curvature flow (VPF), iff it satisfies the equation

$$\partial_t \psi^N = \bar{H}(\psi) - H(\psi), \tag{1.1}$$

where $\partial_t \psi^N$ is the normal component of the velocity vector $\partial_t \psi$, defined as $\partial_t \psi^N = g(\partial_t \psi, \nu(\psi))$, with $\nu(\psi)$, the outward unit normal to S_t , is the normal component of the velocity vector $\partial_t \psi$. Here $H(\psi)$ is the mean curvature of S_t and $\bar{H} = \bar{H}(t)$ its mean value over S_t ,

$$\bar{H} := \frac{\int_{S_t} H d\sigma}{\int_{S_t} d\sigma}. \tag{1.2}$$

The VPF appeared in material sciences almost a century ago in modeling cell, grain and bubble growth, etc. It has been derived—as the sharp interface limit (non-rigorously)—from the Kawasaki dynamics in the Ising or Potts model, or from the phase field models of the Cahn–Hilliard type (see e.g. [11, 12]), which themselves can be considered as a (formal) hydrodynamic limit of a microscopic dynamics, say, the stochastic Ginzburg–Landau model (see Spohn [28]).

The static solutions to (1.1) are exactly the constant mean curvature immersions

$$H = \bar{H} \equiv h. \tag{1.3}$$

They appear in nature as (almost) stationary interfaces, e.g. in block copolymer melts ([23], see Fig. 1). (Because of close packing, surfaces deviate from CMC surfaces.)

As a consequence of (1.1), the enclosed volume of S_t is constant in time, while its area is decreasing (see below).

Let $A(\psi)$ and $V_{\text{enc}}(\psi)$ denote the area and enclosed volume of a closed surface given by an immersion ψ . In what follows we take $\Sigma = \mathbb{S}^n$, the standard unit n -dimensional sphere. Let

$$X_c^k := \{\psi \in H^k(\mathbb{S}^n, M) : V_{\text{enc}}(\psi) = c\}. \tag{1.4}$$


Given $c > 0$ small and $z \in \mathbb{M}^{n+1}$, there exists a unique geodesic sphere, $\theta_{\lambda,z} \in X_c^k$, of a radius $\lambda = \lambda(c)$, centred at z (λ depends on z as well), having enclosed volume c . The function $\lambda = \lambda(c)$ is invertible. We denote its inverse by $c(\lambda)$. The main result of this note is the following

Theorem 1.1 *For λ sufficiently small, there exists a family of maps $\Phi_\lambda : M \rightarrow X_{c(\lambda)}^k, r > n/2 + 2$, defined constructively in (2.16) below, such that*

- (i) *For h large, there is an invertible function $\lambda = \lambda(h)$, s.t. $\psi_{\lambda,z} = \Phi_\lambda(z)$, with $\lambda = \lambda(h)$, solves (1.3), iff z is a critical point of $A(\Phi_\lambda(z))$ on $\{V_{\text{enc}}(\Phi_\lambda(z)) = c\}$.*
- (ii) *Let z_* be a critical point of $A(\Phi_\lambda(z))$ on $\{V_{\text{enc}}(\Phi_\lambda(z)) = c\}$. Then the surface ψ_{λ,z_*} is a (non-degenerate) local minimum/(saddle or maximum) point of the area functional $A(\psi)$ on the set $\{V_{\text{enc}}(\psi) = c\}$ iff z_* is a (non-degenerate) local minimum/(saddle or local maximum) point of the function of $A(\Phi_\lambda(z))$ on $\{V_{\text{enc}}(\Phi_\lambda(z)) = c\}$.*
- (iii) *With z_* as above, the surface ψ_{λ,z_*} is asymptotically stable, if z_* is a strict local minimum point of $A(\Phi_\lambda(z))$ on $\{V_{\text{enc}}(\Phi_\lambda(z)) = c\}$ and unstable if z_* is a saddle or local maximum point.*

² We choose their orientation to be compatible with that of M .

Fig. 1 Interfacial curvature, H , distributions for the C, G, PL and D microstructures. The green patches on the schematic diagrams to the left indicate elementary interfacial units. The detailed shapes of these units are then displayed on the right by means of a SCFT calculation along the L/C phase boundary at χ , $N = 20$ and $f = 0.3378$. The curvature distribution over each unit is specified by the colour scale with the average and standard deviation quoted to the far right. Adapted from [69]



We prove this theorem in Sect. 2. It can also be essentially deduced using the proofs and arguments in Alikakos and Freire [1]. Its main value is a new viewpoint.

The map Φ_λ is constructed using the well-known Lyapunov - Schmidt argument. We call it the *Lyapunov-Schmidt map*.³ Abstractly and informally, it is defined as follows. Let $Y \subset X$ be two Hilbert spaces. Given an orthogonal projection P on X and a point $u \in Y$, we say that a differentiable map $G : Y \rightarrow X$ is in the domain of Φ_{Pu} iff the operator $\bar{L} := \bar{P}dG(u)\bar{P}$, where $\bar{P} := \mathbf{1} - P$, is invertible on $\text{Ran } \bar{P}$. For such a G , we define the new map $\Phi_{Pu}(G) : PY \rightarrow PX$ as

$$\Phi_{Pu}(G)(v) := PG(u + v + w(v)),$$

where $v \in PY$ and $w \equiv w(v)$ is a solution to the equation $\bar{P}G(u + v + w) = 0$ in $w \in \text{Ran } \bar{P}Y$. (The requirement that the operator \bar{L} is invertible on $\text{Ran } \bar{P}$ and elementary smoothness properties guarantee that the equation $\bar{P}G(u + v + w) = 0$ has a unique solution in an appropriate space.) We call Φ_{Pu} the *Lyapunov-Schmidt map*. One can compose such maps with varying P and u (possibly also with rescaling) to obtain a discrete dynamical system - a renormalization group.

Coming to the notion of stability, we will work with the following definitions. Recall (see Proposition 1.5 below) that the Gâteaux derivative dH of the mean curvature H is the hessian A'' of the area functional A .

Definition 1.2 *Linear stability* (or just stability in geometric analysis): We say that a (critical) hypersurface $\psi_* \in X_c^k$ is (linearly) stable, if $dH(\psi_*) = A''(\psi_*) \geq 0$ on $T_{\psi_*} X_c^k$. On the other

³ Like the Feshbach-Schur map, which comes from reconceptualizing of the well-known Feshbach-Schur perturbation theory, the Lyapunov-Schmidt map comes from rethinking the well-known Lyapunov-Schmidt reduction technique.

hand, $\psi_* \in X_c^k$ will be called (linearly) unstable if $dH(\psi_*) = A''(\psi_*)$ on $T_{\psi_*}X_c^k$ has a negative spectrum.

Asymptotic stability A CMC hypersurface $\psi_* \in X_c^k$ is asymptotically stable, iff there exists an $\varepsilon > 0$, such that for every initial hypersurface $\psi_0 \varepsilon$ -close to ψ_* in the H^k norm, $k > \frac{n}{2} + 2$, the solution converges (possibly after reparametrizations) to ψ_* , as $t \rightarrow +\infty$.

The theorem above, together with a refinement (due to [24, 29]) of beautiful results of [18] (see also [30]) about the area and enclosed volume expansions for small geodesic spheres, gives the following results due to Ye [29], Huisken and Yau [22], Alikakos and Freire [1], Pacard and Xu [24] (the existence part) and Alikakos and Freire ([1], the stability part):

Theorem 1.3 *For each non-degenerate critical point z_* of the scalar curvature R of M^{n+1} and for each $c > 0$ sufficiently small:*

- (i) *There exists a constant mean curvature hypersurface of enclosed volume c , enclosing z_* .*
- (ii) *Varying c , these CMC hypersurfaces foliate a neighborhood of z_* .*
- (iii) *The hypersurfaces corresponding to non-degenerate maximum of R are asymptotically stable and those corresponding to saddle points and minima are unstable.*

Let $S_z^n := \{\omega \in T_z M : g_z(\omega, \omega) = 1\}$, the unit sphere in the tangent space $T_z M$, $I_z : \mathbb{S}^n \rightarrow S_z^n$ is an identification of S^n with \mathbb{S}^n and $\exp_z(v) : S_z^n \rightarrow M$ be the exponential map. For $\rho : \mathbb{S}^n \rightarrow \mathbb{R}_+$, we define the map

$$\theta_{\rho, z}(\omega) := \exp_z(\rho(\omega)I_z \omega). \tag{1.5}$$

Let $H^s \equiv H^s(\mathbb{S}^n, \mathbb{R})$. Consider the VPF (1.1) with initial configuration $\psi_0 = \theta_{\rho_0, z_0}$ close to the geodesic sphere, θ_{λ_*, z_*} , for some $z_* \in M^{n+1}$ and $\lambda_* > 0$, small, in the sense that $\|\rho_0 - \lambda_*\|_{H^k} + d_M(z_0, z_*)$, $k > n/2 + 1$, is sufficiently small, see Sect. 2.1. We compare the hypersurface $\psi(t, \cdot)$ with the *adiabatically* evolving hypersurface $\psi_{\lambda(t), z(t)} := \Phi_{\lambda(t)}(z(t))$ for some $z = z(t)$ and $\lambda = \lambda(t)$ to be determined. Let $a_\lambda(z) := A(\Phi_\lambda(z))$ and $v_{\text{enc}}(z) := V_{\text{enc}}(\Phi_\lambda(z))$ and

$$\chi_c = \{z \in M : v_{\text{enc}}(z) = c\}. \tag{1.6}$$

Statement (iii) is derived from the following (cf. [1])

Theorem 1.4 *With the definitions above and for c (or λ) sufficiently small, we have*

$$\psi(t, \omega) = \theta_{\xi(t), \psi_{\lambda(t), z(t)}(\omega)}(\omega), \tag{1.7}$$

$$\dot{z} = -\nabla a_\lambda(z) + O(\lambda^2 e^{-\delta t}), \tag{1.8}$$

$$\dot{\lambda} = O(\lambda^2 e^{-\delta t}), \tag{1.9}$$

where $z \in \chi_c$, $\delta = \frac{n+2}{2} - O(\lambda)$ and $\xi(t)$ satisfies the estimates

$$\|\xi\|_{H^k} \lesssim \lambda e^{-\delta t}. \tag{1.10}$$

We call evolution (1.8)–(1.9) the *adiabatic dynamics* of almost CMC surfaces. Eq (1.7) formulates the *nonlinear perturbation theory* (with $\xi(t)$ being a perturbation of $\psi_{\lambda(t), z(t)}$).

Foliations by CMC surfaces are of interest in general relativity (see e.g. [22]). For material sciences, the opposite scaling regime is of interest. In it, the ambient manifold is almost flat while the CMC surfaces are of size $O(1)$.

In the Euclidean case, $M = \mathbb{R}^{n+1}$, Huisken [21], in the general case, and Gage [17], for curves, proved that the solution to (1.1) exists globally and converges exponentially fast to

a sphere, provided that the initial surface S_0 is uniformly convex and smooth. Athanassenas [4, 5] has shown neckpinching of certain class of rotationally symmetric surfaces under the volume preserving modification of the mean curvature flow. Later Escher and Simonett [16] and Antonopoulou et al. [3] proved the asymptotic stability of spheres (see also [20]). For related works, see [2, 6] and the references therein.

Finally, we present some standard relations (see e.g. [13]) which are helpful in understanding our results and which are used in the proofs.

Proposition 1.5 *The Gateaux derivatives of the area and enclosed volume of $S \hookrightarrow M$ equal*

$$dA(S)\eta = \int_S Hf d\sigma, \quad dV_{\text{enc}}(S)\eta = \int_S f d\sigma, \tag{1.11}$$

where $f := g(\eta, \nu(S))$.

Hence, from the variation formula (1.11) it follows that the tangent space to X_c is

$$T_\psi X_c^k = \{ \xi : S^n \rightarrow TM, \int_{S^n} g(\xi, \nu(\psi)) d\sigma = 0 \} \tag{1.12}$$

Proposition 1.6 (i) *Minimizers of the area functional $A(\psi)$ for a given enclosed volume $V_{\text{enc}}(\psi)$ are critical points of $A(\psi)$ on X_c^k .*

(ii) *The Euler–Lagrange equation for these critical points is exactly the CMC equation (1.3).*

(iii) *These critical points are critical points of the modified functional*

$$A_h(\psi) = A(\psi) - hV_{\text{enc}}(\psi), \tag{1.13}$$

where h is determined by $c = V_{\text{enc}}(\psi)$ and vice versa.

(iv) *The VPF (1.1) is a gradient flow for the area functional on closed surfaces with given enclosed volume.*

Notation In this paper we use the following notation:

- $A \lesssim B$ denotes an inequality of the form $A \leq CB$, where $C > 0$ is a uniform in λ and z constant.
- $\partial_z^\alpha := \prod_{i=1}^{n+1} \partial_{z_i}$, where $z = (z_1, \dots, z_{n+1}) \in \mathbb{R}^{n+1}$ and $\alpha = (\alpha_1, \dots, \alpha_{n+1})$, $|\alpha| = \sum_{i=1}^{n+1} \alpha_i$.
- $S_z^n := \{ \omega \in T_z M : g_z(\omega, \omega) = 1 \}$, the unit sphere in the tangent space $T_z M$ and $H^s \equiv H^s(S_z^n, \mathbb{R})$, the Sobolev space of the order s .

2 The Lyapunov–Schmidt Map: Proof of Theorem 1.1

2.1 Graphs over (Geodesic) Spheres

We construct CMC hypersurfaces by deforming geodesic spheres $\theta_{\lambda,z}$ of radius λ centred at z . The geodesic spheres are the images of spheres, $S_z^n := \{ \omega \in T_z M : g_z(\omega, \omega) = 1 \}$, in the tangent space $T_z M$ under the exponential map $\exp_z(v)$ and can be parametrized by $\theta_{\lambda,z} : S_z^n \rightarrow M$, where

$$\theta_{\lambda,z}(\omega) = \exp_z(\lambda\omega), \quad z \in \mathbb{R}^{n+1}, \quad \omega \in S_z^n \tag{2.1}$$

As usual, $\exp_z(0) = z$ and $\partial_\lambda \exp_z(0) = \omega$. Abusing notation, we define the graph over the sphere S_z^n (or the corresponding geodesic sphere) as (cf. (1.5))

$$\theta_{\rho,z}(\omega) = \exp_z(\rho(\omega)\omega), \quad \rho(\omega) = \lambda(1 + \phi(\omega)) \tag{2.2}$$

We introduce the topology on the space of graphs over S_z^n as follows. We say the graph $\psi' = \theta_{\rho',z}$ is close to $\psi = \theta_{\rho,z}$ iff it can be written as $\psi'(\omega) = \exp_{\psi(\omega)}(\xi(\omega)\omega)$, with ξ sufficiently small in an appropriate norm (H^k).

2.2 The Lyapunov–Schmidt Map

Let dH denote the Gâteaux derivative of a map H ($dH(\psi)\eta := \partial_s H(\psi_s)|_{s=0}$ for ψ_s with $\psi_{s=0} = \psi$ and $\partial_s \psi_s|_{s=0} = \eta$) and let

$$\hat{H}(\theta) = H(\theta) - \bar{H}(\theta) \tag{2.3}$$

Lemma 2.1 *Let $\rho(\omega) = \lambda(1 + \phi(\omega))$. Then the Gâteaux derivative, $L_{0,z} := d\phi(\lambda\hat{H}(\theta_{\rho,z}))|_{\phi=0,\lambda=0}$, of \hat{H} at $\theta_{\lambda,z}$ equals*

$$L_{0,z}u = -(\Delta_{S^n} + n)u + \frac{n}{|S^n|} \int_{S^n} u \tag{2.4}$$

Moreover, $L_{0,z}$ is self-adjoint on $L^2(S^n)$, its spectrum is discrete and non-negative and its kernel is spanned by constants and the coordinate functions ω^i for $\omega = (\omega^1, \dots, \omega^{n+1}) \in S^n$.

Proof The formula (2.4) can be read off from the expansion (A.3) of the mean curvature $H(\theta_{\rho,z})$, included in Appendix A. It also follows from Proposition 2.5 of [9].

The spectral properties of $L_{0,z}$ are part of the standard theory. □

We will look for a solution of (1.3) of the form (2.2) with

$$\rho(\omega) = \lambda(1 + \phi(\omega)), \quad \phi \perp \omega^i, \quad i = 0, \dots, n + 1, \quad \omega^0 := 1. \tag{2.5}$$

The conditions on ϕ imply $\int_{S^n} \phi = 0$ and therefore

$$\lambda = \langle \rho \rangle = \frac{\int_{S^n} \rho}{|S^n|}.$$

(By Lemma B.2, with $\phi_{\lambda,z} = 0$, any $\psi = \theta_{\rho,z}$ sufficiently close to a geodesics sphere is of the form (2.2), with (2.5).)

Definition 2.2 Let P_0 be the $L^2(S^n)$ -orthogonal projection onto $\text{Null } L_{0,z}$, which is the span of the eigenvectors $1, \omega^1, \dots, \omega^{n+1}$ of $-\Delta_{S^n} - n$, with the eigenvalues $-n, 0$, and let $\bar{P}_0 := 1 - P_0$, the orthogonal projection onto $(\text{Null } L_{0,z})^\perp$.

Consider the equation for $\phi \in \bar{P}_0 H^k, k > \frac{n}{2} + 2$,

$$F(\phi, \lambda, z) = 0, \quad \text{where } F(\phi, \lambda, z) := \bar{P}_0 \lambda \hat{H}(\theta_{\lambda(1+\phi),z}) \tag{2.6}$$

Remark 2.3 We observe that the function $\lambda \hat{H}(\theta_{\lambda(1+\phi),z})$ has by definition zero mean, i.e., it is L^2 -orthogonal to 1. Hence, we need only project it onto the L^2 -orthogonal complement of the span of $\omega^1, \dots, \omega^{n+1}$, but we will keep using the projection operator \bar{P}_0 for consistency with other parts of the proof.

Proposition 2.4 For λ sufficiently small, Eq. (2.6) has a unique solution $\phi = \phi_{\lambda,z}$ in a small ball in $\bar{P}_0 H^k$, for $k > \frac{n}{2} + 2$, which satisfies the estimate

$$\|\partial_\lambda^r \partial_z^\alpha \phi_{\lambda,z}\|_{H^k} \lesssim \lambda^{2-r}, \quad r + |\alpha| \leq 2 \tag{2.7}$$

Proof The proof is an application of the inverse function theorem. We have the following properties:

- (i) The map $F : \bar{P}_0 H^k \times \mathbb{R}_{\geq 0} \times \mathbb{R}^n \rightarrow \bar{P}_0 H^{k-2}$ is smooth;
- (ii) $F(0, 0, z) = 0$,
- (iii) $d_\phi F(0, 0, z)$ is invertible.

(i) and (ii) are standard and follow from Lemma A.1 of Appendix A.

For (iii) we mention that, since $L_{0,z}$ is self-adjoint and has purely discrete spectrum and $\text{Null} L_0 = \text{Range}(1 - \bar{P}_0)$, the operator $d_\phi F(0, 0, z) = \bar{P}_0 L_{0,z}$ is invertible on $\text{Ran } \bar{P}_0$. Hence, by the IFT there exists a unique solution ϕ to (2.6) for every λ sufficiently small.

To prove the estimates (2.7), we use the expansion

$$\lambda \hat{H}(\theta_{\lambda(1+\phi),z}) = \lambda \hat{H}(\theta_{\lambda,z}) + L_{\lambda,z} \phi + N_{\lambda,z}(\phi), \tag{2.8}$$

where $L_{\lambda,z} := d_\phi(\lambda \hat{H}(\theta_{\rho,z}))|_{\phi=0}$ and $N_{\lambda,z}(\phi)$ is defined by this expression.

Let $\bar{L}_{\lambda,z} = \bar{P}_0 L_{\lambda,z} \bar{P}_0$. We would like to use the above expansion and invert the operator $\bar{L}_{\lambda,z}$ and rewrite equation $F(\phi, \lambda, z) = 0$ as the fixed point problem, which we use to estimate ϕ . This is done with the help of the following lemma.

Lemma 2.5 For λ sufficiently small, the operator $\bar{L}_{\lambda,z}$ is invertible and we have the following estimates

$$\|\partial_\lambda^m \partial_z^\alpha \bar{L}_{\lambda,z}^{-1} \phi\|_{H^k} \lesssim \|\phi\|_{H^{k-2}}, \tag{2.9}$$

$$\|\partial_\lambda^m \partial_z^\alpha F(0, \lambda, z)\|_{H^k} \lesssim \lambda^{2-m}, \tag{2.10}$$

$$\|\partial_\lambda^m \partial_z^\alpha N_{\lambda,z}(\phi)\|_{H^{k-2}} \lesssim \|\phi\|_{H^k}^2 \tag{2.11}$$

and a similar estimate for $N_{\lambda,z}(\phi') - N_{\lambda,z}(\phi)$, for $m + |\alpha| \leq 2$.

Proof We write

$$L_{\lambda,z} = L_{0,z} + \lambda^2 M_{\lambda,z}, \tag{2.12}$$

where $M_{\lambda,z}$ is defined by this expression. By Lemma A.1, we have that

$$\|(L_{0,z} + 1)^{-1} M_{\lambda,z}\|_{H^k \rightarrow H^k} \lesssim 1. \tag{2.13}$$

By (2.13), $\|\bar{L}_{0,z}^{-1} \lambda^2 \bar{M}_{\lambda,z}\|_{H^k \rightarrow H^k} < 1$ for λ sufficiently small and therefore $\bar{L}_{\lambda,z}$ is invertible.

Next, let $\beta := (m, \alpha)$, $w := (\lambda, z)$ and $\partial_w^\beta := \partial_\lambda^m \partial_z^\alpha$. Using the relation $\partial L^{-1} = -L^{-1} \partial L L^{-1}$, we see that we have that

$$\|\partial_w^\beta \bar{L}_{\lambda,z}^{-1}\|_{H^{k-2} \rightarrow H^k} = \sum \prod \|\partial_w^{\beta_i} (\lambda^2 \bar{L}_{\lambda,z}) \bar{L}_{\lambda,z}^{-1}\|_{H^k \rightarrow H^k} \|\bar{L}_{\lambda,z}^{-1}\|_{H^{k-2} \rightarrow H^k}^{r_i}$$

where the sum is taken over all partitions of $\beta := (m, \alpha)$ into smaller integers and the product over the integers in the given partition which give all the ways of distributing the partial ∂_w^β according to $\partial L^{-1} = -L^{-1} \partial L L^{-1}$. This gives (2.9).

Now, bounds (2.10) and (2.11) follow by comparing the expansion (2.8) with (A.3) of Appendix A. (2.10), follows from (A.3) with $\phi = 0$ and (2.11), from (A.6). \square

Now, using (2.8), we can rewrite Eq. (2.6) as the fixed point problem

$$\phi = -L_{\lambda,z}^{-1}(F(0, \lambda, z) + \bar{P}_0 N_{\lambda,z}(\phi)). \tag{2.14}$$

Lemma 2.5 implies that this equation has a unique fixed point, $\phi = \phi_{\lambda,z}$, in a small ball in $\bar{P}_0 H^k$. Assuming that ϕ lies in a small ball in $\bar{P}_0 H^k$ of radius $\varepsilon > 0$, we see that the bounds in (2.7) follow from (2.14) and Lemma 2.5. Indeed, we derive

$$\begin{aligned} \|\phi\|_{H^k} &\leq \|\bar{L}_{\lambda,z}^{-1}(F(0, \lambda, z))\|_{H^k} + \|\bar{L}_{\lambda,z}^{-1}\bar{P}_0 N_{\lambda,z}(\phi)\|_{H^k} \\ &\lesssim \lambda^2 + \|N_{\lambda,z}(\phi)\|_{H^{k-2}} \\ &\lesssim \lambda^2 + \|\phi\|_{H^k}^2, \end{aligned} \tag{2.15}$$

which implies estimate (2.7), with $r = 0, \alpha = 0$, provided $\|\phi\|_{H^k}$ is sufficiently small. The estimates for the $\partial_\lambda^r \partial_z^\alpha$ derivatives of ϕ are obtained similarly using the fixed point equation and estimates in Lemma 2.5. Indeed, running through the same argument as Eq. (2.15), gives (2.7). \square

Define the *Lyapunov–Schmidt map*, $\Phi_\lambda(z)$, as

$$\Phi_\lambda(z) = \theta_{\rho_{\lambda,z},z}, \text{ where } \rho_{\lambda,z} := \lambda(1 + \phi_{\lambda,z}), \quad \phi_{\lambda,z} \in \bar{P}_0 H^k \text{ solves (2.6)}. \tag{2.16}$$

Note that $\Phi_\lambda(z)$ is a perturbation of the classical exponential map.

2.3 Proof of Theorem 1.1

We begin with part (i). Let $\psi_{\lambda,z} := \Phi_\lambda(z)$. If $\psi_{\lambda,z}$, with $\lambda = \lambda(h)$, solves (1.3), then, by Proposition 1.6, $dA_h(\psi_{\lambda,z}) = 0$, for $h = h(\lambda)$. Furthermore,

$$\partial_{z^i} A_h(\psi_{\lambda,z}) = dA_h(\psi_{\lambda,z}) \partial_{z^i} \psi_{\lambda,z} \tag{2.17}$$

for every i . Hence, $\partial_{z^i} A_h(\psi_{\lambda,z}) = 0$, i.e. z is a critical point of $A_h(\psi_{\lambda,z})$.

Conversely, if $\partial_{z^i} A_h(\psi_{\lambda,z}) = 0$, we will argue that $dA_h(\psi_{\lambda,z}) = 0$. Define

$$\partial_{z^i}^N \psi_{\lambda,z} := g(\partial_{z^i} \psi_{\lambda,z}, \nu(\psi_{\lambda,z})). \tag{2.18}$$

Note that the superscript N in $\partial_{z^i}^N \psi_{\lambda,z}$ depends on $\psi_{\lambda,z}$.

In the case of the geodesic spheres we have $\partial_{z^i} \theta_{\lambda,z}|_{\lambda=0} = \partial_{z^i} \exp_z(\lambda\omega)|_{\lambda=0} = d \exp_z(0)(\partial_{z^i} z) = e_i$, where $\{e_i\}$ is the canonical basis in \mathbb{R}^{n+1} . Therefore $\partial_{z^i}^N \theta_{\lambda,z}|_{\lambda=0} := g(e_i, \omega) = \omega_i, i = 1, \dots, n + 1$, which span the null space of $L_{0,z}$ and P_0 can be written as

$$P_0 f = g_0^{ij} \langle f, \partial_{z^i}^N \theta_{\lambda,z}|_{\lambda=0} \rangle_{L^2} \partial_{z^j}^N \theta_{\lambda,z}|_{\lambda=0}, \tag{2.19}$$

where g_0^{ij} is the inverse matrix of $(g_0)_{ij} := \langle \partial_{z^i}^N \theta_{\lambda,z}|_{\lambda=0}, \partial_{z^j}^N \theta_{\lambda,z}|_{\lambda=0} \rangle_{L^2} = (\int_{\mathbb{S}^n} \omega_1^2) \delta_{ij}$.

We define the normal L^2 gradient, $\text{grad}^N G(\psi)$, of a functional $G(\psi)$ by

$$\int_{\mathbb{S}^n} \text{grad}^N G(\psi) \xi = dG(\psi) \xi,$$

for any normal variation ξ . Then $\text{grad}^N A_h(\psi) = H(\psi) - h$. By definition (2.16) and (2.6), $\bar{P}_0 \text{grad}^N A_h(\psi_{\lambda,z}) = 0$. Hence

$$P_0 \text{grad}^N A_h(\psi_{\lambda,z}) = \text{grad}^N A_h(\psi_{\lambda,z}) \tag{2.20}$$

Remembering (2.18), we define the projection on the span of the non L^2 -orthogonal set $\{\partial_{z_i}^N \psi_{\lambda,z}\}$:

$$P_\lambda f := g_\lambda^{ij} \langle \partial_{z_i}^N \psi_{\lambda,z}, f \rangle_{L^2} \partial_{z_j}^N \psi_{\lambda,z}, \tag{2.21}$$

where g_λ^{ij} is the inverse matrix of $(g_\lambda)_{ij} := \langle \partial_{z_i}^N \psi_{\lambda,z}, \partial_{z_j}^N \psi_{\lambda,z} \rangle_{L^2}$. (For simplicity, we do not display the dependence on z .) Hence, by the assumption $\partial_{z_i} A_h(\psi_{\lambda,z}) = 0$, we have

$$\begin{aligned} P_\lambda \text{grad}^N A_h(\psi_{\lambda,z}) &= g_\lambda^{ij} \langle \partial_{z_i}^N \psi_{\lambda,z}, \text{grad}^N A_h(\psi_{\lambda,z}) \rangle_{L^2} \partial_{z_j}^N \psi_{\lambda,z} \\ &= g_\lambda^{ij} \partial_{z_i}^N A_h(z) \partial_{z_j}^N \psi_{\lambda,z} = 0. \end{aligned} \tag{2.22}$$

Lemma 2.6 *The difference of the projections P_0, P_λ satisfies the estimate*

$$\|P_0 - P_\lambda\|_{L^2 \rightarrow L^2} \lesssim \lambda(1 + \|\phi_{\lambda,k}\|_{H^k} + \lambda \|\partial_\lambda \phi_{\lambda,z}\|_{H^k}) \tag{2.23}$$

for $k > n/2 + 1$.

Proof We show that the eigenfunctions of two projections are close. Let $f(\lambda) := \partial_{z_i}^N \theta_{\rho_{\lambda,z},z}$, with $\rho_{\lambda,z} := \lambda(1 + \phi_{\lambda,z})$. For ω fixed, we note that $f(\lambda)(\omega)$ is smooth in λ , $\phi_{\lambda,z}(\omega)$, and $\partial_z \phi_{\lambda,z}(\omega)$. Notice that $\partial_{z_i} \theta_{\rho_{\lambda,z},z} |_{\lambda=0} = \partial_{z_i} \theta_{\lambda,z} |_{\lambda=0}$. Then

$$\begin{aligned} \|\partial_{z_i}^N \theta_{\lambda,z} |_{\lambda=0} - \partial_{z_i}^N \theta_{\rho_{\lambda,z},z}\|_{H^{k-1}} &= \|f(\lambda) - f(0)\|_{H^{k-1}} \lesssim \lambda \int_0^1 \|f'(t\lambda)\|_{H^{k-1}} dt \\ &\lesssim \lambda(1 + \|\phi_{\lambda,k}\|_{H^k} + \lambda \|\partial_\lambda \phi_{\lambda,z}\|_{H^k}) \end{aligned}$$

The last inequality and definition (2.21) imply (2.23). □

Combining now (2.20), (2.22) and (2.23) and using

$$\text{grad}^N A_h(\psi_{\lambda,z}) = P_0 \text{grad}^N A_h(\psi_{\lambda,z}) = (P_0 - P_\lambda) \text{grad}^N A_h(\psi_{\lambda,z}),$$

we derive

$$\begin{aligned} \|\text{grad}^N A_h(\psi_{\lambda,z})\|_{L^2} &\leq \|P_0 - P_\lambda\|_{L^2} \cdot \|\text{grad}^N A_h(\psi_{\lambda,z})\|_{L^2} \\ &\lesssim \lambda(1 + \|\phi_{\lambda,k}\|_{H^k} + \lambda \|\partial_\lambda \phi_{\lambda,z}\|_{H^k}) \|\text{grad}^N A_h(\psi_{\lambda,z})\|_{L^2} \end{aligned}$$

Thus, for λ sufficiently small it follows that $\text{grad}^N A_h(\psi_{\lambda,z}) = 0$ and therefore $dA_h(\psi_{\lambda,z})\xi = 0$ for any normal variation ξ . Since $dA_h(\psi_{\lambda,z})\xi = 0$ for any tangential variation ξ , we have $dA_h(\psi_{\lambda,z}) = 0$.

Finally, we show that for λ sufficiently small, the map $\lambda \rightarrow h(\lambda) := H(\Phi_\lambda(z))$ is invertible. Using (2.8) and an expansion of $H(\theta_{\lambda,z})$ in λ (see also (A.3)), we find that

$$\lambda h = n + O(\lambda^2 + \|\phi\|_{H^k}) \tag{2.24}$$

$$= n + O(\lambda^2) \tag{2.25}$$

This and a similar estimate for the derivative in λ , together with the IFT, show that $\lambda \rightarrow h(\lambda)$ is invertible.

To prove part (ii) we let $A''_h(\psi) := d \text{grad} A_h(\psi)$ (the hessian of $A_h(\psi)$) and use (2.17) to compute the second derivative of $A_h(\Phi_\lambda(z))$:

$$\partial_{z_i} \partial_{z_j} A_h(\Phi_\lambda(z)) = \langle A''_h(\Phi_\lambda(z)) \partial_{z_j} \Phi_\lambda(z), \partial_{z_i} \Phi_\lambda(z) \rangle_{L^2} + \langle \text{grad} A_h(\Phi_\lambda(z)), \partial_{z_i} \partial_{z_j} \Phi_\lambda(z) \rangle_{L^2} \tag{2.26}$$

By (i), evaluating (2.26) at a critical point z_* of $A_h(\Phi_\lambda(z))$ we have

$$\partial_{z_i} \partial_{z_j} A_h(\Phi_\lambda(z_*)) = (A''_h(\Phi_\lambda(z_*)) \partial_{z_j} \Phi_\lambda(z_*), \partial_{z_i} \Phi_\lambda(z_*))_{L^2} \tag{2.27}$$

As before the one direction is obvious: if $A''_h(\Phi_\lambda(z_*))$ is positive, then so is $\partial_z^2 A_h(\Phi_\lambda(z_*))$.

We show that $\partial_z^2 A_h(\Phi_\lambda(z_*)) \geq 0 (> 0)$ implies $A''_h(\Phi_\lambda(z_*)) \geq 0 (> 0)$. Abbreviate $A''_h \equiv A''_h(\Phi_\lambda(z_*))$. By Proposition 1.5, $A''_h = d_\psi \hat{H}(\psi)|_{\psi=\Phi_\lambda(z_*)}$ and therefore by the definition $L_{\lambda,z} := d_\phi(\lambda \hat{H}(\theta_{\rho,z}))|_{\phi=0}$, we have that $\lambda^2 A''_h = L_{\lambda,z_*} + O(\lambda^3)$ and by relation (2.12),

$$\lambda^2 A''_h = L_{0,z_*} + \lambda^2 M_{\lambda,z_*} + O(\lambda^3) \tag{2.28}$$

This relation and $\bar{P}_0 L_{0,z} \bar{P}_0 \gtrsim \bar{P}_0$ give $\bar{P}_0 A''_h \bar{P}_0 \gtrsim \bar{P}_0$. This and (2.23) imply $\bar{P}_\lambda A''_h \bar{P}_\lambda \gtrsim \bar{P}_\lambda$.

Due to (2.12) and (2.23), the cross-terms $\bar{P}_\lambda A''_h \bar{P}_\lambda$ and $P_\lambda A''_h \bar{P}_\lambda$ are $O(\lambda)$. These estimates can be improved as follows. We recall from (2.20) that $\bar{P}_0 \text{grad}^N A_h(\psi_{\lambda,z}) = 0$ for all z . Hence

$$\bar{P}_0 A''_h P_\lambda = 0. \tag{2.29}$$

Using this identity, we see that

$$\bar{P}_\lambda A''_h P_\lambda = (\bar{P}_\lambda - \bar{P}_0) A''_h P_\lambda = (P_0 - P_\lambda) A''_h P_\lambda = (P_0 - P_\lambda) P_\lambda A''_h P_\lambda + (P_0 - P_\lambda) \bar{P}_\lambda A''_h P_\lambda.$$

Due to (2.23), this equation can be solved for $\bar{P}_\lambda A''_h P_\lambda$, for λ sufficiently small, as $\bar{P}_\lambda A''_h P_\lambda = S P_\lambda A''_h P_\lambda$, where $S = [1 - (P_0 - P_\lambda)]^{-1} (P_0 - P_\lambda)$, which gives

$$\bar{P}_\lambda A''_h P_\lambda = \bar{P}_\lambda S P_\lambda A''_h P_\lambda =: \bar{P}_\lambda S B. \tag{2.30}$$

The definitions (2.21) and $\psi_{\lambda,z} := \Phi_\lambda(z)$ imply

$$\langle \varphi, P_\lambda A''_h P_\lambda \varphi \rangle = \sum \bar{v}^i \langle \partial_{z_i}^N \psi_{\lambda,z}, A''_h \partial_{z_k}^N \psi_{\lambda,z} \rangle v^k,$$

where $v^i := \sum_j g^{ij} \langle \partial_{z_j}^N \psi_{\lambda,z}, \varphi \rangle$. The latter relation at $z = z_*$, equality (2.27) (since the tangential derivatives are zero modes of A''_h one can replace $\partial_{z_i} \psi_{\lambda,z}$ in (2.27) by $\partial_{z_i}^N \psi_{\lambda,z}$) and the assumption that $\partial_z^2 A_h(\Phi_\lambda(z_*)) \geq 0 (> 0)$ imply that $B = P_\lambda A''_h P_\lambda \geq 0 (> 0)$. Using this, (2.30) and that $\|S\| \lesssim \lambda$, we find

$$|\langle \varphi, \bar{P}_\lambda S B \varphi \rangle| \leq \|B^{1/2} S \bar{P}_\lambda \varphi\| \|B^{1/2} \varphi\| \leq \lambda C \|B^{1/2} Q\| \|\bar{P}_\lambda \varphi\| \|B^{1/2} \varphi\|,$$

for some constant $C \lesssim 1$, where Q be the finite rank orthogonal projection onto $\text{Ran} P_0 + \text{Rran} P_\lambda$. This gives

$$2|\langle \varphi, \bar{P}_\lambda A''_h P_\lambda \varphi \rangle| \leq 2\lambda C \|\bar{P}_\lambda \varphi\| \|B^{1/2} \varphi\| \leq \lambda^{2(1-\epsilon)} C \|\bar{P}_\lambda \varphi\|^2 + \lambda^{2\epsilon} \|B^{1/2} \varphi\|^2$$

which implies that

$$\begin{aligned} \langle \varphi, A''_h \varphi \rangle &= \langle \varphi, B \varphi \rangle + \langle \varphi, \bar{P}_\lambda A''_h \bar{P}_\lambda \varphi \rangle + 2 \text{Re} \langle \varphi, \bar{P}_\lambda A''_h P_\lambda \varphi \rangle \\ &\geq \langle \varphi, B \varphi \rangle + c \|\bar{P}_\lambda \varphi\|^2 - \lambda^{2(1-\epsilon)} C \|\bar{P}_\lambda \varphi\|^2 - \lambda^{2\epsilon} \|B^{1/2} \varphi\|^2 \end{aligned}$$

for some $c \gtrsim 1$. Hence $\langle \varphi, A''_h \varphi \rangle \geq 0 (> 0)$ provided λ is sufficiently small. This proves the assertion and with it part (ii).

Finally, we prove part (iii). By the definition, the unstable part of statement (iii) follows directly from statement (ii) and a standard argument. To prove the stability of minimizing CMC's, we use Theorem 1.4 as follows.

Proof of (iii)(stability) By (1.9), $\lambda \rightarrow \lambda_*$ for some λ_* as $t \rightarrow \infty$. If z_* is a strict local minimum point of $A(\Phi_\lambda(z))$ on $\{V_{\text{enc}}(\Phi_\lambda(z)) = c\}$, then $z(t) \rightarrow z_*$ and therefore, by (1.7) and (1.10), $\psi(t) = \theta_{\xi(t), \psi_{\lambda(t), z(t)}} \rightarrow \theta_{0, \psi_{\lambda_*, z_*}} = \psi_{\lambda_*, z_*}$. This shows that the surface ψ_{λ, z_*} is asymptotically stable.

If z_* is a saddle or local maximum point, then the standard argument and (1.8) show that $z(t)$ stays away from z_* however close the initial condition z_0 is to z_* . This implies the same for $\psi(t)$ w.r.to ψ_{λ_*, z_*} . □

Note that if z_* is a non-degenerate local minimum point of $A(\Phi_\lambda(z))$ on $\{V_{\text{enc}}(\Phi_\lambda(z)) = c\}$, then $z(t) \rightarrow z_*$ and therefore $\psi(t) \rightarrow \psi_{\lambda_*, z_*}$ exponentially fast. Indeed, the proof follows from (1.8) in a standard way. Indeed, let a''_h denote the hessian of a_h . Passing to local coordinates, one expands

$$\nabla a_h(z) = a''_h(z_*)(z - z_*) + O(|z - z_*|^2) \tag{2.31}$$

around z_* and uses the Duhamel principle to rewrite (1.8) as the fixed point problem $\beta = \Psi(\beta)$, where $\beta(t) = z(t) - z_*$ and

$$\Psi(\beta)(t) := e^{-a''_h(z_*)t} \beta(0) + \int_0^{+\infty} e^{-a''_h(z_*)(t-s)} [O(\beta^2(s)) + O(e^{-\delta s})] ds. \tag{2.32}$$

If $a''_h(z_*) \geq \alpha > 0$, then the standard fixed point argument in the space $X := C([0, +\infty), e^{-\alpha t} \mathbb{R}^{n+1})$ gives $\sup_t (e^{\alpha t} \|\beta(t)\|) \lesssim 1$ and therefore $|z(t) - z_*| \lesssim e^{-\alpha t}$, which implies that $\psi(t) = \theta_{\xi(t), \psi_{\lambda(t), z(t)}} \rightarrow \theta_{0, \psi_{\lambda_*, z_*}} = \psi_{\lambda_*, z_*}$ exponentially fast, giving the result.

3 Adiabatic Dynamics: Proof of Theorem 1.4

Proof of Theorem 1.4 Let $\psi(\omega) : \mathbb{S}^n \rightarrow M$ be an immersion of the form $\psi(\omega) = \theta_{\rho, z'}(\omega) := \exp_{z'}(\rho(\omega)\omega)$ for some ρ and z' . We assume ψ close to some CMC surface $\psi_{\lambda', z'} := \Phi_{\lambda'(1+\phi_{\lambda', z'})}$ (in the sense of topology induced by ρ and z' , see Sect. 2.1). We look for ρ in the form

$$\rho \equiv \rho(\xi) = \lambda(1 + \phi_{\lambda, z} + \xi), \tag{3.1}$$

for some λ and z close to λ' and z' . We note that ξ depends on λ, z , but we drop this dependence in our notation.

We denote $z^0 := \lambda$ and define

$$\sigma(\xi)(\omega) := g(v(\theta_{\rho, z}(\omega)), \partial_s \theta_{s, z}(\omega) |_{s=\rho}), \tag{3.2}$$

$$f_i(\xi)(\omega) := g(v(\theta_{\rho, z}(\omega)), \partial_{z_i} \theta_{\rho, z}(\omega)), \quad i = 0, 1, \dots, n + 1. \tag{3.3}$$

For $\xi = 0$, we omit the argument 0, e.g. we denote $f_i(0) = f_i$, etc.

Having in mind that f_i and σ depend on λ and z , we define the *barycenter* of $\psi = \theta_{\rho, z'}$ to be (λ, z) that solve the equation

$$\langle f_i, \sigma \xi \rangle = 0 \text{ for } i = 0, \dots, n + 1. \tag{3.4}$$

By Lemma B.2, a unique solution exists provided ψ is sufficiently near some CMC surface $\psi_{\lambda', z'}$.

Suppose that $\psi(\omega, t) = \theta_{\rho(t), z(t)}(\omega)$ is a solution to VPF with some initial condition $\psi_0 = \theta_{\rho_0, z_0}$ near a CMC surface $\psi_{\lambda', z'}$ and let $\lambda(t), z(t)$ be its barycenter. Then we see that

$$\partial_t^N \psi = f_i(\xi) \dot{z}^i + \lambda \sigma(\xi) \dot{\xi}. \tag{3.5}$$

Our first task is to obtain the effective equation for λ and z . We begin with rewriting (1.1) as

$$\partial_t^N \psi = -\nabla^{N(\psi)} A(\psi)$$

on X_c^k and expand the r.h.s. in ξ and divide the resulting equation by $(\sigma(\xi)/\sigma) = 1 + O_{H^k}(\xi)$. Using (3.5), the definition $\psi_{\lambda,z} := \Phi_\lambda(z) = \theta_{\lambda(1+\phi_{\lambda,z}),z}$ and Lemma B.1, we can rewrite our equation $\partial_t^N \psi = -\nabla^{N(\psi)} A(\psi)$ on X_c^k as

$$\tilde{f}_j(\xi) \dot{z}^j + \lambda \sigma \dot{\xi} = -\nabla^{N(\psi_{\lambda,z})} A(\psi_{\lambda,z}) + \lambda L_{\lambda,z} \xi + \lambda^2 O_{H^k}(\xi^2) + \lambda O_{H^k}(|\dot{\xi}| \xi), \tag{3.6}$$

for $z \in \chi_c$, where $\tilde{f}_j(\xi) := f_j(\xi)\sigma/\sigma(\xi) = f_j + O_{H^k}(\xi)$ and $L_{\lambda,z} := \lambda d_\phi \nabla^{N(\theta_{\rho,z})} A(\theta_{\rho,z})|_{\rho=\rho_{\lambda,z}}$ ($\rho = \lambda(1 + \phi)$). Now we multiply the equation by f_i , $i = 0, \dots, n + 1$, and divide the resulting equation by $(h(\xi)/h) = 1 + O_{H^k}(\xi)$, where the matrices h and $h(\xi)$ have the entries $h_{ij} = \langle f_i, f_j \rangle$ and $h_{ij}(\xi) = \langle f_i, \tilde{f}_j(\xi) \rangle$, with $i, j = 0, \dots, n + 1$. We have

$$h_{ij} \dot{z}^j = -\langle f_i, \nabla^{N(\psi_{\lambda,z})} A(\psi_{\lambda,z}) \rangle - \lambda \langle f_i, \sigma \dot{\xi} \rangle - \lambda \langle f_i, L_{\lambda,z} \xi \rangle + \lambda^2 O_{H^k}(\xi^2) + \lambda O_{H^k}(|\dot{\xi}| \xi) + O_{H^k}(|\dot{z}| \xi). \tag{3.7}$$

By Lemma B.1, we see that $h_{ij} \approx \text{id}$ is invertible. Moreover, by the choice of the barycenter, we see that

$$\langle f_i, \sigma \dot{\xi} \rangle = -\langle \partial_t(f_i \sigma), \xi \rangle = O_{H^k}(|\dot{z}| \xi). \tag{3.8}$$

Set $v_i = \langle f_i, \nabla^{N(\psi_{\lambda,z})} A(\psi_{\lambda,z}) \rangle$. Since $\partial_{z^j} A(\psi_{\lambda,z}) = dA(\psi_{\lambda,z}) \partial_{z^j} \psi_{\lambda,z} = \int g(Hv, \partial_{z^j} \psi_{\lambda,z}) = \int H f_j + O_{H^k}(\xi)$ since $H = O(\lambda^{-1})$ (cf. Lemma A.1), we have $v_j = \partial_{z^j} v$. This and the fact that f_j are almost the null eigenvectors of the linear part $L_{\lambda,z}$ imply (after solving for \dot{z})

$$\dot{z} = -h^{ij} v_j + \lambda^2 O_{H^k}(\xi) = -\nabla v + \lambda^2 O_{H^k}(\xi) \tag{3.9}$$

where $-\nabla v = -h^{ij} \partial_{z^j} v$ for $z \in \chi_c$.

Now to get equation for ξ , let P denote the orthogonal projection onto $\text{Span}\{f_i\}$ and let $\bar{P} = \mathbf{1} - P$. Applying \bar{P} to (3.6) and using $\bar{P} f_i = 0$, we derive

$$\lambda \bar{P}(\sigma \dot{\xi}) = -\bar{P} \nabla^{N(\psi_{\lambda,z})} A(\psi_{\lambda,z}) - \lambda L_{\lambda,z} \xi - \bar{P}(\tilde{f}_i(\xi) - f_i) \dot{z}^i + \lambda^2 O_{H^k}(\xi^2) \tag{3.10}$$

By the choice of barycenter again (see (3.4)),

$$\bar{P}(\sigma \dot{\xi}) = \sigma \dot{\xi} - P(\sigma \dot{\xi}) = \sigma \dot{\xi} + \partial_t(P\sigma) \xi \tag{3.11}$$

So we get

$$\dot{\xi} = -\sigma^{-1} L_{\lambda,z} \xi + \text{Rem}, \tag{3.12}$$

$$\text{Rem} := -\lambda^{-1} \sigma^{-1} (\bar{P} \nabla^{N(\psi)} A(\psi) + \lambda \partial_t(P\sigma) \xi + \bar{P}(f_i(\xi) - f_i) \dot{z}^i + \lambda^2 O_{H^k}(\xi^2)) \tag{3.13}$$

Now, using equations (3.9) and (3.12), we estimate ξ . To this end, following [3], we use the Lyapunov functionals:

$$\Lambda_k(\xi) := \frac{1}{2} \langle \xi, (-\Delta - n)^k \xi \rangle. \tag{3.14}$$

These functionals satisfy the inequalities

$$\Lambda_k(\xi) \geq (n + 2) \Lambda_{k-1}(\xi), \quad C \|\xi\|_{H^k}^2 \geq \Lambda_k(\xi) \geq c \|\xi\|_{H^k}^2 \tag{3.15}$$

for some $C, c > 0$. As in [3], one uses the equation on ξ , (3.12), to obtain differential inequalities for $\Lambda_k(u)$, which together with (3.15) imply the bound (1.10) completing the proof of the theorem. \square

4 Proof of Theorem 1.3: Existence and Linear Stability/Instability

Theorem 1.3(iii) follows readily from Theorem 1.1(iii). Thus we address only the first two statements.

4.1 Existence of Foliations

By Theorem 1.1(i), to prove existence of the CMC's, we have to prove existence of critical points of the function $A_h(\Phi_\lambda(a)) := A(\Phi_\lambda(a)) - hV_{enc}(\Phi_\lambda(a))$. To this end, we use the following expansion which follows from Corollary A.2 of Appendix A (see [24], Lemma 2.2):

$$A_h(\Phi_\lambda(z)) = \lambda^n \left(\frac{a_n}{n+1} \left[1 + n - h\lambda + \left(\frac{h\lambda}{6(n+3)} - \frac{1}{6} \right) R(z)\lambda^2 \right] + \lambda^3 B_{\lambda,z}^{(0)}(\phi) + \lambda^2 B_{\lambda,z}^{(2)}(\phi) \right) \tag{4.1}$$

where a_n denotes the area of the Euclidean unit sphere S^n , R is the scalar curvature of M and the remainders $B_{\lambda,z}^{(j)}(\phi)$, $j = 0, 2$, satisfy the estimates

$$|\partial_\lambda^j \partial_z^\alpha B_{\lambda,z}^{(r)}(\phi)| \lesssim \sum_{j \leq i, \beta \leq \alpha} \|\partial_\lambda^j \partial_z^\alpha \phi\|_{H^k}^r \tag{4.2}$$

provided $\|\phi\|_{H^k} \lesssim 1$ and $i + |\alpha| \leq 2$. (If ϕ is independent of λ and z , then only the term with $j = |\beta| = 0$ survives on the r.h.s.)

Utilizing the estimate (2.7) and (4.1), the equations $\partial_{z_j} A_h(\Phi_\lambda(z)) = 0$, $j = 1, \dots, n + 1$ take the form

$$\nabla_z R(z) + O(\lambda^2) = 0 \tag{4.3}$$

If λ is sufficiently small, the preceding equation can be solved for z near a non-degenerate critical point z_* of R by employing the implicit function theorem. The latter yields a solution z satisfying

$$z = z_* + O(\lambda^2) \tag{4.4}$$

Since $\Phi_\lambda(z)$ solves (2.6) and $\partial_z A_h(\Phi_\lambda(z)) = 0$, by Theorem 1.1 (i), it is also a solution of (1.3).

Equation (1.3) and expansion (A.3) imply the relation between h and λ :

$$h = \frac{n}{\lambda} + O(\lambda). \tag{4.5}$$

Now we show that the solutions $\Phi_\lambda(z_*)$ actually foliate the neighborhood of z_* . We exhibit a local homeomorphism from a neighborhood of z_* to in \mathbb{R}^{n+1} that maps each CMC surface into a spherical shell centred at the origin. We simply use the map (2.16):

$$\begin{aligned} \Phi : [0, \epsilon) \times S^n &\rightarrow M, \\ (\lambda, \omega) &\mapsto \Phi_\lambda(z_*). \end{aligned}$$

where, recall, $\Phi_\lambda(z) := \exp_z(\rho_{\lambda,z}\omega)$, $\rho_{\lambda,z} := \lambda(1 + \phi_{\lambda,z})$. We note that $(r, \omega) \mapsto \exp_{z_*}(r\omega)$ is a local diffeomorphism, so it suffices to show that the map $(\lambda, \omega) \mapsto (\rho_{\lambda,z_*}, \omega)$ is a

homeomorphism. Indeed, taking the Jacobian matrix, we get an upper diagonal matrix

$$\begin{pmatrix} \partial_\lambda \rho_{\lambda, z_*} & \cdots \\ 0 & \text{id}_{n \times n} \end{pmatrix}.$$

Since $\|\partial_\lambda \rho_{\lambda, z_*} - 1\|_\infty = \|\phi_{\lambda, z_*} + \lambda \partial_\lambda \phi_{\lambda, z_*}\|_\infty \lesssim \lambda^2$ by Proposition 2.4, we see that the Jacobian matrix is invertible. It follows by the inverse function theorem that Φ is a homeomorphism.

The linear stability/instability statement of the theorem follows from the formula (4.1), the asymptotics (4.5) and Theorem 1.1(ii).

Acknowledgements Ilias Chenn and I. M. Sigal research is supported in part by NSERC Grant No. NA7901. Ilias Chenn is also partly supported by NSERC CGS D Scholarship. G. Fournodavlos is supported by the EPSRC Grant EP/K00865X/1 on ‘‘Singularities of Geometric Partial Differential Equations’’. We are grateful to the anonymous referee for useful remarks. I. M. Sigal is grateful to Stephen Gustafson for many illuminating discussions and insights. The author’s discussions over a period of many years with Jürg Fröhlich, Tom Spencer and Herbert Spohn played a crucial role in forming his understanding of adiabatic and effective dynamics and of mathematical physics in general.

Appendix A: Expansions of the Induced Metric and Mean Curvature

We Taylor expand the induced metric and mean curvature of $\theta_{\rho, z}$ and estimate the non-linear terms in ϕ in H^k , $k > \frac{n}{2} + 2$. One may consult [18] for a thorough study on such classical expansions, we mostly follow the notation in [24]. We abuse slightly notation by denoting the pushforward of ω through $d\theta_{\rho, z}$ by the same symbol ω . For instance, given x^i, ∂_i a set of coordinates and the associated vector fields, we write the corresponding basis of tangent vector fields on S_z^n : $\zeta_i := \lambda(1 + \phi)\partial_i\omega + \lambda\partial_i\phi\omega \in TS_z^n$.

Lemma A.1 (see [24], Lemmas 2.1 and 2.4) *The following expansions are valid:*

$$\begin{aligned} \lambda^{-2}(1 + \phi)^{-2}g(\zeta_i, \zeta_j) &= g(\partial_i\omega, \partial_j\omega) + \frac{1}{3}R(\omega, \partial_i\omega, \omega, \partial_j\omega) \cdot \lambda^2(1 + \phi)^2 \\ &\quad + \frac{\partial_i\phi\partial_j\phi}{(1 + \phi)^2} + \lambda^3R_{\lambda, z}^{(0)}(\phi) + \lambda^2R_{\lambda, z}^{(2)}(\phi) \end{aligned} \tag{A.1}$$

where R is the Riemann curvature tensor and the remainders $R_{\lambda, z}^{(j)}(\phi)$, $j = 0, 2$, are local terms depending on ϕ and $\partial\phi$ and satisfying the estimates

$$\|\partial_\lambda^i \partial_z^\alpha R_{\lambda, z}^{(r)}(\phi, \partial\phi)\|_{H^{k-1}} \lesssim \sum_{j \leq i, \beta \leq \alpha} \|\partial_\lambda^j \partial_z^\beta \phi\|_{H^k}^r \tag{A.2}$$

provided $\|\phi\|_{H^k} \lesssim 1$ and $i + |\alpha| \leq 2$, and

$$\lambda H(\theta_{\rho, z}) = n - \frac{1}{3}Ric_z \lambda^2 - (\Delta_{S^n} + n)\phi + \lambda^2 M'_{\lambda, z} \phi + \lambda^2 N'_{\lambda, z}(\phi) + \lambda^3 H_{\lambda, z}, \tag{A.3}$$

where $Ric_z : \omega \rightarrow Ric_z(\omega, \omega)$ ($Ric(\cdot, \cdot)$ is the Ricci curvature tensor of M), $M'_{\lambda, z} \phi$, $N'_{\lambda, z}(\phi)$ and $H_{\lambda, z}$ are linear non-linear terms in ϕ and in its derivatives up to order two, respectively,

and independent of ϕ term, satisfying the estimates:

$$\|\partial_\lambda^i \partial_z^\alpha M'_{\lambda,z}(\phi)\|_{H^{k-2}} \lesssim \sum_{j \leq i, \beta \leq \alpha} \|\partial_\lambda^j \partial_z^\beta \phi\|_{H^k}, \tag{A.4}$$

$$\|\partial_\lambda^i \partial_z^\alpha N'_{\lambda,z}(\phi)\|_{H^{k-2}} \lesssim \sum_{j \leq i, \beta \leq \alpha} \|\partial_\lambda^j \partial_z^\beta \phi\|_{H^k}^2, \tag{A.5}$$

$$\|\partial_\lambda^j \partial_z^\alpha H_{\lambda,z}\|_{H^{k-2}} \lesssim 1, \tag{A.6}$$

and similarly for $N'_{\lambda,z}(\phi') - N'_{\lambda,z}(\phi)$, for $i + |\alpha| \leq 2, k > \frac{n}{2} + 2$. (Above, if ϕ is independent of λ and z , then only the term with $j = |\beta| = 0$ survives on the r.h.s..)

Proof Both expressions (A.1) and (A.3) can be read from [24]. Bounds (A.4) are immediate by definition. In order to derive estimate (A.6) for the non-linearity $N_{\lambda,z}(\phi)$ we have to first examine its structure. Recall that the mean curvature of $\theta_{\rho,z}$ is given by⁴

$$H(\theta_{\rho,z}) = \sum_{i,j} g^{ij} g(\nabla_{\zeta_i} \nu, \zeta_j) = \operatorname{div}_{\theta_{\rho,z}} \nu \tag{A.8}$$

where g^{ij} is the inverse matrix to the metric $g_{ij} := g(\zeta_i, \zeta_j)$ and ν the outward unit normal vector field on $\theta_{\rho,z}$:

$$\nu = \frac{-\omega + \lambda \sum_{i,j} g^{ij} \partial_i \phi \zeta_j}{\sqrt{1 - \lambda^2 g(\nabla_{\mathbb{S}^n} \phi, \nabla_{\mathbb{S}^n} \phi)}} \tag{A.9}$$

Note that ν is well defined for λ and $\|\phi\|_{H^k}$ appropriately small. Hence, we observe that the non-linear terms in ϕ arising in the expansion of $H(\theta_{\rho,z})$ are of the form $b(\lambda\phi(\omega), \lambda\partial\phi(\omega))\lambda\partial^2\phi(\omega)$, where $b(s, t)$ is a simple function, uniformly bounded together with its derivatives, provided $|s| \ll 1$ and $|t| \ll 1$. Using these estimates it is not hard (but somewhat tedious) to show that $\|b(\lambda\phi, \lambda\partial\phi)\lambda\partial^2\phi\|_{H^{k-2}} \lesssim \lambda^r \|\phi\|_{H^k}^r$ for some $r \geq 2$, provided $\|\phi\|_{H^k} \ll 1$ (here we use the condition $k > \frac{n}{2} + 2$). This completes the proof of the lemma. \square

Eq (A.1) implies

$$\begin{aligned} \lambda^{-n} (1 + \phi)^{-n} \sqrt{\det g_{\theta_{\rho,z}}} &= 1 + \frac{1}{6} \operatorname{Ric}(\omega, \omega) \lambda^2 (1 + \phi)^2 + \frac{1}{2} \frac{|\nabla_{\mathbb{S}^n} \phi|^2}{(1 + \phi)^2} \\ &\quad + \lambda^3 S_{\lambda,z}^{(0)}(\phi) + \lambda^2 S_{\lambda,z}^{(2)}(\phi) \end{aligned} \tag{A.10}$$

where Ric is the Ricci curvature tensor of M and $S_{\lambda,z}^{(r)}, r = 0, 2$, are local terms depending on ϕ and $\partial\phi$ satisfying the estimates

$$\|\partial_\lambda^i \partial_z^\alpha S_{\lambda,z}^{(r)}(\phi, \partial\phi)\|_{H^{k-1}} \lesssim \|\phi\|_{H^k}^r \tag{A.11}$$

⁴ Let g be the ambient metric with its associated Christoffel symbols, Γ_{kl}^j , and \bar{g} be the pull back metric of g onto $\psi(\mathbb{S}^n)$. In any local coordinate, if ψ is any immersion from \mathbb{S}^n to M^{n+1} ,

$$H = \bar{g}^{\alpha\beta} g_{ij} \nu^i (\partial_\alpha \partial_\beta \psi^j + \Gamma_{kl}^j \partial_\alpha \psi^i \partial_\beta \psi^k) \tag{A.7}$$

where the Latin indices $i, j = 1, \dots, n + 1$ are for coordinates in the ambient manifold and the Greek ones $1, \dots, n$ are on \mathbb{S}^n and ν^j is the unit normal vector on $\psi(\mathbb{S}^n)$. At $\psi(\omega)$, we see that all geometric quantities are smooth functions of the ambient metric and at most 2 derivatives of the immersion ψ , all evaluated at ω . We remark that, however, dependence on top (2nd) order derivative is linear and comes from the $\nabla_\alpha \nabla_\beta \psi$ term above.

provided $\|\phi\|_{H^k} \lesssim 1$ and $i + |\alpha| \leq 2$. Furthermore, multiplying (A.10) by $\lambda^n(1 + \phi)^n$, integrating over \mathbb{S}^n and taking into account that $\int_{\mathbb{S}^n} \phi = 0$, and using the co-area formula and polar coordinates, we obtain:

Corollary A.2 *We have the following expansions*

$$A(\Phi_\lambda(z)) = \lambda^n \left(a_n \left[1 - \frac{1}{6(n+1)} R(z)\lambda^2 \right] + \lambda^3 Q_{\lambda,z}^{(0)}(\phi) + \lambda^2 Q_{\lambda,z}^{(2)}(\phi) \right) \tag{A.12}$$

where a_n denotes the area of the Euclidean unit sphere \mathbb{S}^n and R is the scalar curvature of M , and

$$V_{enc}(\Phi_\lambda(z)) = \lambda^{n+1} \left(\frac{a_n}{n+1} \left[1 - \frac{1}{6(n+3)} R(z)\lambda^2 \right] + \lambda^3 T_{\lambda,z}^{(0)}(\phi) + \lambda^2 T_{\lambda,z}^{(2)}(\phi) \right) \tag{A.13}$$

The remainders $Q_{\lambda,z}^{(r)}$ and $T_{\lambda,z}^{(r)}$ above are local expression in $\phi, \partial\phi$ satisfying the estimates of the type of (4.2).

Appendix B: Existence of Barycenter

In this section, we show that barycenter exists. That is, we show that Eq. (3.4) has a solution. To begin, we state some useful estimates.

Lemma B.1 *For the terms defined in equations (3.2)–(3.3), we have the estimate*

$$\|\sigma - 1\|_{H^{k-1}} \lesssim \lambda \|\phi_{\lambda,z}\|_{H^k} \tag{B.1}$$

$$\|\sigma(\xi) - \sigma\|_{H^{k-1}} \lesssim \lambda \|\xi\|_{H^k} \tag{B.2}$$

$$\|f_i - \omega_i\|_{H^{k-1}} \lesssim \lambda(1 + \|\phi_{\lambda,z}\|_{H^k} + \|\partial_{z_i}\phi_{\lambda,z}\|_{H^k}) \text{ for } i = 1, \dots, n + 1 \tag{B.3}$$

$$\|f_i(\xi) - f_i\|_{H^{k-1}} \lesssim \lambda \|\xi\|_{H^k} \text{ for } i = 1, \dots, n + 1 \tag{B.4}$$

$$\|f_0 - 1\|_{H^{k-1}} \lesssim \lambda \|\partial_\lambda \phi_{\lambda,z}\|_{H^k} + \|\phi_{\lambda,z}\|_{H^k} \tag{B.5}$$

$$\|f_i(\xi) - f_i\|_{H^{k-1}} \lesssim \lambda \|\xi\|_{H^k} \tag{B.6}$$

Proof Recall that $\psi(\omega) = \exp_z(\rho(\xi)\omega)$, where $\rho(\xi) := \lambda(1 + \phi_{\lambda,z} + \xi)$. Using the definitions (3.2), we compute the difference, $\sigma(\xi) - \sigma$. To this end, it suffices to compute the differences of each factor in (3.2):

$$\begin{aligned} \sigma(\xi) - \sigma &= g_\psi(v(\theta_{\rho(\xi),z}(\omega)), \partial_s \theta_{s,z}(\omega) |_{s=\rho(\xi)}) - g_{\theta_{\lambda,z}}(v(\theta_{\rho(0),z}(\omega)), \partial_s \theta_{s,z}(\omega) |_{s=\rho(0)}) \\ &= g_\psi(v(\theta_{\rho(\xi),z}(\omega)), \partial_s \theta_{s,z}(\omega) |_{s=\rho(\xi)}) - g_\psi(Tv(\theta_{\rho(0),z}(\omega)), T\partial_s \theta_{s,z}(\omega) |_{s=\rho(0)}) \end{aligned}$$

where T is the parallel transport from $\theta_{\lambda,z}(\omega)$ to $\psi(\omega)$. Continuing the estimate, we have

$$\begin{aligned} &= g_\psi(v(\theta_{\rho(\xi)}) - Tv(\theta_{\rho(0),z}(\omega)), \partial_s \theta_{s,z}(\omega) |_{s=\rho(\xi)}) \\ &\quad + g_\psi(Tv(\theta_{\rho(0),z}(\omega)), \partial_s \theta_{s,z}(\omega) |_{s=\rho(\xi)} - T\partial_s \theta_{s,z}(\omega) |_{s=\rho(0)}). \end{aligned}$$

Recalling that $\theta_{\lambda,z} = \exp_z(\rho(0)\omega)$ and letting $\omega^0 := 1$, we find furthermore

$$\begin{aligned} g(v(\theta_{\rho(0),z}(\omega)), \partial_s \theta_{s,z}(\omega) |_{s=\rho(0)}) |_{\lambda=0} &= g(v(\theta_{\lambda,z}(\omega)) |_{\lambda=0}, \partial_s \theta_{s,z}(\omega) |_{s=0}) = 1 \\ g(v(\theta_{\rho(0),z}(\omega)), \partial_{z_i} \theta_{\rho(0),z}(\omega)) |_{\lambda=0} &= \omega^i, \quad i = 0, 1, \dots, n + 1. \end{aligned}$$

Similarly, we write $f_i(\xi) - f_i$. Hence, using this, we only need to estimate the following items to complete the proof:

$$\begin{aligned}
 &g_p - g_{p'} \\
 &T(s) - T(s') \\
 &\partial_s \exp_z(s\omega) |_{s=s_1} - T(s_1 - s_2, \exp_z(s_2\omega))\partial_s \exp_z(s\omega) |_{s=s_2} \\
 &\partial_z \exp_z(s_1\omega) - T(s_1 - s_2, \exp_z(s_2\omega))\partial_z \exp_z(s_2\omega) \\
 &\rho(\xi) - \rho(0) \\
 &\partial_{\lambda}\rho - \partial_{\lambda}\rho |_{\phi_{\lambda,z}=0} \\
 &\rho(\xi) \\
 &\nu(\psi) - T(\lambda, \theta_{\lambda,z})\nu(\theta_{\lambda,z})
 \end{aligned}$$

where g_p is the value of the metric at p , $T(s, z)$ is parallel transport from z along ω for time s . Then the difference of quantities in the statement of the question are composition and smooth functions of the above with at most two derivatives of $\phi_{\lambda,z}$ and ξ . Note that the first four quantities only depends on the ambient geometry of M^{n+1} . Since we are working locally, we may assume that we are working on a compact subset of M . Thus, the first four are smooth functions, the first four quantity exhibits Lipschitz estimates in the difference in their argument. For example,

$$|\partial_z \exp_z(s_1\omega) - T(s_1 - s_2, \exp_z(s_2\omega))\partial_z \exp_z(s_2\omega)| \lesssim |s_1 - s_2|$$

uniformly on M^{n+1} . The next 3 expressions have the obvious estimate

$$\begin{aligned}
 \|\rho(\xi) - \rho(0)\|_{H^k} &= \lambda\|\xi\|_{H^k} \\
 \|\partial_{\lambda}\rho - \partial_{\lambda}\rho |_{\phi_{\lambda,z}=0}\|_{H^k} &\lesssim \lambda\|\partial_{\lambda}\phi_{\lambda,z}\|_{H^k} + \|\phi_{\lambda,z}\|_{H^k} \\
 \|\partial_z\rho - \partial_z\rho |_{\phi_{\lambda,z}=0}\|_{H^k} &\lesssim \lambda(\|\partial_z\phi_{\lambda,z}\|_{H^k} + \|\phi_{\lambda,z}\|_{H^k}) \\
 \|\rho(\xi)\|_{H^k} &\lesssim \lambda(\|\xi\|_{H^k} + \|\phi_{\lambda,z}\|_{H^k}).
 \end{aligned}$$

Finally, the last one follows from the fact $\nu(\psi)$ is a none singular rational function of ψ . \square

Now, let

$$P(\lambda, z, \psi) : \mathbb{R} \times \mathbb{R}^{n+1} \times H^k \rightarrow \mathbb{R} \times \mathbb{R}^{n+1} = (\langle f_0\sigma, \xi \rangle, \dots, \langle f_{n+1}\sigma, \xi \rangle)$$

where ξ is defined by Eq. (3.1).

Lemma B.2 Any ψ sufficiently close to a CMC $\theta_{\lambda_0, z_0} = \exp_{z_0}(\lambda_0(1 + \phi_{\lambda_0, z_0}))$ can be written in the form (2.2), with (3.1) and λ and z satisfying the equation

$$P(\lambda, z, \psi) = 0$$

Proof We note that $P(\lambda_0, z_0, \theta_{\lambda_0, z_0}) = 0$. By the implicit function theorem, it suffices to show that

- (i) P is C^1 , and
- (ii) $(\partial_{\lambda,z}P)(\lambda_0, z_0, \theta_{\lambda_0, z_0})$ is an invertible matrix.

(i) We check, by definition of f_i and ξ , that they are real functions of λ, z and $\phi_{\lambda,z}$ up to 1 derivative. Since $\phi_{\lambda,z}$ is C^1 in λ and z (cf. Proposition 2.4), we see that f_i and σ are C^1 in λ and z . Since P is linear in ξ , using the estimate of Proposition 2.4 again, we see that P is C^1 in ψ as well.

(ii) We compute

$$(\partial_{\lambda,z} P)(\lambda_0, z_0, \theta_{\lambda_0,z_0}) = (\langle f_i \sigma, \partial_{\lambda,z} \xi \rangle) \tag{B.7}$$

since $\xi = 0$ for $\psi = \theta_{\lambda_0,z_0}$. To compute $\partial_\lambda \xi$, we use the fact that, by definition of ξ ,

$$\exp_{z_0}(\lambda(1 + \phi_{\lambda,z_0} + \xi)\omega) = \exp_{z_0}(\lambda_0(1 + \phi_{\lambda_0,z_0})\omega) \tag{B.8}$$

for λ sufficiently close to λ_0 . (Note that we suppressed the identification I_z between $T_z M$ and \mathbb{R}^{n+1} here as z is not varied.) Taking ∂_λ and evaluate at $\lambda = 0$, we get the result of

$$\partial_s \exp_{z_0}(s\omega) |_{s=\lambda_0(1+\phi_{\lambda_0,z_0})} (1 + \phi_{\lambda_0,z_0} + \lambda_0 \partial_\lambda \phi_{\lambda,z_0} |_{\lambda=\lambda_0} + \lambda_0 \partial_\lambda \xi |_{\lambda=\lambda_0}) = 0 \tag{B.9}$$

Contracting with $\nu(\theta_{\lambda_0,\theta_0})$, we get

$$0 = f_0 + \lambda_0 \sigma \partial_\lambda \xi, \tag{B.10}$$

where the last line follow from Proposition 2.4. To compute $\partial_z \xi |_{\lambda_0,z_0}$, we consider curves $z(t) : (-\epsilon, \epsilon) \rightarrow M$ with $z(0) = z_0$ and $\dot{z}(0) = v$ for any $v \in T_{z_0} M$ fixed. Then any variation of

$$\exp_{z(t)}(\lambda_0(1 + \phi_{\lambda_0,z(t)} + \xi)I_{z(t)}(\omega)) \tag{B.11}$$

is tangential. Taking derivative with respect to t and setting $t = 0$, we see that the expression (B.11) becomes

$$\begin{aligned} & [(\partial_x \exp_x(\lambda_0(1 + \phi_{\lambda_0,z_0})I_x(\omega))) |_{x=z_0} \\ & + \partial_s \exp_{z_0}(s\omega) |_{s=\lambda_0(1+\phi_{\lambda_0,z_0})} (\lambda_0 \partial_z \phi_{\lambda_0,z} |_{z=z_0} + \lambda_0 \partial_z \xi |_{z=z_0})] \dot{z}(0) \end{aligned}$$

Using the fact that varying z is only tangential, it follows that

$$0 = g(\nu(\theta_{\lambda_0,z_0}), [(\partial_x \exp_x(\lambda_0(1 + \phi_{\lambda_0,z_0})I_x(\omega))) |_{x=z_0} \tag{B.12}$$

$$+ \partial_s \exp_{z_0}(s\omega) |_{s=\lambda_0(1+\phi_{\lambda_0,z_0})} (\lambda_0 \partial_z \phi_{\lambda_0,z} |_{z=z_0})]v) \tag{B.13}$$

$$+ g(\nu(\theta_{\lambda_0,z_0}), \lambda_0 \partial_z \xi |_{z=z_0})v) \tag{B.14}$$

for any $v \in T_{z_0} M$. By definition of f_i ,

$$f_i + \lambda_0 \sigma \partial_{z^i} \xi = 0 \tag{B.15}$$

It follows that

$$(\partial_{\lambda,z} P)(\lambda_0, z_0, \theta_{\lambda_0,z_0}) = -\lambda_0^{-1} \langle f_i, f_j \rangle \tag{B.16}$$

is invertible by Lemma B.1. □

References

1. Alikakos, N., Freire, A.: The normalized mean curvature flow for a small bubble in a Riemannian manifold. *J. Differ. Geom.* **64**(2), 247–303 (2003)
2. Andrews, B.: Volume-preserving anisotropic mean curvature flow. *Indiana Univ. Math. J.* **50**(2), 783–827 (2001)
3. Antonopoulou, D.C., Karali, G.D., Sigal, I.M.: Stability of spheres under volume preserving mean curvature flow. *Dyn. PDE* **7**, 327–344 (2010)
4. Athanassenas, M.: Volume-preserving mean curvature flow of rotationally symmetric surfaces. *Comment. Math. Helv.* **72**(1), 52–66 (1997)

5. Athanassenas, M.: Behaviour of singularities of the rotationally symmetric, volume-preserving mean curvature flow. *Calc. Var. Partial Differ. Equ.* **17**(1), 1–16 (2003)
6. Athanassenas, M., Kandanaarachchi, S.: Convergence of axially symmetric volume-preserving mean curvature flow. *Pac. J. Math.* **259**(1), 41–54 (2012)
7. Bach, V., Fröhlich, J., Sigal, I.M.: Quantum electrodynamics of confined non-relativistic particles. *Adv. Math.* **137**, 299–395 (1998)
8. Bach, V., Chen, T., Fröhlich, J., Sigal, I.M.: Smooth Feshbach map and operator-theoretic renormalization group methods. *J. Funct. Anal.* **203**, 44–92 (2003)
9. Barbosa, J.L., do Carmo, M., Eschenburg, J.: Stability of hypersurfaces of constant mean curvature in Riemannian manifolds. *Math. Z.* **197**(1), 123–138 (1988)
10. Bourgain, J.: On a homogenization problem. Preprint (2017)
11. Bronsard, L., Stoth, B.: Volume-preserving mean curvature flow as a limit of a nonlocal Ginzburg–Landau equation. *SIAM J. Math. Anal.* **28**(4), 769–807 (1997)
12. Chen, X., Hilhorst, D., Logak, E.: Mass conserved Allen–Cahn equation and volume preserving mean curvature flow (2009). [arXiv:0902.3625v1](https://arxiv.org/abs/0902.3625v1)
13. Colding, T.H., Minicozzi II, W.P.: Minimal submanifolds. *Bull. Lond. Math. Soc.* **38**, 353 (2006)
14. Conlon, J.G., Spencer, T.: A strong central limit theorem for a class of random surfaces. *Commun. Math. Phys.* **325**(1), 1–15 (2014)
15. Conlon, J.G., Spencer, T.: Strong convergence to the homogenized limit of elliptic equations with random coefficients. *Trans. Am. Math. Soc.* **366**(3), 1257–1288 (2014)
16. Escher, J., Simonett, G.: The volume preserving mean curvature flow near spheres. *Proc. Am. Math. Soc.* **126**(9), 2789–2796 (1998)
17. Gage, M.: On an area-preserving evolution equation for plane curves. In: De Turck, D.M. (ed.), *Nonlinear Problems in Geometry*. Contemporary Mathematics, vol. 51. AMS, Providence, pp. 51–62 (1986)
18. Gray, A.: The volume of a small geodesic ball of a Riemannian manifold. *Mich. Math. J.* **20**, 329–344 (1973)
19. Griesemer, M., Hasler, D.: On the smooth Feshbach–Schur map. *J. Funct. Anal.* **254**(9), 2329–2335 (2008)
20. Hartley, D.: Motion by volume preserving mean curvature flow near cylinders. *Commun. Anal. Geom.* **21**(5), 873–889 (2013)
21. Huisken, G.: The volume preserving mean curvature flow. *J. Reine Angew. Math.* **382**, 35–48 (1987)
22. Huisken, G., Yau, S.T.: Definition of center of mass for isolated physical systems and unique foliation by stable spheres with constant mean curvature flow. *Invent. Math.* **124**, 281–311 (1996)
23. Matsen, M.W.: The standard Gaussian model for block copolymer melts. *J. Phys. Condens. Matter* **14**, R21–R47 (2002)
24. Pacard, F., Xu, X.: Constant mean curvature spheres in Riemannian manifolds. *Manuscr. Math.* **128**(3), 275–295 (2009)
25. Pizzo, A.: Bose particles in a box I. A convergent expansion of the ground state of a three-modes Bogoliubov Hamiltonian. [arXiv:1511.07022](https://arxiv.org/abs/1511.07022)
26. Pizzo, A.: Bose particles in a box II. A convergent expansion of the ground state of the Bogoliubov Hamiltonian in the mean field limiting regime. [arXiv:1511.07025](https://arxiv.org/abs/1511.07025)
27. Pizzo, A.: Bose particles in a box III. A convergent expansion of the ground state of the Hamiltonian in the mean field limiting regime. [arXiv:1511.07026](https://arxiv.org/abs/1511.07026)
28. Spohn, H.: Interface motion in models with stochastic dynamics. *J. Stat. Phys.* **71**(5/6), 1081 (1993)
29. Ye, R.: Foliation by constant mean curvature spheres. *Pac. J. Math.* **147**(2), 381–396 (1991)
30. Willmore, T.J.: *Riemannian Geometry*. Oxford University Press, Oxford (1993)