

# ELEMENTS BOOK 2

*Fundamentals of Geometric Algebra*


Ὅροι.

α'. Πᾶν παραλληλόγραμμον ὀρθογώνιον περιέχεσθαι λέγεται ὑπὸ δύο τῶν τὴν ὀρθὴν γωνίαν περιεχουσῶν εὐθειῶν.

β'. Παντὸς δὲ παραλληλογράμμου χωρίου τῶν περὶ τὴν διάμετρον αὐτοῦ παραλληλογράμμων ἐν ὁποιοοῦν σὺν τοῖς δυὶ παραπληρώμασι γνῶμων καλείσθω.

α'.

Ἐὰν ὄσι δύο εὐθεῖαι, τμηθῆ δὲ ἡ ἑτέρα αὐτῶν εἰς ὅσα δεητοῦν τμήματα, τὸ περιεχόμενον ὀρθογώνιον ὑπὸ τῶν δύο εὐθειῶν ἴσον ἐστὶ τοῖς ὑπὸ τε τῆς ἀτμήτου καὶ ἐκάστου τῶν τμημάτων περιεχομένοις ὀρθογωνίσις.



Ἐστωσαν δύο εὐθεῖαι αἱ  $A$ ,  $BΓ$ , καὶ τετμήσθω ἡ  $BΓ$ , ὡς ἔτυχεν, κατὰ τὰ  $Δ$ ,  $Ε$  σημεία· λέγω, ὅτι τὸ ὑπὸ τῶν  $A$ ,  $BΓ$  περιεχόμενον ὀρθογώνιον ἴσον ἐστὶ τῶν ὑπὸ τῶν  $A$ ,  $BΔ$  περιεχομένῳ ὀρθογωνίῳ καὶ τῶν ὑπὸ τῶν  $A$ ,  $ΔΕ$  καὶ ἔτι τῶν ὑπὸ τῶν  $A$ ,  $ΕΓ$ .

Ἦχθω γὰρ ἀπὸ τοῦ  $B$  τῆς  $BΓ$  πρὸς ὀρθὰς ἡ  $BZ$ , καὶ κείσθω τῆς  $A$  ἴση ἡ  $BH$ , καὶ διὰ μὲν τοῦ  $H$  τῆς  $BΓ$  παράλληλος ἡ  $HK$  ἡ  $HΘ$ , διὰ δὲ τῶν  $Δ$ ,  $Ε$ ,  $Γ$  τῆς  $BH$  παράλληλοι ἡ  $ΔK$ ,  $ΕΛ$ ,  $ΓΘ$ .

Ἴσον δὴ ἐστὶ τὸ  $BΘ$  τοῖς  $BK$ ,  $ΔΛ$ ,  $ΕΘ$ . καὶ ἐστὶ τὸ μὲν  $BΘ$  τὸ ὑπὸ τῶν  $A$ ,  $BΓ$ · περιέχεται μὲν γὰρ ὑπὸ τῶν  $HB$ ,  $BΓ$ , ἴση δὲ ἡ  $BH$  τῆς  $A$ · τὸ δὲ  $BK$  τὸ ὑπὸ τῶν  $A$ ,  $BΔ$ · περιέχεται μὲν γὰρ ὑπὸ τῶν  $HB$ ,  $BΔ$ , ἴση δὲ ἡ  $BH$  τῆς  $A$ . τὸ δὲ  $ΔΛ$  τὸ ὑπὸ τῶν  $A$ ,  $ΔΕ$ · ἴση γὰρ ἡ  $ΔK$ , τουτέστιν ἡ  $BH$ , τῆς  $A$ . καὶ ἔτι ὁμοίως τὸ  $ΕΘ$  τὸ ὑπὸ τῶν  $A$ ,  $ΕΓ$ · τὸ ἄρα ὑπὸ τῶν  $A$ ,  $BΓ$  ἴσον ἐστὶ τῶν ὑπὸ  $A$ ,  $BΔ$  καὶ τῶν ὑπὸ  $A$ ,  $ΔΕ$  καὶ ἔτι τῶν ὑπὸ  $A$ ,  $ΕΓ$ .

Ἐὰν ἄρα ὄσι δύο εὐθεῖαι, τμηθῆ δὲ ἡ ἑτέρα αὐτῶν εἰς ὅσα δεητοῦν τμήματα, τὸ περιεχόμενον ὀρθογώνιον ὑπὸ τῶν δύο εὐθειῶν ἴσον ἐστὶ τοῖς ὑπὸ τε τῆς ἀτμήτου καὶ ἐκάστου τῶν τμημάτων περιεχομένοις ὀρθογωνίσις· ὅπερ


Definitions

1. Any rectangular parallelogram is said to be contained by the two straight-lines containing the right-angle.

2. And in any parallelogrammic figure, let any one whatsoever of the parallelograms about its diagonal, (taken) with its two complements, be called a gnomon.

Proposition 1<sup>†</sup>

If there are two straight-lines, and one of them is cut into any number of pieces whatsoever, then the rectangle contained by the two straight-lines is equal to the (sum of the) rectangles contained by the uncut (straight-line), and every one of the pieces (of the cut straight-line).



Let  $A$  and  $BC$  be the two straight-lines, and let  $BC$  be cut, at random, at points  $D$  and  $E$ . I say that the rectangle contained by  $A$  and  $BC$  is equal to the rectangle(s) contained by  $A$  and  $BD$ , by  $A$  and  $DE$ , and, finally, by  $A$  and  $EC$ .

For let  $BF$  have been drawn from point  $B$ , at right-angles to  $BC$  [Prop. 1.11], and let  $BG$  be made equal to  $A$  [Prop. 1.3], and let  $GH$  have been drawn through (point)  $G$ , parallel to  $BC$  [Prop. 1.31], and let  $DK$ ,  $EL$ , and  $CH$  have been drawn through (points)  $D$ ,  $E$ , and  $C$  (respectively), parallel to  $BG$  [Prop. 1.31].

So the (rectangle)  $BH$  is equal to the (rectangles)  $BK$ ,  $DL$ , and  $EH$ . And  $BH$  is the (rectangle contained) by  $A$  and  $BC$ . For it is contained by  $GB$  and  $BC$ , and  $BG$  (is) equal to  $A$ . And  $BK$  (is) the (rectangle contained) by  $A$  and  $BD$ . For it is contained by  $GB$  and  $BD$ , and  $BG$  (is) equal to  $A$ . And  $DL$  (is) the (rectangle contained) by  $A$  and  $DE$ . For  $DK$ , that is to say  $BG$  [Prop. 1.34], (is) equal to  $A$ . Similarly,  $EH$  (is) also the (rectangle contained) by  $A$  and  $EC$ . Thus, the (rectangle contained) by  $A$  and  $BC$  is equal to the (rectangles contained) by  $A$

ἔδει δεῖξαι.


and  $BD$ , by  $A$  and  $DE$ , and, finally, by  $A$  and  $EC$ .

Thus, if there are two straight-lines, and one of them is cut into any number of pieces whatsoever, then the rectangle contained by the two straight-lines is equal to the (sum of the) rectangles contained by the uncut (straight-line), and every one of the pieces (of the cut straight-line). (Which is) the very thing it was required to show.

† This proposition is a geometric version of the algebraic identity:  $a(b + c + d + \dots) = ab + ac + ad + \dots$ .

β'.

Ἐάν εὐθεῖα γραμμὴ τμηθῆ, ὡς ἔτυχεν, τὸ ὑπὸ τῆς ὅλης καὶ ἑκατέρου τῶν τμημάτων περιεχόμενον ὀρθογώνιον ἴσον ἐστὶ τῷ ἀπὸ τῆς ὅλης τετραγώνῳ.



Εὐθεῖα γὰρ ἡ  $AB$  τετμήσθω, ὡς ἔτυχεν, κατὰ τὸ  $\Gamma$  σημεῖον· λέγω, ὅτι τὸ ὑπὸ τῶν  $AB$ ,  $B\Gamma$  περιεχόμενον ὀρθογώνιον μετὰ τοῦ ὑπὸ  $BA$ ,  $A\Gamma$  περιεχομένου ὀρθογωνίου ἴσον ἐστὶ τῷ ἀπὸ τῆς  $AB$  τετραγώνῳ.


Ἀναγεγράφθω γὰρ ἀπὸ τῆς  $AB$  τετράγωνον τὸ  $ADEB$ , καὶ ἦχθω διὰ τοῦ  $\Gamma$  ὀποτέρᾳ τῶν  $AD$ ,  $BE$  παράλληλος ἡ  $\Gamma Z$ .

ἴσον δὴ ἐστὶ τὸ  $AE$  τοῖς  $AZ$ ,  $GE$ . καὶ ἐστὶ τὸ μὲν  $AE$  τὸ ἀπὸ τῆς  $AB$  τετράγωνον, τὸ δὲ  $AZ$  τὸ ὑπὸ τῶν  $BA$ ,  $A\Gamma$  περιεχόμενον ὀρθογώνιον· περιέχεται μὲν γὰρ ὑπὸ τῶν  $DA$ ,  $A\Gamma$ , ἴση δὲ ἡ  $AD$  τῇ  $AB$ : τὸ δὲ  $GE$  τὸ ὑπὸ τῶν  $AB$ ,  $B\Gamma$ : ἴση γὰρ ἡ  $BE$  τῇ  $AB$ . τὸ ἄρα ὑπὸ τῶν  $BA$ ,  $A\Gamma$  μετὰ τοῦ ὑπὸ τῶν  $AB$ ,  $B\Gamma$  ἴσον ἐστὶ τῷ ἀπὸ τῆς  $AB$  τετραγώνῳ.

Ἐάν ἄρα εὐθεῖα γραμμὴ τμηθῆ, ὡς ἔτυχεν, τὸ ὑπὸ τῆς ὅλης καὶ ἑκατέρου τῶν τμημάτων περιεχόμενον ὀρθογώνιον ἴσον ἐστὶ τῷ ἀπὸ τῆς ὅλης τετραγώνῳ· ὅπερ ἔδει δεῖξαι.

Proposition 2†

If a straight-line is cut at random then the (sum of the) rectangle(s) contained by the whole (straight-line), and each of the pieces (of the straight-line), is equal to the square on the whole.



For let the straight-line  $AB$  have been cut, at random, at point  $C$ . I say that the rectangle contained by  $AB$  and  $BC$ , plus the rectangle contained by  $BA$  and  $AC$ , is equal to the square on  $AB$ .

For let the square  $ADEB$  have been described on  $AB$  [Prop. 1.46], and let  $CF$  have been drawn through  $C$ , parallel to either of  $AD$  or  $BE$  [Prop. 1.31].


So the (square)  $AE$  is equal to the (rectangles)  $AF$  and  $CE$ . And  $AE$  is the square on  $AB$ . And  $AF$  (is) the rectangle contained by the (straight-lines)  $BA$  and  $AC$ . For it is contained by  $DA$  and  $AC$ , and  $AD$  (is) equal to  $AB$ . And  $CE$  (is) the (rectangle contained) by  $AB$  and  $BC$ . For  $BE$  (is) equal to  $AB$ . Thus, the (rectangle contained) by  $BA$  and  $AC$ , plus the (rectangle contained) by  $AB$  and  $BC$ , is equal to the square on  $AB$ .

Thus, if a straight-line is cut at random then the (sum of the) rectangle(s) contained by the whole (straight-line), and each of the pieces (of the straight-line), is equal to the square on the whole. (Which is) the very thing it was required to show.

† This proposition is a geometric version of the algebraic identity:  $ab + ac = a^2$  if  $a = b + c$ .

γ'.

Ἐάν εὐθεῖα γραμμὴ τμηθῆ, ὡς ἔτυχεν, τὸ ὑπὸ τῆς ὅλης καὶ ἐνὸς τῶν τμημάτων περιεχόμενον ὀρθογώνιον ἴσον ἐστὶ τῷ τε ὑπὸ τῶν τμημάτων περιεχομένῳ ὀρθογώνιῳ καὶ τῷ ἀπὸ τοῦ προειρημένου τμήματος τετραγώνῳ.




Εὐθεῖα γὰρ ἡ  $AB$  τετμήσθω, ὡς ἔτυχεν, κατὰ τὸ  $\Gamma$ . λέγω, ὅτι τὸ ὑπὸ τῶν  $AB$ ,  $BE$  περιεχόμενον ὀρθογώνιον ἴσον ἐστὶ τῷ τε ὑπὸ τῶν  $AG$ ,  $GB$  περιεχομένῳ ὀρθογώνιῳ μετὰ τοῦ ἀπὸ τῆς  $BE$  τετραγώνου.

Ἀναγεγράφθω γὰρ ἀπὸ τῆς  $GB$  τετράγωνον τὸ  $\Gamma\Delta EB$ , καὶ διήχθω ἡ  $E\Delta$  ἐπὶ τὸ  $Z$ , καὶ διὰ τοῦ  $A$  ὁποτέρῳ τῶν  $\Gamma\Delta$ ,  $BE$  παράλληλος ἦχθω ἡ  $AZ$ . ἴσον δὲ ἐστὶ τὸ  $AE$  τοῖς  $A\Delta$ ,  $\Gamma E$ : καὶ ἐστὶ τὸ μὲν  $AE$  τὸ ὑπὸ τῶν  $AB$ ,  $BE$  περιεχόμενον ὀρθογώνιον: περιέχεται μὲν γὰρ ὑπὸ τῶν  $AB$ ,  $BE$ , ἴση δὲ ἡ  $BE$  τῇ  $BE$ : τὸ δὲ  $A\Delta$  τὸ ὑπὸ τῶν  $AG$ ,  $GB$ : ἴση γὰρ ἡ  $\Delta\Gamma$  τῇ  $GB$ : τὸ δὲ  $\Delta B$  τὸ ἀπὸ τῆς  $GB$  τετράγωνον: τὸ ἄρα ὑπὸ τῶν  $AB$ ,  $BE$  περιεχόμενον ὀρθογώνιον ἴσον ἐστὶ τῷ ὑπὸ τῶν  $AG$ ,  $GB$  περιεχομένῳ ὀρθογώνιῳ μετὰ τοῦ ἀπὸ τῆς  $BE$  τετραγώνου.

Ἐάν ἄρα εὐθεῖα γραμμὴ τμηθῆ, ὡς ἔτυχεν, τὸ ὑπὸ τῆς ὅλης καὶ ἐνὸς τῶν τμημάτων περιεχόμενον ὀρθογώνιον ἴσον ἐστὶ τῷ τε ὑπὸ τῶν τμημάτων περιεχομένῳ ὀρθογώνιῳ καὶ τῷ ἀπὸ τοῦ προειρημένου τμήματος τετραγώνῳ· ὅπερ ἔδει δεῖξαι.

Proposition 3†

If a straight-line is cut at random then the rectangle contained by the whole (straight-line), and one of the pieces (of the straight-line), is equal to the rectangle contained by (both of) the pieces, and the square on the aforementioned piece.



For let the straight-line  $AB$  have been cut, at random, at (point)  $C$ . I say that the rectangle contained by  $AB$  and  $BC$  is equal to the rectangle contained by  $AC$  and  $CB$ , plus the square on  $BC$ .

For let the square  $CDEB$  have been described on  $CB$  [Prop. 1.46], and let  $ED$  have been drawn through to  $F$ , and let  $AF$  have been drawn through  $A$ , parallel to either of  $CD$  or  $BE$  [Prop. 1.31]. So the (rectangle)  $AE$  is equal to the (rectangle)  $AD$  and the (square)  $CE$ . And  $AE$  is the rectangle contained by  $AB$  and  $BC$ . For it is contained by  $AB$  and  $BE$ , and  $BE$  (is) equal to  $BC$ . And  $AD$  (is) the (rectangle contained) by  $AC$  and  $CB$ . For  $DC$  (is) equal to  $CB$ . And  $DB$  (is) the square on  $CB$ . Thus, the rectangle contained by  $AB$  and  $BC$  is equal to the rectangle contained by  $AC$  and  $CB$ , plus the square on  $BC$ .

Thus, if a straight-line is cut at random then the rectangle contained by the whole (straight-line), and one of the pieces (of the straight-line), is equal to the rectangle contained by (both of) the pieces, and the square on the aforementioned piece. (Which is) the very thing it was required to show.

† This proposition is a geometric version of the algebraic identity:  $(a + b)a = ab + a^2$ .


δ'.

Ἐάν εὐθεῖα γραμμὴ τμηθῆ, ὡς ἔτυχεν, τὸ ἀπὸ τῆς ὅλης τετράγωνον ἴσον ἐστὶ τοῖς τε ἀπὸ τῶν τμημάτων τετραγώνοις καὶ τῷ δις ὑπὸ τῶν τμημάτων περιεχομένῳ ὀρθο-

Proposition 4†

If a straight-line is cut at random then the square on the whole (straight-line) is equal to the (sum of the) squares on the pieces (of the straight-line), and twice the

γωνίω.




Εὐθεΐα γὰρ γραμμὴ ἡ  $AB$  τετμήσθω, ὡς ἔτυχεν, κατὰ τὸ  $\Gamma$ . λέγω, ὅτι τὸ ἀπὸ τῆς  $AB$  τετράγωνον ἴσον ἐστὶ τοῖς τε ἀπὸ τῶν  $AG$ ,  $GB$  τετραγώνοις καὶ τῷ δις ὑπὸ τῶν  $AG$ ,  $GB$  περιεχομένῳ ὀρθογώνιῳ.

Ἀναγεγράφθω γὰρ ἀπὸ τῆς  $AB$  τετράγωνον τὸ  $ADEB$ , καὶ ἐπεζεύχθω ἡ  $BD$ , καὶ διὰ μὲν τοῦ  $\Gamma$  ὀποτέρᾳ τῶν  $AD$ ,  $EB$  παράλληλος ἦχθω ἡ  $\Gamma Z$ , διὰ δὲ τοῦ  $H$  ὀποτέρᾳ τῶν  $AB$ ,  $DE$  παράλληλος ἦχθω ἡ  $\Theta K$ . καὶ ἐπεὶ παράλληλός ἐστιν ἡ  $\Gamma Z$  τῇ  $AD$ , καὶ εἰς αὐτὰς ἐμπέπτωκεν ἡ  $BD$ , ἡ ἐκτὸς γωνία ἢ ὑπὸ  $\Gamma HB$  ἴση ἐστὶ τῇ ἐντὸς καὶ ἀπεναντίον τῇ ὑπὸ  $ADB$ . ἀλλ' ἡ ὑπὸ  $ADB$  τῇ ὑπὸ  $ABD$  ἐστὶν ἴση, ἐπεὶ καὶ πλευρὰ ἢ  $BA$  τῇ  $AD$  ἐστὶν ἴση· καὶ ἡ ὑπὸ  $\Gamma HB$  ἄρα γωνία τῇ ὑπὸ  $HBF$  ἐστὶν ἴση· ὥστε καὶ πλευρὰ ἢ  $BG$  πλευρᾷ τῇ  $\Gamma H$  ἐστὶν ἴση· ἀλλ' ἡ μὲν  $GB$  τῇ  $HK$  ἐστὶν ἴση. ἡ δὲ  $\Gamma H$  τῇ  $KB$ · καὶ ἡ  $HK$  ἄρα τῇ  $KB$  ἐστὶν ἴση· ἰσόπλευρον ἄρα ἐστὶ τὸ  $\Gamma HKB$ . λέγω δὴ, ὅτι καὶ ὀρθογώνιον. ἐπεὶ γὰρ παράλληλός ἐστιν ἡ  $\Gamma H$  τῇ  $BK$  [καὶ εἰς αὐτὰς ἐμπέπτωκεν εὐθεΐα ἡ  $GB$ ], αἱ ἄρα ὑπὸ  $KBF$ ,  $HGB$  γωνίαι δύο ὀρθαῖς εἰσὶν ἴσαι. ὀρθὴ δὲ ἡ ὑπὸ  $KBF$ · ὀρθὴ ἄρα καὶ ἡ ὑπὸ  $BGH$ · ὥστε καὶ αἱ ἀπεναντίον αἱ ὑπὸ  $\Gamma HK$ ,  $HKB$  ὀρθαῖς εἰσιν. ὀρθογώνιον ἄρα ἐστὶ τὸ  $\Gamma HKB$ · ἐδείχθη δὲ καὶ ἰσόπλευρον· τετράγωνον ἄρα ἐστὶν· καὶ ἐστὶν ἀπὸ τῆς  $GB$ . διὰ τὰ αὐτὰ δὴ καὶ τὸ  $\Theta Z$  τετράγωνόν ἐστιν· καὶ ἐστὶν ἀπὸ τῆς  $\Theta H$ , τουτέστιν [ἀπὸ] τῆς  $AG$ · τὰ ἄρα  $\Theta Z$ ,  $KF$  τετράγωνα ἀπὸ τῶν  $AG$ ,  $GB$  εἰσιν. καὶ ἐπεὶ ἴσον ἐστὶ τὸ  $AH$  τῷ  $HE$ , καὶ ἐστὶ τὸ  $AH$  τὸ ὑπὸ τῶν  $AG$ ,  $GB$ · ἴση γὰρ ἡ  $HG$  τῇ  $GB$ · καὶ τὸ  $HE$  ἄρα ἴσον ἐστὶ τῷ ὑπὸ  $AG$ ,  $GB$ · τὰ ἄρα  $AH$ ,  $HE$  ἴσα ἐστὶ τῷ δις ὑπὸ τῶν  $AG$ ,  $GB$ . ἔστι δὲ καὶ τὰ  $\Theta Z$ ,  $KF$  τετράγωνα ἀπὸ τῶν  $AG$ ,  $GB$ · τὰ ἄρα τέσσαρα τὰ  $\Theta Z$ ,  $KF$ ,  $AH$ ,  $HE$  ἴσα ἐστὶ τοῖς τε ἀπὸ τῶν  $AG$ ,  $GB$  τετραγώνοις καὶ τῷ δις ὑπὸ τῶν  $AG$ ,  $GB$  περιεχομένῳ ὀρθογώνιῳ. ἀλλὰ τὰ  $\Theta Z$ ,  $KF$ ,  $AH$ ,  $HE$  ὅλον ἐστὶ τὸ  $ADEB$ , ὃ ἐστὶν ἀπὸ τῆς  $AB$  τετράγωνον· τὸ ἄρα ἀπὸ τῆς  $AB$  τετράγωνον ἴσον ἐστὶ τοῖς τε ἀπὸ τῶν  $AG$ ,  $GB$  τετραγώνοις καὶ τῷ δις ὑπὸ τῶν  $AG$ ,  $GB$  περιεχομένῳ ὀρθογώνιῳ.

Ἐὰν ἄρα εὐθεΐα γραμμὴ τμηθῇ, ὡς ἔτυχεν, τὸ ἀπὸ τῆς

rectangle contained by the pieces.



For let the straight-line  $AB$  have been cut, at random, at (point)  $C$ . I say that the square on  $AB$  is equal to the (sum of the) squares on  $AC$  and  $CB$ , and twice the rectangle contained by  $AC$  and  $CB$ .

For let the square  $ADEB$  have been described on  $AB$  [Prop. 1.46], and let  $BD$  have been joined, and let  $CF$  have been drawn through  $C$ , parallel to either of  $AD$  or  $EB$  [Prop. 1.31], and let  $HK$  have been drawn through  $G$ , parallel to either of  $AB$  or  $DE$  [Prop. 1.31]. And since  $CF$  is parallel to  $AD$ , and  $BD$  has fallen across them, the external angle  $CGB$  is equal to the internal and opposite (angle)  $ADB$  [Prop. 1.29]. But,  $ADB$  is equal to  $ABD$ , since the side  $BA$  is also equal to  $AD$  [Prop. 1.5]. Thus, angle  $CGB$  is also equal to  $GBC$ . So the side  $BC$  is equal to the side  $CG$  [Prop. 1.6]. But,  $CB$  is equal to  $GK$ , and  $CG$  to  $KB$  [Prop. 1.34]. Thus,  $GK$  is also equal to  $KB$ . Thus,  $CGKB$  is equilateral. So I say that (it is) also right-angled. For since  $CG$  is parallel to  $BK$  [and the straight-line  $CB$  has fallen across them], the angles  $KBC$  and  $GCB$  are thus equal to two right-angles [Prop. 1.29]. But  $KBC$  (is) a right-angle. Thus,  $BCG$  (is) also a right-angle. So the opposite (angles)  $CGK$  and  $GKB$  are also right-angles [Prop. 1.34]. Thus,  $CGKB$  is right-angled. And it was also shown (to be) equilateral. Thus, it is a square. And it is on  $CB$ . So, for the same (reasons),  $HF$  is also a square. And it is on  $HG$ , that is to say [on]  $AC$  [Prop. 1.34]. Thus, the squares  $HF$  and  $KC$  are on  $AC$  and  $CB$  (respectively). And the (rectangle)  $AG$  is equal to the (rectangle)  $GE$  [Prop. 1.43]. And  $AG$  is the (rectangle contained) by  $AC$  and  $CB$ . For  $GC$  (is) equal to  $CB$ . Thus,  $GE$  is also equal to the (rectangle contained) by  $AC$  and  $CB$ . Thus, the (rectangles)  $AG$  and  $GE$  are equal to twice the (rectangle contained) by  $AC$  and  $CB$ . And  $HF$  and  $CK$  are the squares on  $AC$  and  $CB$  (respectively). Thus, the four (figures)  $HF$ ,  $CK$ ,  $AG$ , and  $GE$  are equal to the (sum of the) squares on

ὅλης τετράγωνον ἴσον ἐστὶ τοῖς τε ἀπὸ τῶν τμημάτων τετραγώνοις καὶ τῷ δις ὑπὸ τῶν τμημάτων περιεχομένῳ ὀρθογώνῳ· ὅπερ ἔδει δεῖξαι.


$AC$  and  $BC$ , and twice the rectangle contained by  $AC$  and  $CB$ . But, the (figures)  $HF$ ,  $CK$ ,  $AG$ , and  $GE$  are (equivalent to) the whole of  $ADEB$ , which is the square on  $AB$ . Thus, the square on  $AB$  is equal to the (sum of the) squares on  $AC$  and  $CB$ , and twice the rectangle contained by  $AC$  and  $CB$ .

Thus, if a straight-line is cut at random then the square on the whole (straight-line) is equal to the (sum of the) squares on the pieces (of the straight-line), and twice the rectangle contained by the pieces. (Which is) the very thing it was required to show.

† This proposition is a geometric version of the algebraic identity:  $(a + b)^2 = a^2 + b^2 + 2ab$ .

ε'.

Ἐὰν εὐθεῖα γραμμὴ τμηθῆ εἰς ἴσα καὶ ἄνισα, τὸ ὑπὸ τῶν ἀνίσων τῆς ὅλης τμημάτων περιεχόμενον ὀρθογώνιον μετὰ τοῦ ἀπὸ τῆς μεταξὺ τῶν τομῶν τετραγώνου ἴσον ἐστὶ τῷ ἀπὸ τῆς ἡμισείας τετραγώνῳ.




Εὐθεῖα γὰρ τις ἡ  $AB$  τετμήσθω εἰς μὲν ἴσα κατὰ τὸ  $\Gamma$ , εἰς δὲ ἄνισα κατὰ τὸ  $\Delta$ . λέγω, ὅτι τὸ ὑπὸ τῶν  $A\Delta$ ,  $\Delta B$  περιεχόμενον ὀρθογώνιον μετὰ τοῦ ἀπὸ τῆς  $\Gamma\Delta$  τετραγώνου ἴσον ἐστὶ τῷ ἀπὸ τῆς  $GB$  τετραγώνῳ.

Ἀναγεγράφθω γὰρ ἀπὸ τῆς  $GB$  τετράγωνον τὸ  $\Gamma EZB$ , καὶ ἐπεζεύχθω ἡ  $BE$ , καὶ διὰ μὲν τοῦ  $\Delta$  ὁποτέρᾳ τῶν  $\Gamma E$ ,  $BZ$  παράλληλος ἦχθω ἡ  $\Delta H$ , διὰ δὲ τοῦ  $\Theta$  ὁποτέρᾳ τῶν  $AB$ ,  $EZ$  παράλληλος πάλιν ἦχθω ἡ  $KM$ , καὶ πάλιν διὰ τοῦ  $A$  ὁποτέρᾳ τῶν  $\Gamma\Lambda$ ,  $BM$  παράλληλος ἦχθω ἡ  $AK$ . καὶ ἐπεὶ ἴσον ἐστὶ τὸ  $\Gamma\Theta$  παραπλήρωμα τῷ  $\Theta Z$  παραπληρώματι, κοινὸν προσκείσθω τὸ  $\Delta M$ . ὅλον ἄρα τὸ  $\Gamma M$  ὅλῳ τῷ  $\Delta Z$  ἴσον ἐστίν. ἀλλὰ τὸ  $\Gamma M$  τῷ  $A\Lambda$  ἴσον ἐστίν, ἐπεὶ καὶ ἡ  $A\Gamma$  τῆ  $\Gamma B$  ἐστὶν ἴση· καὶ τὸ  $A\Lambda$  ἄρα τῷ  $\Delta Z$  ἴσον ἐστίν. κοινὸν προσκείσθω τὸ  $\Gamma\Theta$ . ὅλον ἄρα τὸ  $A\Theta$  τῷ  $MN\Xi$  γνόμωνι ἴσον ἐστίν. ἀλλὰ τὸ  $A\Theta$  τὸ ὑπὸ τῶν  $A\Delta$ ,  $\Delta B$  ἐστίν· ἴση γὰρ ἡ  $\Delta\Theta$  τῆ  $\Delta B$ · καὶ ὁ  $MN\Xi$  ἄρα γνόμων ἴσος ἐστὶ τῷ ὑπὸ  $A\Delta$ ,  $\Delta B$ . κοινὸν προσκείσθω τὸ  $\Lambda H$ , ὃ ἐστὶν ἴσον τῷ ἀπὸ τῆς  $\Gamma\Delta$ . ὁ ἄρα  $MN\Xi$  γνόμων καὶ τὸ  $\Lambda H$  ἴσα ἐστὶ τῷ ὑπὸ τῶν  $A\Delta$ ,  $\Delta B$  περιεχομένῳ ὀρθογώνῳ καὶ τῷ ἀπὸ τῆς

Proposition 5<sup>‡</sup>

If a straight-line is cut into equal and unequal (pieces) then the rectangle contained by the unequal pieces of the whole (straight-line), plus the square on the (difference) between the (equal and unequal) pieces, is equal to the square on half (of the straight-line).



For let any straight-line  $AB$  have been cut—equally at  $C$ , and unequally at  $D$ . I say that the rectangle contained by  $AD$  and  $DB$ , plus the square on  $CD$ , is equal to the square on  $CB$ .

For let the square  $CEFB$  have been described on  $CB$  [Prop. 1.46], and let  $BE$  have been joined, and let  $DG$  have been drawn through  $D$ , parallel to either of  $CE$  or  $BF$  [Prop. 1.31], and again let  $KM$  have been drawn through  $H$ , parallel to either of  $AB$  or  $EF$  [Prop. 1.31], and again let  $AK$  have been drawn through  $A$ , parallel to either of  $CL$  or  $BM$  [Prop. 1.31]. And since the complement  $CH$  is equal to the complement  $HF$  [Prop. 1.43], let the (square)  $DM$  have been added to both. Thus, the whole (rectangle)  $CM$  is equal to the whole (rectangle)  $DF$ . But, (rectangle)  $CM$  is equal to (rectangle)  $AL$ , since  $AC$  is also equal to  $CB$  [Prop. 1.36]. Thus, (rectangle)  $AL$  is also equal to (rectangle)  $DF$ . Let (rectangle)  $CH$  have been added to both. Thus, the whole (rectangle)  $AH$  is equal to the gnomon  $NOP$ . But,  $AH$

ΓΔ τετραγώνω. ἀλλὰ ὁ ΜΝΞ γνώμων καὶ τὸ ΛΗ ὄλον ἐστὶ τὸ ΓΕΖΒ τετράγωνον, ὃ ἐστὶν ἀπὸ τῆς ΓΒ· τὸ ἄρα ὑπὸ τῶν ΑΔ, ΔΒ περιεχόμενον ὀρθογώνιον μετὰ τοῦ ἀπὸ τῆς ΓΔ τετραγώνου ἴσον ἐστὶ τῷ ἀπὸ τῆς ΓΒ τετραγώνω.

Ἐὰν ἄρα εὐθεῖα γραμμὴ τμηθῆ εἰς ἴσα καὶ ἄνισα, τὸ ὑπὸ τῶν ἀνίσων τῆς ὅλης τμημάτων περιεχόμενον ὀρθογώνιον μετὰ τοῦ ἀπὸ τῆς μεταξὺ τῶν τομῶν τετραγώνου ἴσον ἐστὶ τῷ ἀπὸ τῆς ἡμισείας τετραγώνω. ὅπερ ἔδει δεῖξαι.

is the (rectangle contained) by  $AD$  and  $DB$ . For  $DH$  (is) equal to  $DB$ . Thus, the gnomon  $NOP$  is also equal to the (rectangle contained) by  $AD$  and  $DB$ . Let  $LG$ , which is equal to the (square) on  $CD$ , have been added to both. Thus, the gnomon  $NOP$  and the (square)  $LG$  are equal to the rectangle contained by  $AD$  and  $DB$ , and the square on  $CD$ . But, the gnomon  $NOP$  and the (square)  $LG$  is (equivalent to) the whole square  $CEFB$ , which is on  $CB$ . Thus, the rectangle contained by  $AD$  and  $DB$ , plus the square on  $CD$ , is equal to the square on  $CB$ .


Thus, if a straight-line is cut into equal and unequal (pieces) then the rectangle contained by the unequal pieces of the whole (straight-line), plus the square on the (difference) between the (equal and unequal) pieces, is equal to the square on half (of the straight-line). (Which is) the very thing it was required to show.

† Note the (presumably mistaken) double use of the label  $M$  in the Greek text.

‡ This proposition is a geometric version of the algebraic identity:  $ab + [(a + b)/2 - b]^2 = [(a + b)/2]^2$ .

ζ'.

Ἐὰν εὐθεῖα γραμμὴ τμηθῆ δίχα, προστεθῆ δὲ τις αὐτῇ εὐθεῖα ἐπ' εὐθείας, τὸ ὑπὸ τῆς ὅλης σὺν τῇ προσκειμένῃ καὶ τῆς προσκειμένης περιεχόμενον ὀρθογώνιον μετὰ τοῦ ἀπὸ τῆς ἡμισείας τετραγώνου ἴσον ἐστὶ τῷ ἀπὸ τῆς συγκεκλιμένης ἔκ τε τῆς ἡμισείας καὶ τῆς προσκειμένης τετραγώνω.




Εὐθεῖα γάρ τις ἡ  $AB$  τεμηθῆτω δίχα κατὰ τὸ  $\Gamma$  σημεῖον, προσκείσθω δὲ τις αὐτῇ εὐθεῖα ἐπ' εὐθείας ἡ  $B\Delta$ . λέγω, ὅτι τὸ ὑπὸ τῶν  $A\Delta$ ,  $\Delta B$  περιεχόμενον ὀρθογώνιον μετὰ τοῦ ἀπὸ τῆς  $\Gamma B$  τετραγώνου ἴσον ἐστὶ τῷ ἀπὸ τῆς  $\Gamma\Delta$  τετραγώνω.

Ἀναγεγράφθω γὰρ ἀπὸ τῆς  $\Gamma\Delta$  τετράγωνον τὸ  $\Gamma E Z \Delta$ , καὶ ἐπεζεύχθω ἡ  $\Delta E$ , καὶ διὰ μὲν τοῦ  $B$  σημείου ὁποτέρᾳ τῶν  $E\Gamma$ ,  $\Delta Z$  παράλληλος ἦχθω ἡ  $BH$ , διὰ δὲ τοῦ  $\Theta$  σημείου ὁποτέρᾳ τῶν  $AB$ ,  $EZ$  παράλληλος ἦχθω ἡ  $KM$ , καὶ ἔτι διὰ τοῦ  $A$  ὁποτέρᾳ τῶν  $\Gamma\Lambda$ ,  $\Delta M$  παράλληλος ἦχθω ἡ  $AK$ .

Ἐπεὶ οὖν ἴση ἐστὶν ἡ  $A\Gamma$  τῇ  $\Gamma B$ , ἴσον ἐστὶ καὶ τὸ  $AK$

Proposition 6†

If a straight-line is cut in half, and any straight-line added to it straight-on, then the rectangle contained by the whole (straight-line) with the (straight-line) having been added, and the (straight-line) having been added, plus the square on half (of the original straight-line), is equal to the square on the sum of half (of the original straight-line) and the (straight-line) having been added.



For let any straight-line  $AB$  have been cut in half at point  $C$ , and let any straight-line  $BD$  have been added to it straight-on. I say that the rectangle contained by  $AD$  and  $DB$ , plus the square on  $CB$ , is equal to the square on  $CD$ .

For let the square  $CEFD$  have been described on  $CD$  [Prop. 1.46], and let  $DE$  have been joined, and let  $BG$  have been drawn through point  $B$ , parallel to either of  $EC$  or  $DF$  [Prop. 1.31], and let  $KM$  have been drawn through point  $H$ , parallel to either of  $AB$  or  $EF$  [Prop. 1.31], and finally let  $AK$  have been drawn

τῶ ΓΘ. ἀλλὰ τὸ ΓΘ τῶ ΘΖ ἴσον ἐστίν. καὶ τὸ ΑΛ ἄρα τῶ ΘΖ ἐστὶν ἴσον. κοινὸν προσκείσθω τὸ ΓΜ· ὅλον ἄρα τὸ ΑΜ τῶ ΝΞΟ γνώμονι ἐστὶν ἴσον. ἀλλὰ τὸ ΑΜ ἐστὶ τὸ ὑπὸ τῶν ΑΔ, ΔΒ· ἴση γάρ ἐστὶν ἡ ΔΜ τῆι ΔΒ· καὶ ὁ ΝΞΟ ἄρα γνώμων ἴσος ἐστὶ τῶ ὑπὸ τῶν ΑΔ, ΔΒ [περιεχομένῳ ὀρθογωνίῳ]. κοινὸν προσκείσθω τὸ ΛΗ, ὃ ἐστὶν ἴσον τῶ ἀπὸ τῆς ΒΓ τετραγώνῳ· τὸ ἄρα ὑπὸ τῶν ΑΔ, ΔΒ περιεχόμενον ὀρθογώνιον μετὰ τοῦ ἀπὸ τῆς ΓΒ τετραγώνου ἴσον ἐστὶ τῶ ΝΞΟ γνώμονι καὶ τῶ ΛΗ. ἀλλὰ ὁ ΝΞΟ γνώμων καὶ τὸ ΛΗ ὅλον ἐστὶ τὸ ΓΕΖΔ τετράγωνον, ὃ ἐστὶν ἀπὸ τῆς ΓΔ· τὸ ἄρα ὑπὸ τῶν ΑΔ, ΔΒ περιεχόμενον ὀρθογώνιον μετὰ τοῦ ἀπὸ τῆς ΓΒ τετραγώνου ἴσον ἐστὶ τῶ ἀπὸ τῆς ΓΔ τετραγώνῳ.

Ἐὰν ἄρα εὐθεῖα γραμμὴ τμηθῆ διχα, προστεθῆ δέ τις αὐτῇ εὐθεῖα ἐπ' εὐθείας, τὸ ὑπὸ τῆς ὅλης σὺν τῇ προσκειμένη καὶ τῆς προσκειμένης περιεχόμενον ὀρθογώνιον μετὰ τοῦ ἀπὸ τῆς ἡμισείας τετραγώνου ἴσον ἐστὶ τῶ ἀπὸ τῆς συγκεκλιμένης ἕκ τε τῆς ἡμισείας καὶ τῆς προσκειμένης τετραγώνῳ· ὅπερ ἔδει δεῖξαι.

through  $A$ , parallel to either of  $CL$  or  $DM$  [Prop. 1.31].


Therefore, since  $AC$  is equal to  $CB$ , (rectangle)  $AL$  is also equal to (rectangle)  $CH$  [Prop. 1.36]. But, (rectangle)  $CH$  is equal to (rectangle)  $HF$  [Prop. 1.43]. Thus, (rectangle)  $AL$  is also equal to (rectangle)  $HF$ . Let (rectangle)  $CM$  have been added to both. Thus, the whole (rectangle)  $AM$  is equal to the gnomon  $NOP$ . But,  $AM$  is the (rectangle contained) by  $AD$  and  $DB$ . For  $DM$  is equal to  $DB$ . Thus, gnomon  $NOP$  is also equal to the [rectangle contained] by  $AD$  and  $DB$ . Let  $LG$ , which is equal to the square on  $BC$ , have been added to both. Thus, the rectangle contained by  $AD$  and  $DB$ , plus the square on  $CB$ , is equal to the gnomon  $NOP$  and the (square)  $LG$ . But the gnomon  $NOP$  and the (square)  $LG$  is (equivalent to) the whole square  $CEFD$ , which is on  $CD$ . Thus, the rectangle contained by  $AD$  and  $DB$ , plus the square on  $CB$ , is equal to the square on  $CD$ .

Thus, if a straight-line is cut in half, and any straight-line added to it straight-on, then the rectangle contained by the whole (straight-line) with the (straight-line) having being added, and the (straight-line) having being added, plus the square on half (of the original straight-line), is equal to the square on the sum of half (of the original straight-line) and the (straight-line) having been added. (Which is) the very thing it was required to show.

† This proposition is a geometric version of the algebraic identity:  $(2a + b)b + a^2 = (a + b)^2$ .

ζ'.


Ἐὰν εὐθεῖα γραμμὴ τμηθῆ, ὡς ἔτυχεν, τὸ ἀπὸ τῆς ὅλης καὶ τὸ ἀφ' ἑνὸς τῶν τμημάτων τὰ συναμφοτέρα τετράγωνα ἴσα ἐστὶ τῶ τε δις ὑπὸ τῆς ὅλης καὶ τοῦ εἰρημένου τμήματος περιεχομένῳ ὀρθογωνίῳ καὶ τῶ ἀπὸ τοῦ λοιποῦ τμήματος τετραγώνῳ.



Εὐθεῖα γάρ τις ἡ  $AB$  τεμηθῆσθω, ὡς ἔτυχεν, κατὰ τὸ  $\Gamma$  σημεῖον· λέγω, ὅτι τὰ ἀπὸ τῶν  $AB$ ,  $B\Gamma$  τετράγωνα ἴσα ἐστὶ τῶ τε δις ὑπὸ τῶν  $AB$ ,  $B\Gamma$  περιεχομένῳ ὀρθογωνίῳ καὶ τῶ

Proposition 7†

If a straight-line is cut at random then the sum of the squares on the whole (straight-line), and one of the pieces (of the straight-line), is equal to twice the rectangle contained by the whole, and the said piece, and the square on the remaining piece.



For let any straight-line  $AB$  have been cut, at random, at point  $C$ . I say that the (sum of the) squares on  $AB$  and  $BC$  is equal to twice the rectangle contained by  $AB$  and



ἀπὸ τῆς ΓΑ τετραγώνω.

Ἀναγεγράφθω γὰρ ἀπὸ τῆς ΑΒ τετράγωνον τὸ ΑΔΕΒ· καὶ καταγεγράφθω τὸ σχῆμα.

Ἐπεὶ οὖν ἴσον ἐστὶ τὸ ΑΗ τῷ ΗΕ, κοινὸν προσκείσθω τὸ ΓΖ· ὅλον ἄρα τὸ ΑΖ ὅλω τῷ ΓΕ ἴσον ἐστίν· τὰ ἄρα ΑΖ, ΓΕ διπλάσιά ἐστι τοῦ ΑΖ. ἀλλὰ τὰ ΑΖ, ΓΕ ὁ ΚΑΜ ἐστὶ γνῶμων καὶ τὸ ΓΖ τετράγωνον· ὁ ΚΑΜ ἄρα γνῶμων καὶ τὸ ΓΖ διπλάσιά ἐστι τοῦ ΑΖ. ἐστὶ δὲ τοῦ ΑΖ διπλάσιον καὶ τὸ δις ὑπὸ τῶν ΑΒ, ΒΓ· ἴση γὰρ ἡ ΒΖ τῇ ΒΓ· ὁ ἄρα ΚΑΜ γνῶμων καὶ τὸ ΓΖ τετράγωνον ἴσον ἐστὶ τῷ δις ὑπὸ τῶν ΑΒ, ΒΓ. κοινὸν προσκείσθω τὸ ΔΗ, ὃ ἐστὶν ἀπὸ τῆς ΑΓ τετράγωνον· ὁ ἄρα ΚΑΜ γνῶμων καὶ τὰ ΒΗ, ΗΔ τετράγωνα ἴσα ἐστὶ τῷ τε δις ὑπὸ τῶν ΑΒ, ΒΓ περιεχομένῳ ὀρθογωνίῳ καὶ τῷ ἀπὸ τῆς ΑΓ τετραγώνῳ. ἀλλὰ ὁ ΚΑΜ γνῶμων καὶ τὰ ΒΗ, ΗΔ τετράγωνα ὅλον ἐστὶ τὸ ΑΔΕΒ καὶ τὸ ΓΖ, ἃ ἐστὶν ἀπὸ τῶν ΑΒ, ΒΓ τετράγωνα· τὰ ἄρα ἀπὸ τῶν ΑΒ, ΒΓ τετράγωνα ἴσα ἐστὶ τῷ [τε] δις ὑπὸ τῶν ΑΒ, ΒΓ περιεχομένῳ ὀρθογωνίῳ μετὰ τοῦ ἀπὸ τῆς ΑΓ τετραγώνου.

Ἐὰν ἄρα εὐθεῖα γραμμὴ τμηθῆ, ὡς ἔτυχεν, τὸ ἀπὸ τῆς ὅλης καὶ τὸ ἀφ' ἑνὸς τῶν τμημάτων τὰ συναμφοτέρα τετράγωνα ἴσα ἐστὶ τῷ τε δις ὑπὸ τῆς ὅλης καὶ τοῦ εἰρημένου τμήματος περιεχομένῳ ὀρθογωνίῳ καὶ τῷ ἀπὸ τοῦ λοιποῦ τμήματος τετραγώνῳ· ὅπερ εἶδει δεῖξαι.

*BC*, and the square on *CA*.

For let the square *ADEB* have been described on *AB* [Prop. 1.46], and let the (rest of) the figure have been drawn.

Therefore, since (rectangle) *AG* is equal to (rectangle) *GE* [Prop. 1.43], let the (square) *CF* have been added to both. Thus, the whole (rectangle) *AF* is equal to the whole (rectangle) *CE*. Thus, (rectangle) *AF* plus (rectangle) *CE* is double (rectangle) *AF*. But, (rectangle) *AF* plus (rectangle) *CE* is the gnomon *KLM*, and the square *CF*. Thus, the gnomon *KLM*, and the square *CF*, is double the (rectangle) *AF*. But double the (rectangle) *AF* is also twice the (rectangle contained) by *AB* and *BC*. For *BF* (is) equal to *BC*. Thus, the gnomon *KLM*, and the square *CF*, are equal to twice the (rectangle contained) by *AB* and *BC*. Let *DG*, which is the square on *AC*, have been added to both. Thus, the gnomon *KLM*, and the squares *BG* and *GD*, are equal to twice the rectangle contained by *AB* and *BC*, and the square on *AC*. But, the gnomon *KLM* and the squares *BG* and *GD* is (equivalent to) the whole of *ADEB* and *CF*, which are the squares on *AB* and *BC* (respectively). Thus, the (sum of the) squares on *AB* and *BC* is equal to twice the rectangle contained by *AB* and *BC*, and the square on *AC*.

Thus, if a straight-line is cut at random then the sum of the squares on the whole (straight-line), and one of the pieces (of the straight-line), is equal to twice the rectangle contained by the whole, and the said piece, and the square on the remaining piece. (Which is) the very thing it was required to show.

† This proposition is a geometric version of the algebraic identity:  $(a + b)^2 + a^2 = 2(a + b)a + b^2$ .

η'.

Ἐὰν εὐθεῖα γραμμὴ τμηθῆ, ὡς ἔτυχεν, τὸ τετράκις ὑπὸ τῆς ὅλης καὶ ἑνὸς τῶν τμημάτων περιεχόμενον ὀρθογώνιον μετὰ τοῦ ἀπὸ τοῦ λοιποῦ τμήματος τετραγώνου ἴσον ἐστὶ τῷ ἀπὸ τε τῆς ὅλης καὶ τοῦ εἰρημένου τμήματος ὡς ἀπὸ μιᾶς ἀναγραφέντι τετραγώνῳ.

Εὐθεῖα γὰρ τις ἡ ΑΒ τετμήσθω, ὡς ἔτυχεν, κατὰ τὸ Γ σημεῖον· λέγω, ὅτι τὸ τετράκις ὑπὸ τῶν ΑΒ, ΒΓ περιεχόμενον ὀρθογώνιον μετὰ τοῦ ἀπὸ τῆς ΑΓ τετραγώνου ἴσον ἐστὶ τῷ ἀπὸ τῆς ΑΒ, ΒΓ ὡς ἀπὸ μιᾶς ἀναγραφέντι τετραγώνῳ.



Ἐκβεβλήσθω γὰρ ἐπ' εὐθείας [τῇ ΑΒ εὐθείᾳ] ἡ ΒΔ, καὶ κείσθω τῇ ΓΒ ἴση ἡ ΒΔ, καὶ ἀναγεγράφθω ἀπὸ τῆς ΑΔ τετράγωνον τὸ ΑΕΖΔ, καὶ καταγεγράφθω διπλοῦν τὸ σχῆμα.

### Proposition 8<sup>†</sup>

If a straight-line is cut at random then four times the rectangle contained by the whole (straight-line), and one of the pieces (of the straight-line), plus the square on the remaining piece, is equal to the square described on the whole and the former piece, as on one (complete straight-line).

For let any straight-line *AB* have been cut, at random, at point *C*. I say that four times the rectangle contained by *AB* and *BC*, plus the square on *AC*, is equal to the square described on *AB* and *BC*, as on one (complete straight-line).

For let *BD* have been produced in a straight-line [with the straight-line *AB*], and let *BD* be made equal to *CB* [Prop. 1.3], and let the square *Aefd* have been described on *AD* [Prop. 1.46], and let the (rest of the) figure have been drawn double.



Ἐπει οὖν ἴση ἐστὶν ἡ ΓΒ τῆ ΒΔ, ἀλλὰ ἡ μὲν ΓΒ τῆ ΗΚ ἐστὶν ἴση, ἡ δὲ ΒΔ τῆ ΚΝ, καὶ ἡ ΗΚ ἄρα τῆ ΚΝ ἐστὶν ἴση. διὰ τὰ αὐτὰ δὴ καὶ ἡ ΠΡ τῆ ΡΟ ἐστὶν ἴση. καὶ ἐπει ἴση ἐστὶν ἡ ΒΓ τῆ ΒΔ, ἡ δὲ ΗΚ τῆ ΚΝ, ἴσον ἄρα ἐστὶ καὶ τὸ μὲν ΓΚ τῷ ΚΔ, τὸ δὲ ΗΡ τῷ ΡΝ. ἀλλὰ τὸ ΓΚ τῷ ΡΝ ἐστὶν ἴσον· παραπληρώματα γὰρ τοῦ ΓΟ παραλληλογράμμου· καὶ τὸ ΚΔ ἄρα τῷ ΗΡ ἴσον ἐστίν· τὰ τέσσαρα ἄρα τὰ ΔΚ, ΓΚ, ΗΡ, ΡΝ ἴσα ἀλλήλοις ἐστίν. τὰ τέσσαρα ἄρα τετραπλάσια ἐστὶ τοῦ ΓΚ. πάλιν ἐπει ἴση ἐστὶν ἡ ΓΒ τῆ ΒΔ, ἀλλὰ ἡ μὲν ΒΔ τῆ ΒΚ, τουτέστι τῆ ΓΗ ἴση, ἡ δὲ ΓΒ τῆ ΗΚ, τουτέστι τῆ ΗΠ, ἐστὶν ἴση, καὶ ἡ ΓΗ ἄρα τῆ ΗΠ ἴση ἐστίν. καὶ ἐπει ἴση ἐστὶν ἡ μὲν ΓΗ τῆ ΗΠ, ἡ δὲ ΠΡ τῆ ΡΟ, ἴσον ἐστὶ καὶ τὸ μὲν ΑΗ τῷ ΜΠ, τὸ δὲ ΠΛ τῷ ΡΖ. ἀλλὰ τὸ ΜΠ τῷ ΠΛ ἐστὶν ἴσον· παραπληρώματα γὰρ τοῦ ΜΛ παραλληλογράμμου· καὶ τὸ ΑΗ ἄρα τῷ ΡΖ ἴσον ἐστίν· τὰ τέσσαρα ἄρα τὰ ΑΗ, ΜΠ, ΠΛ, ΡΖ ἴσα ἀλλήλοις ἐστίν· τὰ τέσσαρα ἄρα τοῦ ΑΗ ἐστὶ τετραπλάσια. ἐδείχθη δὲ καὶ τὰ τέσσαρα τὰ ΓΚ, ΚΔ, ΗΡ, ΡΝ τοῦ ΓΚ τετραπλάσια· τὰ ἄρα ὀκτώ, ἃ περιέχει τὸν ΣΤΥ γνῶμονα, τετραπλάσια ἐστὶ τοῦ ΑΚ. καὶ ἐπει τὸ ΑΚ τὸ ὑπὸ τῶν ΑΒ, ΒΔ ἐστίν· ἴση γὰρ ἡ ΒΚ τῆ ΒΔ· τὸ ἄρα τετράκις ὑπὸ τῶν ΑΒ, ΒΔ τετραπλάσιόν ἐστὶ τοῦ ΑΚ. ἐδείχθη δὲ τοῦ ΑΚ τετραπλάσιος καὶ ὁ ΣΤΥ γνῶμων· τὸ ἄρα τετράκις ὑπὸ τῶν ΑΒ, ΒΔ ἴσον ἐστὶ τῷ ΣΤΥ γνῶμονι. κοινὸν προσκείσθω τὸ ΞΘ, ὃ ἐστὶν ἴσον τῷ ἀπὸ τῆς ΑΓ τετραγώνῳ· τὸ ἄρα τετράκις ὑπὸ τῶν ΑΒ, ΒΔ περιεχόμενον ὀρθογώνιον μετὰ τοῦ ἀπὸ ΑΓ τετραγώνου ἴσον ἐστὶ τῷ ΣΤΥ γνῶμονι καὶ τῷ ΞΘ. ἀλλὰ ὁ ΣΤΥ γνῶμων καὶ τὸ ΞΘ ὅλον ἐστὶ τὸ ΑΕΖΔ τετραγώνον, ὃ ἐστὶν ἀπὸ τῆς ΑΔ· τὸ ἄρα τετράκις ὑπὸ τῶν ΑΒ, ΒΔ μετὰ τοῦ ἀπὸ ΑΓ ἴσον ἐστὶ τῷ ἀπὸ ΑΔ τετραγώνῳ· ἴση δὲ ἡ ΒΔ τῆ ΒΓ. τὸ ἄρα τετράκις ὑπὸ τῶν ΑΒ, ΒΓ περιεχόμενον ὀρθογώνιον μετὰ τοῦ ἀπὸ ΑΓ τετραγώνου ἴσον ἐστὶ τῷ ἀπὸ τῆς ΑΔ, τουτέστι τῷ ἀπὸ τῆς ΑΒ καὶ ΒΓ ὡς ἀπὸ μιᾶς ἀναγραφέντι τετραγώνῳ.

Ἐάν ἄρα εὐθεῖα γραμμὴ τμηθῆ, ὡς ἔτυχεν, τὸ τετράκις ὑπὸ τῆς ὅλης καὶ ἐνὸς τῶν τμημάτων περιεχόμενον ὀρθογώνιον μετὰ τοῦ ἀπὸ τοῦ λοιποῦ τμήματος τετραγώνου ἴσου

Therefore, since  $CB$  is equal to  $BD$ , but  $CB$  is equal to  $GK$  [Prop. 1.34], and  $BD$  to  $KN$  [Prop. 1.34],  $GK$  is thus also equal to  $KN$ . So, for the same (reasons),  $QR$  is equal to  $RP$ . And since  $BC$  is equal to  $BD$ , and  $GK$  to  $KN$ , (square)  $CK$  is thus also equal to (square)  $KD$ , and (square)  $GR$  to (square)  $RN$  [Prop. 1.36]. But, (square)  $CK$  is equal to (square)  $RN$ . For (they are) complements in the parallelogram  $CP$  [Prop. 1.43]. Thus, (square)  $KD$  is also equal to (square)  $GR$ . Thus, the four (squares)  $DK, CK, GR,$  and  $RN$  are equal to one another. Thus, the four (taken together) are quadruple (square)  $CK$ . Again, since  $CB$  is equal to  $BD$ , but  $BD$  (is) equal to  $BK$ —that is to say,  $CG$ —and  $CB$  is equal to  $GK$ —that is to say,  $GQ$ — $CG$  is thus also equal to  $GQ$ . And since  $CG$  is equal to  $GQ$ , and  $QR$  to  $RP$ , (rectangle)  $AG$  is also equal to (rectangle)  $MQ$ , and (rectangle)  $QL$  to (rectangle)  $RF$  [Prop. 1.36]. But, (rectangle)  $MQ$  is equal to (rectangle)  $QL$ . For (they are) complements in the parallelogram  $ML$  [Prop. 1.43]. Thus, (rectangle)  $AG$  is also equal to (rectangle)  $RF$ . Thus, the four (rectangles)  $AG, MQ, QL,$  and  $RF$  are equal to one another. Thus, the four (taken together) are quadruple (rectangle)  $AG$ . And it was also shown that the four (squares)  $CK, KD, GR,$  and  $RN$  (taken together are) quadruple (square)  $CK$ . Thus, the eight (figures taken together), which comprise the gnomon  $STU$ , are quadruple (rectangle)  $AK$ . And since  $AK$  is the (rectangle contained) by  $AB$  and  $BD$ , for  $BK$  (is) equal to  $BD$ , four times the (rectangle contained) by  $AB$  and  $BD$  is quadruple (rectangle)  $AK$ . But the gnomon  $STU$  was also shown (to be equal to) quadruple (rectangle)  $AK$ . Thus, four times the (rectangle contained) by  $AB$  and  $BD$  is equal to the gnomon  $STU$ . Let  $OH$ , which is equal to the square on  $AC$ , have been added to both. Thus, four times the rectangle contained by  $AB$  and  $BD$ , plus the square on  $AC$ , is equal to the gnomon  $STU$ , and the (square)  $OH$ . But,

ἔστι τῷ ἀπό τε τῆς ὅλης καὶ τοῦ εἰρημένου τμήματος ὡς ἀπὸ μιᾶς ἀναγραφέντι τετραγώνῳ· ὅπερ ἔδει δεῖξαι.


the gnomon  $STU$  and the (square)  $OH$  is (equivalent to) the whole square  $AEFD$ , which is on  $AD$ . Thus, four times the (rectangle contained) by  $AB$  and  $BD$ , plus the (square) on  $AC$ , is equal to the square on  $AD$ . And  $BD$  (is) equal to  $BC$ . Thus, four times the rectangle contained by  $AB$  and  $BC$ , plus the square on  $AC$ , is equal to the (square) on  $AD$ , that is to say the square described on  $AB$  and  $BC$ , as on one (complete straight-line).

Thus, if a straight-line is cut at random then four times the rectangle contained by the whole (straight-line), and one of the pieces (of the straight-line), plus the square on the remaining piece, is equal to the square described on the whole and the former piece, as on one (complete straight-line). (Which is) the very thing it was required to show.

† This proposition is a geometric version of the algebraic identity:  $4(a + b)a + b^2 = [(a + b) + a]^2$ .

θ'.

Ἐὰν εὐθεῖα γραμμὴ τμηθῆ εἰς ἴσα καὶ ἄνισα, τὰ ἀπὸ τῶν ἀνίσων τῆς ὅλης τμημάτων τετράγωνα διπλάσιά ἐστι τοῦ τε ἀπὸ τῆς ἡμίσειας καὶ τοῦ ἀπὸ τῆς μεταξύ τῶν τομῶν τετραγώνου.




Εὐθεῖα γὰρ τις ἡ  $AB$  τετμήσθω εἰς μὲν ἴσα κατὰ τὸ  $\Gamma$ , εἰς δὲ ἄνισα κατὰ τὸ  $\Delta$ . λέγω, ὅτι τὰ ἀπὸ τῶν  $A\Delta$ ,  $\Delta B$  τετράγωνα διπλάσιά ἐστι τῶν ἀπὸ τῶν  $A\Gamma$ ,  $\Gamma\Delta$  τετραγώνων.

Ἦχθω γὰρ ἀπὸ τοῦ  $\Gamma$  τῆς  $AB$  πρὸς ὀρθὰς ἡ  $\Gamma E$ , καὶ κείσθω ἴση ἑκατέρω τῶν  $A\Gamma$ ,  $\Gamma B$ , καὶ ἐπεζεύχθωσαν αἱ  $EA$ ,  $EB$ , καὶ διὰ μὲν τοῦ  $\Delta$  τῆς  $EG$  παράλληλος ἦχθω ἡ  $\Delta Z$ , διὰ δὲ τοῦ  $Z$  τῆς  $AB$  ἡ  $ZH$ , καὶ ἐπεζεύχθω ἡ  $AZ$ . καὶ ἐπεὶ ἴση ἐστὶν ἡ  $A\Gamma$  τῆς  $\Gamma E$ , ἴση ἐστὶ καὶ ἡ ὑπὸ  $EAG$  γωνία τῆς ὑπὸ  $AEG$ . καὶ ἐπεὶ ὀρθὴ ἐστὶν ἡ πρὸς τῷ  $\Gamma$ , λοιπαὶ ἄρα αἱ ὑπὸ  $EAG$ ,  $AEG$  μιᾶ ὀρθῇ ἴσαι εἰσὶν· καὶ εἰσὶν ἴσαι· ἡμίσεια ἄρα ὀρθῆς ἐστὶν ἑκατέρω τῶν ὑπὸ  $GEA$ ,  $FAE$ . διὰ τὰ αὐτὰ δὲ καὶ ἑκατέρω τῶν ὑπὸ  $GEB$ ,  $EBF$  ἡμίσειά ἐστὶν ὀρθῆς· ὅλη ἄρα ἡ ὑπὸ  $AEB$  ὀρθὴ ἐστὶν. καὶ ἐπεὶ ἡ ὑπὸ  $HEZ$  ἡμίσειά ἐστὶν ὀρθῆς, ὀρθὴ δὲ ἡ ὑπὸ  $EHZ$ · ἴση γὰρ ἐστὶ τῆς ἐντὸς καὶ ἀπεναντίον τῆς ὑπὸ  $EGB$ · λοιπὴ ἄρα ἡ ὑπὸ  $EZH$  ἡμίσειά ἐστὶν

Proposition 9†

If a straight-line is cut into equal and unequal (pieces) then the (sum of the) squares on the unequal pieces of the whole (straight-line) is double the (sum of the) square on half (the straight-line) and (the square) on the (difference) between the (equal and unequal) pieces.



For let any straight-line  $AB$  have been cut—equally at  $C$ , and unequally at  $D$ . I say that the (sum of the) squares on  $AD$  and  $DB$  is double the (sum of the squares) on  $AC$  and  $CD$ .

For let  $CE$  have been drawn from (point)  $C$ , at right-angles to  $AB$  [Prop. 1.11], and let it be made equal to each of  $AC$  and  $CB$  [Prop. 1.3], and let  $EA$  and  $EB$  have been joined. And let  $DF$  have been drawn through (point)  $D$ , parallel to  $EC$  [Prop. 1.31], and (let)  $FG$  (have been drawn) through (point)  $F$ , (parallel) to  $AB$  [Prop. 1.31]. And let  $AF$  have been joined. And since  $AC$  is equal to  $CE$ , the angle  $EAC$  is also equal to the (angle)  $AEC$  [Prop. 1.5]. And since the (angle) at  $C$  is a right-angle, the (sum of the) remaining angles (of triangle  $AEC$ ),  $EAC$  and  $AEC$ , is thus equal to one right-

ὀρθῆς· ἴση ἄρα [ἐστίν] ἡ ὑπὸ HEZ γωνία τῇ ὑπὸ EZH· ὥστε καὶ πλευρὰ ἡ EH τῇ HZ ἐστίν ἴση. πάλιν ἐπεὶ ἡ πρὸς τῷ B γωνία ἡμίσειά ἐστιν ὀρθῆς, ὀρθὴ δὲ ἡ ὑπὸ ZΔB· ἴση γὰρ πάλιν ἐστὶ τῇ ἐντὸς καὶ ἀπεναντίον τῇ ὑπὸ EΓB· λοιπὴ ἄρα ἡ ὑπὸ BZΔ ἡμίσειά ἐστιν ὀρθῆς· ἴση ἄρα ἡ πρὸς τῷ B γωνία τῇ ὑπὸ ΔZB· ὥστε καὶ πλευρὰ ἡ ZΔ πλευρᾷ τῇ ΔB ἐστίν ἴση. καὶ ἐπεὶ ἴση ἐστίν ἡ ΑΓ τῇ ΓΕ, ἴσον ἐστὶ καὶ τὸ ἀπὸ ΑΓ τῷ ἀπὸ ΓΕ· τὰ ἄρα ἀπὸ τῶν ΑΓ, ΓΕ τετράγωνα διπλάσιά ἐστι τοῦ ἀπὸ ΑΓ. τοῖς δὲ ἀπὸ τῶν ΑΓ, ΓΕ ἴσον ἐστὶ τὸ ἀπὸ τῆς ΕΑ τετράγωνον· ὀρθὴ γὰρ ἡ ὑπὸ ΑΓΕ γωνία· τὸ ἄρα ἀπὸ τῆς ΕΑ διπλάσιόν ἐστι τοῦ ἀπὸ τῆς ΑΓ. πάλιν, ἐπεὶ ἴση ἐστίν ἡ ΕΗ τῇ ΗΖ, ἴσον καὶ τὸ ἀπὸ τῆς ΕΗ τῷ ἀπὸ τῆς ΗΖ· τὰ ἄρα ἀπὸ τῶν ΕΗ, ΗΖ τετράγωνα διπλάσιά ἐστι τοῦ ἀπὸ τῆς ΗΖ τετραγώνου. τοῖς δὲ ἀπὸ τῶν ΕΗ, ΗΖ τετραγώνοις ἴσον ἐστὶ τὸ ἀπὸ τῆς ΕΖ τετράγωνον· τὸ ἄρα ἀπὸ τῆς ΕΖ τετράγωνον διπλάσιόν ἐστι τοῦ ἀπὸ τῆς ΗΖ. ἴση δὲ ἡ ΗΖ τῇ ΓΔ· τὸ ἄρα ἀπὸ τῆς ΕΖ διπλάσιόν ἐστι τοῦ ἀπὸ τῆς ΓΔ. ἔστι δὲ καὶ τὸ ἀπὸ τῆς ΕΑ διπλάσιον τοῦ ἀπὸ τῆς ΑΓ· τὰ ἄρα ἀπὸ τῶν ΑΕ, ΕΖ τετράγωνα διπλάσιά ἐστι τῶν ἀπὸ τῶν ΑΓ, ΓΔ τετραγώνων. τοῖς δὲ ἀπὸ τῶν ΑΕ, ΕΖ ἴσον ἐστὶ τὸ ἀπὸ τῆς ΑΖ τετράγωνον· ὀρθὴ γὰρ ἐστίν ἡ ὑπὸ ΑΕΖ γωνία· τὸ ἄρα ἀπὸ τῆς ΑΖ τετράγωνον διπλάσιόν ἐστι τῶν ἀπὸ τῶν ΑΓ, ΓΔ. τῷ δὲ ἀπὸ τῆς ΑΖ ἴσα τὰ ἀπὸ τῶν ΑΔ, ΔΖ· ὀρθὴ γὰρ ἡ πρὸς τῷ Δ γωνία· τὰ ἄρα ἀπὸ τῶν ΑΔ, ΔΖ διπλάσιά ἐστι τῶν ἀπὸ τῶν ΑΓ, ΓΔ τετραγώνων. ἴση δὲ ἡ ΔΖ τῇ ΔB· τὰ ἄρα ἀπὸ τῶν ΑΔ, ΔB τετράγωνα διπλάσιά ἐστι τῶν ἀπὸ τῶν ΑΓ, ΓΔ τετραγώνων.

Ἐὰν ἄρα εὐθεῖα γραμμὴ τμηθῇ εἰς ἴσα καὶ ἄνισα, τὰ ἀπὸ τῶν ἀνίσων τῆς ὅλης τμημάτων τετράγωνα διπλάσιά ἐστι τοῦ τε ἀπὸ τῆς ἡμίσειας καὶ τοῦ ἀπὸ τῆς μεταξὺ τῶν τομῶν τετραγώνου· ὅπερ ἔδει δεῖξαι.


angle [Prop. 1.32]. And they are equal. Thus, (angles)  $CEA$  and  $CAE$  are each half a right-angle. So, for the same (reasons), (angles)  $CEB$  and  $EBC$  are also each half a right-angle. Thus, the whole (angle)  $AEB$  is a right-angle. And since  $GEF$  is half a right-angle, and  $EGF$  (is) a right-angle—for it is equal to the internal and opposite (angle)  $ECB$  [Prop. 1.29]—the remaining (angle)  $EFG$  is thus half a right-angle [Prop. 1.32]. Thus, angle  $GEF$  [is] equal to  $EFG$ . So the side  $EG$  is also equal to the (side)  $GF$  [Prop. 1.6]. Again, since the angle at  $B$  is half a right-angle, and (angle)  $FDB$  (is) a right-angle—for again it is equal to the internal and opposite (angle)  $ECB$  [Prop. 1.29]—the remaining (angle)  $BFD$  is half a right-angle [Prop. 1.32]. Thus, the angle at  $B$  (is) equal to  $DFB$ . So the side  $FD$  is also equal to the side  $DB$  [Prop. 1.6]. And since  $AC$  is equal to  $CE$ , the (square) on  $AC$  (is) also equal to the (square) on  $CE$ . Thus, the (sum of the) squares on  $AC$  and  $CE$  is double the (square) on  $AC$ . And the square on  $EA$  is equal to the (sum of the) squares on  $AC$  and  $CE$ . For angle  $ACE$  (is) a right-angle [Prop. 1.47]. Thus, the (square) on  $EA$  is double the (square) on  $AC$ . Again, since  $EG$  is equal to  $GF$ , the (square) on  $EG$  (is) also equal to the (square) on  $GF$ . Thus, the (sum of the squares) on  $EG$  and  $GF$  is double the square on  $GF$ . And the square on  $EF$  is equal to the (sum of the) squares on  $EG$  and  $GF$  [Prop. 1.47]. Thus, the square on  $EF$  is double the (square) on  $GF$ . And  $GF$  (is) equal to  $CD$  [Prop. 1.34]. Thus, the (square) on  $EF$  is double the (square) on  $CD$ . And the (square) on  $EA$  is also double the (square) on  $AC$ . Thus, the (sum of the) squares on  $AE$  and  $EF$  is double the (sum of the) squares on  $AC$  and  $CD$ . And the square on  $AF$  is equal to the (sum of the squares) on  $AE$  and  $EF$ . For the angle  $AEF$  is a right-angle [Prop. 1.47]. Thus, the square on  $AF$  is double the (sum of the squares) on  $AC$  and  $CD$ . And the (sum of the squares) on  $AD$  and  $DF$  (is) equal to the (square) on  $AF$ . For the angle at  $D$  is a right-angle [Prop. 1.47]. Thus, the (sum of the squares) on  $AD$  and  $DF$  is double the (sum of the) squares on  $AC$  and  $CD$ . And  $DF$  (is) equal to  $DB$ . Thus, the (sum of the) squares on  $AD$  and  $DB$  is double the (sum of the) squares on  $AC$  and  $CD$ .

Thus, if a straight-line is cut into equal and unequal (pieces) then the (sum of the) squares on the unequal pieces of the whole (straight-line) is double the (sum of the) square on half (the straight-line) and (the square) on the (difference) between the (equal and unequal) pieces. (Which is) the very thing it was required to show.

† This proposition is a geometric version of the algebraic identity:  $a^2 + b^2 = 2[(a+b)/2]^2 + [(a+b)/2 - b]^2$ .

ι'.

Ἐάν εὐθεῖα γραμμὴ τμηθῆ διχα, προστεθῆ δέ τις αὐτῇ εὐθεῖα ἐπ' εὐθείας, τὸ ἀπὸ τῆς ὅλης σὺν τῇ προσκειμένῃ καὶ τὸ ἀπὸ τῆς προσκειμένης τὰ συναμφότερα τετράγωνα διπλάσιά ἐστι τοῦ τε ἀπὸ τῆς ἡμισείας καὶ τοῦ ἀπὸ τῆς συγκειμένης ἕκ τε τῆς ἡμισείας καὶ τῆς προσκειμένης ὡς ἀπὸ μιᾶς ἀναγραφέντος τετραγώνων.




Εὐθεῖα γάρ τις ἡ  $AB$  τεμηθῶ διχα κατὰ τὸ  $\Gamma$ , προσκεισθῶ δέ τις αὐτῇ εὐθεῖα ἐπ' εὐθείας ἡ  $BD$ . λέγω, ὅτι τὰ ἀπὸ τῶν  $A\Delta$ ,  $\Delta B$  τετράγωνα διπλάσιά ἐστι τῶν ἀπὸ τῶν  $A\Gamma$ ,  $\Gamma\Delta$  τετραγώνων.

Ἦχθω γὰρ ἀπὸ τοῦ  $\Gamma$  σημείου τῆς  $AB$  πρὸς ὀρθὰς ἡ  $GE$ , καὶ κείσθω ἴση ἑκάτερα τῶν  $A\Gamma$ ,  $\Gamma B$ , καὶ ἐπεζεύχθωσαν αἱ  $EA$ ,  $EB$ . καὶ διὰ μὲν τοῦ  $E$  τῆς  $AD$  παράλληλος ἤχθω ἡ  $EZ$ , διὰ δὲ τοῦ  $\Delta$  τῆς  $GE$  παράλληλος ἤχθω ἡ  $Z\Delta$ . καὶ ἐπεὶ εἰς παραλλήλους εὐθείας τὰς  $EG$ ,  $Z\Delta$  εὐθεῖά τις ἐνέπεσεν ἡ  $EZ$ , αἱ ὑπὸ  $GEZ$ ,  $EZ\Delta$  ἄρα δυσὶν ὀρθαῖς ἴσαι εἰσίν· αἱ ἄρα ὑπὸ  $ZEB$ ,  $EZ\Delta$  δύο ὀρθῶν ἐλάσσονές εἰσιν· αἱ δὲ ἀπ' ἐλασσόνων ἡ δύο ὀρθῶν ἐκβαλλόμεναι συμπίπτουσιν· αἱ ἄρα  $EB$ ,  $Z\Delta$  ἐκβαλλόμεναι ἐπὶ τὰ  $B$ ,  $\Delta$  μέρη συμπεσοῦνται. ἐκβεβλήσθωσαν καὶ συμπιπέτωσαν κατὰ τὸ  $H$ , καὶ ἐπεζεύχθω ἡ  $AH$ . καὶ ἐπεὶ ἴση ἐστὶν ἡ  $A\Gamma$  τῆς  $GE$ , ἴση ἐστὶ καὶ γωνία ἡ ὑπὸ  $EAG$  τῆς ὑπὸ  $AEG$ . καὶ ὀρθὴ ἡ πρὸς τῷ  $\Gamma$ . ἡμίσεια ἄρα ὀρθῆς [ἐστὶν] ἑκάτερα τῶν ὑπὸ  $EAG$ ,  $AEG$ . διὰ τὰ αὐτὰ δὴ καὶ ἑκάτερα τῶν ὑπὸ  $GEB$ ,  $EBG$  ἡμίσειά ἐστὶν ὀρθῆς· ὀρθὴ ἄρα ἐστὶν ἡ ὑπὸ  $AEB$ . καὶ ἐπεὶ ἡμίσεια ὀρθῆς ἐστὶν ἡ ὑπὸ  $EBG$ , ἡμίσεια ἄρα ὀρθῆς καὶ ἡ ὑπὸ  $\Delta BH$ . ἔστι δὲ καὶ ἡ ὑπὸ  $B\Delta H$  ὀρθή· ἴση γάρ ἐστι τῆς ὑπὸ  $\Delta GE$ . ἐναλλάξ γάρ· λοιπὴ ἄρα ἡ ὑπὸ  $\Delta HB$  ἡμίσειά ἐστὶν ὀρθῆς· ἡ ἄρα ὑπὸ  $\Delta HB$  τῆς ὑπὸ  $\Delta BH$  ἐστὶν ἴση· ὥστε καὶ πλευρὰ ἡ  $B\Delta$  πλευρᾶ τῆς  $H\Delta$  ἐστὶν ἴση. πάλιν, ἐπεὶ ἡ ὑπὸ  $EHZ$  ἡμίσειά ἐστὶν ὀρθῆς, ὀρθὴ δὲ ἡ πρὸς τῷ  $Z$ . ἴση γάρ ἐστι τῆς ἀπεναντίον τῆς πρὸς τῷ  $\Gamma$ . λοιπὴ ἄρα ἡ ὑπὸ  $ZEH$  ἡμίσειά ἐστὶν ὀρθῆς· ἴση ἄρα ἡ ὑπὸ  $EHZ$  γωνία τῆς ὑπὸ  $ZEH$ . ὥστε καὶ πλευρὰ ἡ  $HZ$  πλευρᾶ τῆς  $EZ$  ἐστὶν ἴση. καὶ ἐπεὶ [ἴση ἐστὶν ἡ  $EG$  τῆς  $\Gamma A$ ], ἴσον ἐστὶ [καὶ] τὸ ἀπὸ τῆς  $EG$  τετραγώνων τῷ ἀπὸ τῆς  $\Gamma A$

Proposition 10†

If a straight-line is cut in half, and any straight-line added to it straight-on, then the sum of the square on the whole (straight-line) with the (straight-line) having been added, and the (square) on the (straight-line) having been added, is double the (sum of the square) on half (the straight-line), and the square described on the sum of half (the straight-line) and (straight-line) having been added, as on one (complete straight-line).



For let any straight-line  $AB$  have been cut in half at (point)  $C$ , and let any straight-line  $BD$  have been added to it straight-on. I say that the (sum of the) squares on  $AD$  and  $DB$  is double the (sum of the) squares on  $AC$  and  $CD$ .

For let  $CE$  have been drawn from point  $C$ , at right-angles to  $AB$  [Prop. 1.11], and let it be made equal to each of  $AC$  and  $CB$  [Prop. 1.3], and let  $EA$  and  $EB$  have been joined. And let  $EF$  have been drawn through  $E$ , parallel to  $AD$  [Prop. 1.31], and let  $FD$  have been drawn through  $D$ , parallel to  $CE$  [Prop. 1.31]. And since some straight-line  $EF$  falls across the parallel straight-lines  $EC$  and  $FD$ , the (internal angles)  $CEF$  and  $EFD$  are thus equal to two right-angles [Prop. 1.29]. Thus,  $FEB$  and  $EFD$  are less than two right-angles. And (straight-lines) produced from (internal angles whose sum is) less than two right-angles meet together [Post. 5]. Thus, being produced in the direction of  $B$  and  $D$ , the (straight-lines)  $EB$  and  $FD$  will meet. Let them have been produced, and let them meet together at  $G$ , and let  $AG$  have been joined. And since  $AC$  is equal to  $CE$ , angle  $EAC$  is also equal to (angle)  $AEC$  [Prop. 1.5]. And the (angle) at  $C$  (is) a right-angle. Thus,  $EAC$  and  $AEC$  [are] each half a right-angle [Prop. 1.32]. So, for the same (reasons),  $CEB$  and  $EBC$  are also each half a right-angle. Thus, (angle)  $AEB$  is a right-angle. And since  $EBC$  is half a right-angle,  $DBG$  (is) thus also half a right-angle [Prop. 1.15]. And  $BDG$  is also a right-angle. For it is equal to  $DCE$ . For (they are) alternate (angles)

τετραγώνω· τὰ ἄρα ἀπὸ τῶν ΕΓ, ΓΑ τετράγωνα διπλάσιά ἐστι τοῦ ἀπὸ τῆς ΓΑ τετραγώνου. τοῖς δὲ ἀπὸ τῶν ΕΓ, ΓΑ ἴσον ἐστὶ τὸ ἀπὸ τῆς ΕΑ· τὸ ἄρα ἀπὸ τῆς ΕΑ τετράγωνον διπλάσιόν ἐστι τοῦ ἀπὸ τῆς ΑΓ τετραγώνου. πάλιν, ἐπεὶ ἴση ἐστὶν ἡ ΖΗ τῆς ΕΖ, ἴσον ἐστὶ καὶ τὸ ἀπὸ τῆς ΖΗ τῷ ἀπὸ τῆς ΖΕ· τὰ ἄρα ἀπὸ τῶν ΗΖ, ΖΕ διπλάσιά ἐστι τοῦ ἀπὸ τῆς ΕΖ. τοῖς δὲ ἀπὸ τῶν ΗΖ, ΖΕ ἴσον ἐστὶ τὸ ἀπὸ τῆς ΕΗ· τὸ ἄρα ἀπὸ τῆς ΕΗ διπλάσιόν ἐστι τοῦ ἀπὸ τῆς ΕΖ. ἴση δὲ ἡ ΕΖ τῆς ΓΔ· τὸ ἄρα ἀπὸ τῆς ΕΗ τετράγωνον διπλάσιόν ἐστι τοῦ ἀπὸ τῆς ΓΔ. ἐδείχθη δὲ καὶ τὸ ἀπὸ τῆς ΕΑ διπλάσιον τοῦ ἀπὸ τῆς ΑΓ· τὰ ἄρα ἀπὸ τῶν ΑΕ, ΕΗ τετράγωνα διπλάσιά ἐστι τῶν ἀπὸ τῶν ΑΓ, ΓΔ τετραγώνων. τοῖς δὲ ἀπὸ τῶν ΑΕ, ΕΗ τετραγώνοις ἴσον ἐστὶ τὸ ἀπὸ τῆς ΑΗ τετράγωνον· τὸ ἄρα ἀπὸ τῆς ΑΗ διπλάσιόν ἐστι τῶν ἀπὸ τῶν ΑΓ, ΓΔ. τῷ δὲ ἀπὸ τῆς ΑΗ ἴσα ἐστὶ τὰ ἀπὸ τῶν ΑΔ, ΔΗ· τὰ ἄρα ἀπὸ τῶν ΑΔ, ΔΗ [τετράγωνα] διπλάσιά ἐστι τῶν ἀπὸ τῶν ΑΓ, ΓΔ [τετραγώνων]. ἴση δὲ ἡ ΔΗ τῆς ΔΒ· τὰ ἄρα ἀπὸ τῶν ΑΔ, ΔΒ [τετράγωνα] διπλάσιά ἐστι τῶν ἀπὸ τῶν ΑΓ, ΓΔ τετραγώνων.

Ἐὰν ἄρα εὐθεῖα γραμμὴ τμηθῆ διχα, προστεθῆ δὲ τις αὐτῇ εὐθεῖα ἐπ' εὐθείας, τὸ ἀπὸ τῆς ὅλης σὺν τῇ προσκειμένη καὶ τὸ ἀπὸ τῆς προσκειμένης τὰ συναμφοτέρα τετράγωνα διπλάσιά ἐστι τοῦ τε ἀπὸ τῆς ἡμισείας καὶ τοῦ ἀπὸ τῆς συγκειμένης ἕκ τε τῆς ἡμισείας καὶ τῆς προσκειμένης ὡς ἀπὸ μιᾶς ἀναγραφέντος τετραγώνου· ὅπερ εἶδει δεῖξαι.

[Prop. 1.29]. Thus, the remaining (angle)  $DGB$  is half a right-angle. Thus,  $DGB$  is equal to  $DBG$ . So side  $BD$  is also equal to side  $GD$  [Prop. 1.6]. Again, since  $EGF$  is half a right-angle, and the (angle) at  $F$  (is) a right-angle, for it is equal to the opposite (angle) at  $C$  [Prop. 1.34], the remaining (angle)  $FEG$  is thus half a right-angle. Thus, angle  $EGF$  (is) equal to  $FEG$ . So the side  $GF$  is also equal to the side  $EF$  [Prop. 1.6]. And since [ $EC$  is equal to  $CA$ ] the square on  $EC$  is [also] equal to the square on  $CA$ . Thus, the (sum of the) squares on  $EC$  and  $CA$  is double the square on  $CA$ . And the (square) on  $EA$  is equal to the (sum of the squares) on  $EC$  and  $CA$  [Prop. 1.47]. Thus, the square on  $EA$  is double the square on  $AC$ . Again, since  $FG$  is equal to  $EF$ , the (square) on  $FG$  is also equal to the (square) on  $FE$ . Thus, the (sum of the squares) on  $GF$  and  $FE$  is double the (square) on  $EF$ . And the (square) on  $EG$  is equal to the (sum of the squares) on  $GF$  and  $FE$  [Prop. 1.47]. Thus, the (square) on  $EG$  is double the (square) on  $EF$ . And  $EF$  (is) equal to  $CD$  [Prop. 1.34]. Thus, the square on  $EG$  is double the (square) on  $CD$ . But it was also shown that the (square) on  $EA$  (is) double the (square) on  $AC$ . Thus, the (sum of the) squares on  $AE$  and  $EG$  is double the (sum of the) squares on  $AC$  and  $CD$ . And the square on  $AG$  is equal to the (sum of the) squares on  $AE$  and  $EG$  [Prop. 1.47]. Thus, the (square) on  $AG$  is double the (sum of the squares) on  $AC$  and  $CD$ . And the (sum of the squares) on  $AD$  and  $DG$  is equal to the (square) on  $AG$  [Prop. 1.47]. Thus, the (sum of the) [squares] on  $AD$  and  $DG$  is double the (sum of the) [squares] on  $AC$  and  $CD$ . And  $DG$  (is) equal to  $DB$ . Thus, the (sum of the) [squares] on  $AD$  and  $DB$  is double the (sum of the) squares on  $AC$  and  $CD$ .

Thus, if a straight-line is cut in half, and any straight-line added to it straight-on, then the sum of the square on the whole (straight-line) with the (straight-line) having been added, and the (square) on the (straight-line) having been added, is double the (sum of the square) on half (the straight-line), and the square described on the sum of half (the straight-line) and (straight-line) having been added, as on one (complete straight-line). (Which is) the very thing it was required to show.



† This proposition is a geometric version of the algebraic identity:  $(2a + b)^2 + b^2 = 2[a^2 + (a + b)^2]$ .

ια'.

### Proposition 11<sup>†</sup>

Τὴν δοθεῖσαν εὐθεῖαν τεμεῖν ὥστε τὸ ὑπὸ τῆς ὅλης καὶ τοῦ ἐτέρου τῶν τμημάτων περιεχόμενον ὀρθογώνιον ἴσον εἶναι τῷ ἀπὸ τοῦ λοιποῦ τμήματος τετραγώνω.

To cut a given straight-line such that the rectangle contained by the whole (straight-line), and one of the pieces (of the straight-line), is equal to the square on the remaining piece.



Ἐστω ἡ δοθεῖσα εὐθεῖα ἡ  $AB$ . δεῖ δὴ τὴν  $AB$  τεμεῖν ὥστε τὸ ὑπὸ τῆς ὅλης καὶ τοῦ ἑτέρου τῶν τμημάτων περιεχόμενον ὀρθογώνιον ἴσον εἶναι τῷ ἀπὸ τοῦ λοιποῦ τμήματος τετραγώνῳ.

Ἀναγεγράφθω γὰρ ἀπὸ τῆς  $AB$  τετράγωνον τὸ  $AB\Delta\Gamma$ , καὶ τεμηθῶ ἡ  $AG$  δίχα κατὰ τὸ  $E$  σημεῖον, καὶ ἐπεζεύχθω ἡ  $BE$ , καὶ διήχθω ἡ  $GA$  ἐπὶ τὸ  $Z$ , καὶ κείσθω τῇ  $BE$  ἴση ἡ  $EZ$ , καὶ ἀναγεγράφθω ἀπὸ τῆς  $AZ$  τετράγωνον τὸ  $Z\Theta$ , καὶ διήχθω ἡ  $H\Theta$  ἐπὶ τὸ  $K$ . λέγω, ὅτι ἡ  $AB$  τέτμηται κατὰ τὸ  $\Theta$ , ὥστε τὸ ὑπὸ τῶν  $AB$ ,  $B\Theta$  περιεχόμενον ὀρθογώνιον ἴσον ποιεῖν τῷ ἀπὸ τῆς  $A\Theta$  τετραγώνῳ.

Ἐπεὶ γὰρ εὐθεῖα ἡ  $AG$  τέτμηται δίχα κατὰ τὸ  $E$ , πρόσκειται δὲ αὐτῇ ἡ  $ZA$ , τὸ ἄρα ὑπὸ τῶν  $\Gamma Z$ ,  $ZA$  περιεχόμενον ὀρθογώνιον μετὰ τοῦ ἀπὸ τῆς  $AE$  τετραγώνου ἴσον ἐστὶ τῷ ἀπὸ τῆς  $EZ$  τετραγώνῳ. ἴση δὲ ἡ  $EZ$  τῇ  $EB$ . τὸ ἄρα ὑπὸ τῶν  $\Gamma Z$ ,  $ZA$  μετὰ τοῦ ἀπὸ τῆς  $AE$  ἴσον ἐστὶ τῷ ἀπὸ  $EB$ . ἀλλὰ τῷ ἀπὸ  $EB$  ἴσα ἐστὶ τὰ ἀπὸ τῶν  $BA$ ,  $AE$ . ὀρθὴ γὰρ ἡ πρὸς τῷ  $A$  γωνία. τὸ ἄρα ὑπὸ τῶν  $\Gamma Z$ ,  $ZA$  μετὰ τοῦ ἀπὸ τῆς  $AE$  ἴσον ἐστὶ τοῖς ἀπὸ τῶν  $BA$ ,  $AE$ . κοινὸν ἀφῆρήσθω τὸ ἀπὸ τῆς  $AE$ . λοιπὸν ἄρα τὸ ὑπὸ τῶν  $\Gamma Z$ ,  $ZA$  περιεχόμενον ὀρθογώνιον ἴσον ἐστὶ τῷ ἀπὸ τῆς  $AB$  τετραγώνῳ. καὶ ἐστὶ τὸ μὲν ὑπὸ τῶν  $\Gamma Z$ ,  $ZA$  τὸ  $ZK$ . ἴση γὰρ ἡ  $AZ$  τῇ  $ZH$ . τὸ δὲ ἀπὸ τῆς  $AB$  τὸ  $A\Delta$ . τὸ ἄρα  $ZK$  ἴσον ἐστὶ τῷ  $A\Delta$ . κοινὸν ἀφῆρήσθω τὸ  $AK$ . λοιπὸν ἄρα τὸ  $Z\Theta$  τῷ  $\Theta\Delta$  ἴσον ἐστίν. καὶ ἐστὶ τὸ μὲν  $\Theta\Delta$  τὸ ὑπὸ τῶν  $AB$ ,  $B\Theta$ . ἴση γὰρ ἡ  $AB$  τῇ  $B\Delta$ . τὸ δὲ  $Z\Theta$  τὸ ἀπὸ τῆς  $A\Theta$ . τὸ ἄρα ὑπὸ τῶν  $AB$ ,  $B\Theta$  περιεχόμενον ὀρθογώνιον ἴσον ἐστὶ τῷ ἀπὸ  $\Theta A$  τετραγώνῳ.

Ἡ ἄρα δοθεῖσα εὐθεῖα ἡ  $AB$  τέτμηται κατὰ τὸ  $\Theta$  ὥστε τὸ ὑπὸ τῶν  $AB$ ,  $B\Theta$  περιεχόμενον ὀρθογώνιον ἴσον ποιεῖν τῷ ἀπὸ τῆς  $\Theta A$  τετραγώνῳ. ὅπερ ἔδει ποιῆσαι.

Let  $AB$  be the given straight-line. So it is required to cut  $AB$  such that the rectangle contained by the whole (straight-line), and one of the pieces (of the straight-line), is equal to the square on the remaining piece.

For let the square  $ABDC$  have been described on  $AB$  [Prop. 1.46], and let  $AC$  have been cut in half at point  $E$  [Prop. 1.10], and let  $BE$  have been joined. And let  $CA$  have been drawn through to (point)  $F$ , and let  $EF$  be made equal to  $BE$  [Prop. 1.3]. And let the square  $FH$  have been described on  $AF$  [Prop. 1.46], and let  $GH$  have been drawn through to (point)  $K$ . I say that  $AB$  has been cut at  $H$  such as to make the rectangle contained by  $AB$  and  $BH$  equal to the square on  $AH$ .

For since the straight-line  $AC$  has been cut in half at  $E$ , and  $FA$  has been added to it, the rectangle contained by  $CF$  and  $FA$ , plus the square on  $AE$ , is thus equal to the square on  $EF$  [Prop. 2.6]. And  $EF$  (is) equal to  $EB$ . Thus, the (rectangle contained) by  $CF$  and  $FA$ , plus the (square) on  $AE$ , is equal to the (square) on  $EB$ . But, the (sum of the squares) on  $BA$  and  $AE$  is equal to the (square) on  $EB$ . For the angle at  $A$  (is) a right-angle [Prop. 1.47]. Thus, the (rectangle contained) by  $CF$  and  $FA$ , plus the (square) on  $AE$ , is equal to the (sum of the squares) on  $BA$  and  $AE$ . Let the square on  $AE$  have been subtracted from both. Thus, the remaining rectangle contained by  $CF$  and  $FA$  is equal to the square on  $AB$ . And  $FK$  is the (rectangle contained) by  $CF$  and  $FA$ . For  $AF$  (is) equal to  $FG$ . And  $AD$  (is) the (square) on  $AB$ . Thus, the (rectangle)  $FK$  is equal to the (square)  $AD$ . Let (rectangle)  $AK$  have been subtracted from both. Thus, the remaining (square)  $FH$  is equal to the (rectangle)  $HD$ . And  $HD$  is the (rectangle contained) by  $AB$  and  $BH$ . For  $AB$  (is) equal to  $BD$ . And  $FH$  (is) the (square) on  $AH$ . Thus, the rectangle contained by  $AB$


and  $BH$  is equal to the square on  $HA$ .

Thus, the given straight-line  $AB$  has been cut at (point)  $H$  such as to make the rectangle contained by  $AB$  and  $BH$  equal to the square on  $HA$ . (Which is) the very thing it was required to do.

† This manner of cutting a straight-line—so that the ratio of the whole to the larger piece is equal to the ratio of the larger to the smaller piece—is sometimes called the “Golden Section”.

ιβ'.

Ἐν τοῖς ἀμβλυγωνίοις τριγώνοις τὸ ἀπὸ τῆς τῆν ἀμβλείαν γωνίαν ὑποτείνουσας πλευρᾶς τετράγωνον μεῖζόν ἐστὶ τῶν ἀπὸ τῶν τῆν ἀμβλείαν γωνίαν περιεχουσῶν πλευρῶν τετραγώνων τῷ περιεχομένῳ δις ὑπὸ τε μιᾶς τῶν περὶ τῆν ἀμβλείαν γωνίαν, ἐφ' ἣν ἡ κάθετος πίπτει, καὶ τῆς ἀπολαμβανομένης ἐκτὸς ὑπὸ τῆς καθέτου πρὸς τῇ ἀμβλείᾳ γωνία.




Ἐστω ἀμβλυγώνιον τρίγωνον τὸ  $AB\Gamma$  ἀμβλείαν ἔχον τὴν ὑπὸ  $BAG$ , καὶ ἤχθω ἀπὸ τοῦ  $B$  σημείου ἐπὶ τὴν  $GA$  ἐκβληθεῖσαν κάθετος ἡ  $BD$ . λέγω, ὅτι τὸ ἀπὸ τῆς  $B\Gamma$  τετράγωνον μεῖζόν ἐστὶ τῶν ἀπὸ τῶν  $BA$ ,  $A\Gamma$  τετραγώνων τῷ δις ὑπὸ τῶν  $GA$ ,  $A\Delta$  περιεχομένῳ ὀρθογωνίῳ.

Ἐπεὶ γὰρ εὐθεῖα ἡ  $\Gamma\Delta$  τέμνεται, ὡς ἔτυχεν, κατὰ τὸ  $A$  σημεῖον, τὸ ἄρα ἀπὸ τῆς  $\Delta\Gamma$  ἴσον ἐστὶ τοῖς ἀπὸ τῶν  $GA$ ,  $A\Delta$  τετραγώνοις καὶ τῷ δις ὑπὸ τῶν  $GA$ ,  $A\Delta$  περιεχομένῳ ὀρθογωνίῳ. κοινὸν προσκείσθω τὸ ἀπὸ τῆς  $\Delta B$ : τὰ ἄρα ἀπὸ τῶν  $\Gamma\Delta$ ,  $\Delta B$  ἴσα ἐστὶ τοῖς τε ἀπὸ τῶν  $GA$ ,  $A\Delta$ ,  $\Delta B$  τετραγώνοις καὶ τῷ δις ὑπὸ τῶν  $GA$ ,  $A\Delta$  [περιεχομένῳ ὀρθογωνίῳ]. ἀλλὰ τοῖς μὲν ἀπὸ τῶν  $\Gamma\Delta$ ,  $\Delta B$  ἴσον ἐστὶ τὸ ἀπὸ τῆς  $GB$ : ὀρθὴ γὰρ ἡ πρὸς τῷ  $\Delta$  γωνία: τοῖς δὲ ἀπὸ τῶν  $A\Delta$ ,  $\Delta B$  ἴσον τὸ ἀπὸ τῆς  $AB$ : τὸ ἄρα ἀπὸ τῆς  $GB$  τετράγωνον ἴσον ἐστὶ τοῖς τε ἀπὸ τῶν  $GA$ ,  $AB$  τετραγώνοις καὶ τῷ δις ὑπὸ τῶν  $GA$ ,  $A\Delta$  περιεχομένῳ ὀρθογωνίῳ· ὥστε τὸ ἀπὸ τῆς  $GB$  τετράγωνον τῶν ἀπὸ τῶν  $GA$ ,  $AB$  τετραγώνων μεῖζόν ἐστὶ τῷ δις ὑπὸ τῶν  $GA$ ,  $A\Delta$  περιεχομένῳ ὀρθογωνίῳ.

Ἐν ἄρα τοῖς ἀμβλυγωνίοις τριγώνοις τὸ ἀπὸ τῆς τῆν ἀμβλείαν γωνίαν ὑποτείνουσας πλευρᾶς τετράγωνον μεῖζόν ἐστὶ τῶν ἀπὸ τῶν τῆν ἀμβλείαν γωνίαν περιεχουσῶν

Proposition 12†

In obtuse-angled triangles, the square on the side subtending the obtuse angle is greater than the (sum of the) squares on the sides containing the obtuse angle by twice the (rectangle) contained by one of the sides around the obtuse angle, to which a perpendicular (straight-line) falls, and the (straight-line) cut off outside (the triangle) by the perpendicular (straight-line) towards the obtuse angle.



Let  $ABC$  be an obtuse-angled triangle, having the angle  $BAC$  obtuse. And let  $BD$  be drawn from point  $B$ , perpendicular to  $CA$  produced [Prop. 1.12]. I say that the square on  $BC$  is greater than the (sum of the) squares on  $BA$  and  $AC$ , by twice the rectangle contained by  $CA$  and  $AD$ .

For since the straight-line  $CD$  has been cut, at random, at point  $A$ , the (square) on  $DC$  is thus equal to the (sum of the) squares on  $CA$  and  $AD$ , and twice the rectangle contained by  $CA$  and  $AD$  [Prop. 2.4]. Let the (square) on  $DB$  have been added to both. Thus, the (sum of the squares) on  $CD$  and  $DB$  is equal to the (sum of the) squares on  $CA$ ,  $AD$ , and  $DB$ , and twice the [rectangle contained] by  $CA$  and  $AD$ . But, the (square) on  $CB$  is equal to the (sum of the squares) on  $CD$  and  $DB$ . For the angle at  $D$  (is) a right-angle [Prop. 1.47]. And the (square) on  $AB$  (is) equal to the (sum of the squares) on  $AD$  and  $DB$  [Prop. 1.47]. Thus, the square on  $CB$  is equal to the (sum of the) squares on  $CA$  and  $AB$ , and twice the rectangle contained by  $CA$  and  $AD$ . So the square on  $CB$  is greater than the (sum of the) squares on



πλευρῶν τετραγῶνων τῷ περιχομένῳ δις ὑπό τε μιᾶς τῶν περι τὴν ἀμβλείαν γωνίαν, ἐφ' ἣν ἡ κάθετος πίπτει, καὶ τῆς ἀπολαμβανομένης ἐκτὸς ὑπὸ τῆς καθέτου πρὸς τῇ ἀμβλείᾳ γωνίᾳ· ὅπερ ἔδει δεῖξαι.

$CA$  and  $AB$  by twice the rectangle contained by  $CA$  and  $AD$ .

Thus, in obtuse-angled triangles, the square on the side subtending the obtuse angle is greater than the (sum of the) squares on the sides containing the obtuse angle by twice the (rectangle) contained by one of the sides around the obtuse angle, to which a perpendicular (straight-line) falls, and the (straight-line) cut off outside (the triangle) by the perpendicular (straight-line) towards the obtuse angle. (Which is) the very thing it was required to show.



† This proposition is equivalent to the well-known cosine formula:  $BC^2 = AB^2 + AC^2 - 2 AB AC \cos BAC$ , since  $\cos BAC = -AD/AB$ .

ιγ'.

Ἐν τοῖς ὀξυγωνίοις τριγώνοις τὸ ἀπὸ τῆς τὴν ὀξείαν γωνίαν ὑποτείνουσας πλευρᾶς τετράγωνον ἔλαττόν ἐστι τῶν ἀπὸ τῶν τὴν ὀξείαν γωνίαν περιεχουσῶν πλευρῶν τετραγῶνων τῷ περιχομένῳ δις ὑπό τε μιᾶς τῶν περι τὴν ὀξείαν γωνίαν, ἐφ' ἣν ἡ κάθετος πίπτει, καὶ τῆς ἀπολαμβανομένης ἐντὸς ὑπὸ τῆς καθέτου πρὸς τῇ ὀξείᾳ γωνίᾳ.

Proposition 13†

In acute-angled triangles, the square on the side subtending the acute angle is less than the (sum of the) squares on the sides containing the acute angle by twice the (rectangle) contained by one of the sides around the acute angle, to which a perpendicular (straight-line) falls, and the (straight-line) cut off inside (the triangle) by the perpendicular (straight-line) towards the acute angle.



Ἐστω ὀξυγώνιον τρίγωνον τὸ  $AB\Gamma$  ὀξείαν ἔχον τὴν πρὸς τῷ  $B$  γωνίαν, καὶ ἤχθω ἀπὸ τοῦ  $A$  σημείου ἐπὶ τὴν  $B\Gamma$  κάθετος ἡ  $AD$ . λέγω, ὅτι τὸ ἀπὸ τῆς  $AG$  τετράγωνον ἔλαττόν ἐστι τῶν ἀπὸ τῶν  $GB$ ,  $BA$  τετραγῶνων τῷ δις ὑπὸ τῶν  $GB$ ,  $BD$  περιχομένῳ ὀρθογωνίῳ.

Let  $ABC$  be an acute-angled triangle, having the angle at (point)  $B$  acute. And let  $AD$  have been drawn from point  $A$ , perpendicular to  $BC$  [Prop. 1.12]. I say that the square on  $AC$  is less than the (sum of the) squares on  $CB$  and  $BA$ , by twice the rectangle contained by  $CB$  and  $BD$ .

Ἐπεὶ γὰρ εὐθεῖα ἡ  $GB$  τέμνεται, ὡς ἔτυχεν, κατὰ τὸ  $\Delta$ , τὰ ἄρα ἀπὸ τῶν  $GB$ ,  $BD$  τετράγωνα ἴσα ἐστὶ τῷ τε δις ὑπὸ τῶν  $GB$ ,  $BD$  περιχομένῳ ὀρθογωνίῳ καὶ τῷ ἀπὸ τῆς  $\Delta\Gamma$  τετραγῶνῳ. κοινὸν προσκείσθω τὸ ἀπὸ τῆς  $\Delta A$  τετράγωνον· τὰ ἄρα ἀπὸ τῶν  $GB$ ,  $BD$ ,  $\Delta A$  τετράγωνα ἴσα ἐστὶ τῷ τε δις ὑπὸ τῶν  $GB$ ,  $BD$  περιχομένῳ ὀρθογωνίῳ καὶ τοῖς ἀπὸ τῶν  $A\Delta$ ,  $\Delta\Gamma$  τετραγῶνις. ἀλλὰ τοῖς μὲν ἀπὸ τῶν  $B\Delta$ ,  $\Delta A$  ἴσον τὸ ἀπὸ τῆς  $AB$ · ὀρθὴ γὰρ ἡ πρὸς τῷ  $\Delta$  γωνία· τοῖς δὲ ἀπὸ τῶν  $A\Delta$ ,  $\Delta\Gamma$  ἴσον τὸ ἀπὸ τῆς  $AG$ · τὰ ἄρα ἀπὸ τῶν  $GB$ ,  $BA$  ἴσα ἐστὶ τῷ τε ἀπὸ τῆς  $AG$  καὶ τῷ δις ὑπὸ τῶν  $GB$ ,  $BD$ · ὥστε μόνον τὸ ἀπὸ τῆς  $AG$  ἔλαττόν ἐστι

For since the straight-line  $CB$  has been cut, at random, at (point)  $D$ , the (sum of the) squares on  $CB$  and  $BD$  is thus equal to twice the rectangle contained by  $CB$  and  $BD$ , and the square on  $DC$  [Prop. 2.7]. Let the square on  $DA$  have been added to both. Thus, the (sum of the) squares on  $CB$ ,  $BD$ , and  $DA$  is equal to twice the rectangle contained by  $CB$  and  $BD$ , and the (sum of the) squares on  $AD$  and  $DC$ . But, the (square) on  $AB$  (is) equal to the (sum of the squares) on  $BD$  and  $DA$ . For the angle at (point)  $D$  is a right-angle [Prop. 1.47].

τῶν ἀπὸ τῶν ΓΒ, ΒΑ τετραγώνων τῶ δις ὑπὸ τῶν ΓΒ, ΒΔ περιεχομένῳ ὀρθογώνιῳ.

Ἐν ἄρα τοῖς ὀξυγωνίοις τριγώνοις τὸ ἀπὸ τῆς τὴν ὀξείαν γωνίαν ὑποτείνουσας πλευρᾶς τετραγώνον ἔλαττόν ἐστι τῶν ἀπὸ τῶν τὴν ὀξείαν γωνίαν περιεχουσῶν πλευρῶν τετραγώνων τῶ περιεχομένῳ δις ὑπὸ τε μιᾶς τῶν περὶ τὴν ὀξείαν γωνίαν, ἐφ' ἣν ἡ κάθετος πίπτει, καὶ τῆς ἀπολαμβανομένης ἐντὸς ὑπὸ τῆς καθέτου πρὸς τῇ ὀξείᾳ γωνίᾳ: ὅπερ ἔδει δεῖξαι.

And the (square) on  $AC$  (is) equal to the (sum of the squares) on  $AD$  and  $DC$  [Prop. 1.47]. Thus, the (sum of the squares) on  $CB$  and  $BA$  is equal to the (square) on  $AC$ , and twice the (rectangle contained) by  $CB$  and  $BD$ . So the (square) on  $AC$  alone is less than the (sum of the) squares on  $CB$  and  $BA$  by twice the rectangle contained by  $CB$  and  $BD$ .

Thus, in acute-angled triangles, the square on the side subtending the acute angle is less than the (sum of the) squares on the sides containing the acute angle by twice the (rectangle) contained by one of the sides around the acute angle, to which a perpendicular (straight-line) falls, and the (straight-line) cut off inside (the triangle) by the perpendicular (straight-line) towards the acute angle. (Which is) the very thing it was required to show.



† This proposition is equivalent to the well-known cosine formula:  $AC^2 = AB^2 + BC^2 - 2 AB BC \cos ABC$ , since  $\cos ABC = BD/AB$ .

ιδ'.

Proposition 14

Τῶ δοθέντι εὐθύγραμμῳ ἴσον τετράγωνον συστήσασθαι.

To construct a square equal to a given rectilinear figure.



Ἐστω τὸ δοθὲν εὐθύγραμμον τὸ Α: δεῖ δὴ τῶ Α εὐθύγραμμῳ ἴσον τετράγωνον συστήσασθαι.

Let  $A$  be the given rectilinear figure. So it is required to construct a square equal to the rectilinear figure  $A$ .

Συνεστάτω γάρ τῶ Α εὐθύγραμμῳ ἴσον παραλληλόγραμμον ὀρθογώνιον τὸ ΒΔ: εἰ μὲν οὖν ἴση ἐστὶν ἡ ΒΕ τῇ ΕΔ, γεγονόςς ἂν εἴη τὸ ἐπιταχθέν. συνέσταται γάρ τῶ Α εὐθύγραμμῳ ἴσον τετράγωνον τὸ ΒΔ: εἰ δὲ οὐ, μία τῶν ΒΕ, ΕΔ μείζων ἐστίν. ἔστω μείζων ἡ ΒΕ, καὶ ἐκβεβλήσθω ἐπὶ τὸ Ζ, καὶ κείσθω τῇ ΕΔ ἴση ἡ ΕΖ, καὶ τετμήσθω ἡ ΒΖ δίχα κατὰ τὸ Η, καὶ κέντρῳ τῶ Η, διαστήματι δὲ ἐνὶ τῶν ΗΒ, ΗΖ ἡμικύκλιον γεγράφθω τὸ ΒΘΖ, καὶ ἐκβεβλήσθω ἡ ΔΕ ἐπὶ τὸ Θ, καὶ ἐπεξεύχθω ἡ ΗΘ.

For let the right-angled parallelogram  $BD$ , equal to the rectilinear figure  $A$ , have been constructed [Prop. 1.45]. Therefore, if  $BE$  is equal to  $ED$  then that (which) was prescribed has taken place. For the square  $BD$ , equal to the rectilinear figure  $A$ , has been constructed. And if not, then one of the (straight-lines)  $BE$  or  $ED$  is greater (than the other). Let  $BE$  be greater, and let it have been produced to  $F$ , and let  $EF$  be made equal to  $ED$  [Prop. 1.3]. And let  $BF$  have been cut in half at (point)  $G$  [Prop. 1.10]. And, with center  $G$ , and radius one of the (straight-lines)  $GB$  or  $GF$ , let the semi-circle  $BHF$  have been drawn. And let  $DE$  have been produced to  $H$ , and let  $GH$  have been joined.

Ἐπεὶ οὖν εὐθεία ἡ ΒΖ τέτμηται εἰς μὲν ἴσα κατὰ τὸ Η, εἰς δὲ ἄνισα κατὰ τὸ Ε, τὸ ἄρα ὑπὸ τῶν ΒΕ, ΕΖ περιεχόμενον ὀρθογώνιον μετὰ τοῦ ἀπὸ τῆς ΕΗ τετραγώνου ἴσον ἐστὶ τῶ ἀπὸ τῆς ΗΖ τετραγώνῳ. ἴση δὲ ἡ ΗΖ τῇ ΗΘ: τὸ ἄρα ὑπὸ τῶν ΒΕ, ΕΖ μετὰ τοῦ ἀπὸ τῆς ΗΕ ἴσον ἐστὶ τῶ ἀπὸ τῆς ΗΘ. τῶ δὲ ἀπὸ τῆς ΗΘ ἴσα ἐστὶ τὰ ἀπὸ τῶν ΘΕ, ΕΗ

Therefore, since the straight-line  $BF$  has been cut—equally at  $G$ , and unequally at  $E$ —the rectangle con-

τετράγωνα· τὸ ἄρα ὑπὸ τῶν  $BE$ ,  $EZ$  μετὰ τοῦ ἀπὸ  $HE$  ἴσα ἐστὶ τοῖς ἀπὸ τῶν  $ΘE$ ,  $EH$ . κοινὸν ἀφηρήσθω τὸ ἀπὸ τῆς  $HE$  τετράγωνον· λοιπὸν ἄρα τὸ ὑπὸ τῶν  $BE$ ,  $EZ$  περιεχόμενον ὀρθογώνιον ἴσον ἐστὶ τῷ ἀπὸ τῆς  $EΘ$  τετραγώνῳ. ἀλλὰ τὸ ὑπὸ τῶν  $BE$ ,  $EZ$  τὸ  $BΔ$  ἐστίν· ἴση γὰρ ἡ  $EZ$  τῇ  $EΔ$ · τὸ ἄρα  $BΔ$  παραλληλόγραμμον ἴσον ἐστὶ τῷ ἀπὸ τῆς  $ΘE$  τετραγώνῳ. ἴσον δὲ τὸ  $BΔ$  τῷ  $A$  εὐθύγραμμῳ. καὶ τὸ  $A$  ἄρα εὐθύγραμμον ἴσον ἐστὶ τῷ ἀπὸ τῆς  $EΘ$  ἀναγραφησομένῳ τετραγώνῳ.

Τῷ ἄρα δοθέντι εὐθύγραμμῳ τῷ  $A$  ἴσον τετράγωνον συνέσταται τὸ ἀπὸ τῆς  $EΘ$  ἀναγραφησόμενον· ὅπερ ἔδει ποιῆσαι.

tained by  $BE$  and  $EF$ , plus the square on  $EG$ , is thus equal to the square on  $GF$  [Prop. 2.5]. And  $GF$  (is) equal to  $GH$ . Thus, the (rectangle contained) by  $BE$  and  $EF$ , plus the (square) on  $GE$ , is equal to the (square) on  $GH$ . And the (sum of the) squares on  $HE$  and  $EG$  is equal to the (square) on  $GH$  [Prop. 1.47]. Thus, the (rectangle contained) by  $BE$  and  $EF$ , plus the (square) on  $GE$ , is equal to the (sum of the squares) on  $HE$  and  $EG$ . Let the square on  $GE$  have been taken from both. Thus, the remaining rectangle contained by  $BE$  and  $EF$  is equal to the square on  $EH$ . But,  $BD$  is the (rectangle contained) by  $BE$  and  $EF$ . For  $EF$  (is) equal to  $ED$ . Thus, the parallelogram  $BD$  is equal to the square on  $HE$ . And  $BD$  (is) equal to the rectilinear figure  $A$ . Thus, the rectilinear figure  $A$  is also equal to the square (which) can be described on  $EH$ .

Thus, a square—(namely), that (which) can be described on  $EH$ —has been constructed, equal to the given rectilinear figure  $A$ . (Which is) the very thing it was required to do.

