

ELEMENTS BOOK 6

Similar Figures

Ὅροι.


α'. Ὅμοια σχήματα εὐθύγραμμά ἐστιν, ὅσα τὰς τε γωνίας ἴσας ἔχει κατὰ μίαν καὶ τὰς περὶ τὰς ἴσας γωνίας πλευρὰς ἀνάλογον.

β'. Ἄκρον καὶ μέσον λόγον εὐθεῖα τετυμησθαι λέγεται, ὅταν ἢ ὡς ἡ ὅλη πρὸς τὸ μείζον τμήμα, οὕτως τὸ μείζον πρὸς τὸ ἔλαττον.

γ'. Ὑψος ἐστὶ πάντος σχήματος ἢ ἀπὸ τῆς κορυφῆς ἐπὶ τὴν βᾶσιν κάθετος ἀγομένη.

α'.

Τὰ τρίγωνα καὶ τὰ παραλληλόγραμμα τὰ ὑπὸ τὸ αὐτὸ ὕψος ὄντα πρὸς ἄλληλά ἐστιν ὡς αἱ βᾶσεις.



Ἐστω τρίγωνα μὲν τὰ ΑΒΓ, ΑΓΔ, παραλληλόγραμμα δὲ τὰ ΕΓ, ΓΖ ὑπὸ τὸ αὐτὸ ὕψος τὸ ΑΓ· λέγω, ὅτι ἐστὶν ὡς ἡ ΒΓ βᾶσις πρὸς τὴν ΓΔ βᾶσις, οὕτως τὸ ΑΒΓ τρίγωνον πρὸς τὸ ΑΓΔ τρίγωνον, καὶ τὸ ΕΓ παραλληλόγραμμον πρὸς τὸ ΓΖ παραλληλόγραμμον.

Ἐκβεβλήσθω γὰρ ἡ ΒΔ ἐφ' ἑκάτερα τὰ μέρη ἐπὶ τὰ Θ, Λ σημεῖα, καὶ κείσθωσαν τῇ μὲν ΒΓ βᾶσει ἴσαι [ὁσαιδηποτοῦν] αἱ ΒΗ, ΗΘ, τῇ δὲ ΓΔ βᾶσει ἴσαι ὁσαιδηποτοῦν αἱ ΔΚ, ΚΛ, καὶ ἐπεζεύχθωσαν αἱ ΑΗ, ΑΘ, ΑΚ, ΑΛ.

Καὶ ἐπεὶ ἴσαι εἰσὶν αἱ ΓΒ, ΒΗ, ΗΘ ἀλλήλαις, ἴσα ἐστὶ καὶ τὰ ΑΘΗ, ΑΗΒ, ΑΒΓ τρίγωνα ἀλλήλοις. ὁσαπλασίον ἔρα ἐστὶν ἡ ΘΓ βᾶσις τῆς ΒΓ βᾶσεως, τοσαυταπλασίον ἐστὶ καὶ τὸ ΑΘΓ τρίγωνον τοῦ ΑΒΓ τριγώνου. διὰ τὰ αὐτὰ δὴ ὁσαπλασίον ἐστὶν ἡ ΛΓ βᾶσις τῆς ΓΔ βᾶσεως, τοσαυταπλασίον ἐστὶ καὶ τὸ ΑΛΓ τρίγωνον τοῦ ΑΓΔ τριγώνου· καὶ εἰ ἴση ἐστὶν ἡ ΘΓ βᾶσις τῇ ΓΔ βᾶσει, ἴσον ἐστὶ καὶ τὸ ΑΘΓ τρίγωνον τῷ ΑΛΓ τριγώνῳ, καὶ εἰ ὑπερέχει ἡ ΘΓ βᾶσις τῆς ΓΔ βᾶσεως, ὑπερέχει καὶ τὸ ΑΘΓ τρίγωνον τοῦ ΑΛΓ τριγώνου, καὶ εἰ ἐλάσσων, ἔλασσον. τεσσάρων δὲ ὄντων μεγεθῶν δύο μὲν βᾶσεων τῶν ΒΓ, ΓΔ, δύο δὲ τριγώνων τῶν ΑΒΓ, ΑΓΔ εἴληπται ἰσάκως πολλαπλασία τῆς μὲν ΒΓ βᾶσεως καὶ τοῦ ΑΒΓ τριγώνου ἢ τε ΘΓ βᾶσις καὶ τὸ ΑΘΓ τρίγωνον, τῆς δὲ ΓΔ βᾶσεως καὶ τοῦ ΑΛΓ τριγώνου ἄλλα,

Definitions


1. Similar rectilinear figures are those (which) have (their) angles separately equal and the (corresponding) sides about the equal angles proportional.

2. A straight-line is said to have been cut in extreme and mean ratio when as the whole is to the greater segment so the greater (segment is) to the lesser.

3. The height of any figure is the (straight-line) drawn from the vertex perpendicular to the base.

Proposition 1[†]

Triangles and parallelograms which are of the same height are to one another as their bases.



Let ABC and ACD be triangles, and EC and CF parallelograms, of the same height AC . I say that as base BC is to base CD , so triangle ABC (is) to triangle ACD , and parallelogram EC to parallelogram CF .

For let the (straight-line) BD have been produced in each direction to points H and L , and let [any number] (of straight-lines) BG and GH be made equal to base BC , and any number (of straight-lines) DK and KL equal to base CD . And let AG , AH , AK , and AL have been joined.

And since CB , BG , and GH are equal to one another, triangles AHG , AGB , and ABC are also equal to one another [Prop. 1.38]. Thus, as many times as base HC is (divisible by) base BC , so many times is triangle AHC also (divisible) by triangle ABC . So, for the same (reasons), as many times as base LC is (divisible) by base CD , so many times is triangle ALC also (divisible) by triangle ACD . And if base HC is equal to base CL then triangle AHC is also equal to triangle ACL [Prop. 1.38]. And if base HC exceeds base CL then triangle AHC also exceeds triangle ACL .[‡] And if (HC is) less (than CL then AHC is also) less (than ACL). So, their being four magnitudes, two bases, BC and CD , and two trian-

ἂ ἔτυχεν, ἰσάκεις πολλαπλάσια ἢ τε $\Lambda\Gamma$ βάσις καὶ τὸ $\Lambda\Lambda\Gamma$ τρίγωνον· καὶ δέδεικται, ὅτι, εἰ ὑπερέχει ἡ $\Theta\Gamma$ βάσις τῆς $\Gamma\Lambda$ βάσεως, ὑπερέχει καὶ τὸ $\Lambda\Theta\Gamma$ τρίγωνον τοῦ $\Lambda\Lambda\Gamma$ τριγώνου, καὶ εἰ ἴση, ἴσον, καὶ εἰ ἔλασσων, ἔλασσον· ἔστιν ἄρα ὡς ἡ $B\Gamma$ βάσις πρὸς τὴν $\Gamma\Delta$ βάσιν, οὕτως τὸ $AB\Gamma$ τρίγωνον πρὸς τὸ $A\Gamma\Delta$ τρίγωνον.

Καὶ ἐπεὶ τοῦ μὲν $AB\Gamma$ τριγώνου διπλάσιόν ἐστι τὸ $E\Gamma$ παραλληλόγραμμον, τοῦ δὲ $A\Gamma\Delta$ τριγώνου διπλάσιόν ἐστι τὸ $Z\Gamma$ παραλληλόγραμμον, τὰ δὲ μέρη τοῖς ὡσαύτως πολλαπλασίοις τὸν αὐτὸν ἔχει λόγον, ἔστιν ἄρα ὡς τὸ $AB\Gamma$ τρίγωνον πρὸς τὸ $A\Gamma\Delta$ τρίγωνον, οὕτως τὸ $E\Gamma$ παραλληλόγραμμον πρὸς τὸ $Z\Gamma$ παραλληλόγραμμον. ἐπεὶ οὖν ἐδείχθη, ὡς μὲν ἡ $B\Gamma$ βάσις πρὸς τὴν $\Gamma\Delta$, οὕτως τὸ $AB\Gamma$ τρίγωνον πρὸς τὸ $A\Gamma\Delta$ τρίγωνον, ὡς δὲ τὸ $AB\Gamma$ τρίγωνον πρὸς τὸ $A\Gamma\Delta$ τρίγωνον, οὕτως τὸ $E\Gamma$ παραλληλόγραμμον πρὸς τὸ $Z\Gamma$ παραλληλόγραμμον, καὶ ὡς ἄρα ἡ $B\Gamma$ βάσις πρὸς τὴν $\Gamma\Delta$ βάσιν, οὕτως τὸ $E\Gamma$ παραλληλόγραμμον πρὸς τὸ $Z\Gamma$ παραλληλόγραμμον.

Τὰ ἄρα τρίγωνα καὶ τὰ παραλληλόγραμμα τὰ ὑπὸ τὸ αὐτὸ ὕψος ὄντα πρὸς ἄλληλά ἐστιν ὡς αἱ βάσεις· ὅπερ ἔδει δεῖξαι.

gles, ABC and ACD , equal multiples have been taken of base BC and triangle ABC —(namely), base HC and triangle AHC —and other random equal multiples of base CD and triangle ADC —(namely), base LC and triangle ALC . And it has been shown that if base HC exceeds base CL then triangle AHC also exceeds triangle ALC , and if (HC is) equal (to CL then AHC is also) equal (to ALC), and if (HC is) less (than CL then AHC is also) less (than ALC). Thus, as base BC is to base CD , so triangle ABC (is) to triangle ACD [Def. 5.5]. And since parallelogram EC is double triangle ABC , and parallelogram FC is double triangle ACD [Prop. 1.34], and parts have the same ratio as similar multiples [Prop. 5.15], thus as triangle ABC is to triangle ACD , so parallelogram EC (is) to parallelogram FC . In fact, since it was shown that as base BC (is) to CD , so triangle ABC (is) to triangle ACD , and as triangle ABC (is) to triangle ACD , so parallelogram EC (is) to parallelogram CF , thus, also, as base BC (is) to base CD , so parallelogram EC (is) to parallelogram FC [Prop. 5.11].


Thus, triangles and parallelograms which are of the same height are to one another as their bases. (Which is) the very thing it was required to show.

† As is easily demonstrated, this proposition holds even when the triangles, or parallelograms, do not share a common side, and/or are not right-angled.

‡ This is a straight-forward generalization of Prop. 1.38.

β'.


Ἐὰν τριγώνου παρὰ μίαν τῶν πλευρῶν ἀχθῆ τις εὐθεῖα, ἀνάλογον τεμεῖ τὰς τοῦ τριγώνου πλευράς· καὶ ἐὰν αἱ τοῦ τριγώνου πλευραὶ ἀνάλογον τμηθῶσιν, ἢ ἐπὶ τὰς τομὰς ἐπιζευγνυμένη εὐθεῖα παρὰ τὴν λοιπὴν ἔσται τοῦ τριγώνου πλευράν.



Τριγώνου γὰρ τοῦ $AB\Gamma$ παράλληλος μὲ τῶν πλευρῶν τῆ $B\Gamma$ ἤχθη ἡ ΔE · λέγω, ὅτι ἐστὶν ὡς ἡ $B\Delta$ πρὸς τὴν ΔA , οὕτως ἡ ΓE πρὸς τὴν $E A$.

Proposition 2

If some straight-line is drawn parallel to one of the sides of a triangle then it will cut the (other) sides of the triangle proportionally. And if (two of) the sides of a triangle are cut proportionally then the straight-line joining the cutting (points) will be parallel to the remaining side of the triangle.



For let DE have been drawn parallel to one of the sides BC of triangle ABC . I say that as BD is to DA , so CE (is) to EA .

Ἐπεζεύχθωσαν γὰρ αἱ BE , $\Gamma\Delta$.

Ἴσον ἄρα ἐστὶ τὸ $B\Delta E$ τρίγωνον τῷ $\Gamma\Delta E$ τριγώνῳ· ἐπὶ γὰρ τῆς αὐτῆς βάσεως ἐστὶ τῆς ΔE καὶ ἐν ταῖς αὐταῖς παραλλήλοις ταῖς ΔE , $B\Gamma$ · ἄλλο δέ τι τὸ $A\Delta E$ τρίγωνον. τὰ δὲ ἴσα πρὸς τὸ αὐτὸ τὸν αὐτὸν ἔχει λόγον· ἐστὶν ἄρα ὡς τὸ $B\Delta E$ τρίγωνον πρὸς τὸ $A\Delta E$ [τρίγωνον], οὕτως τὸ $\Gamma\Delta E$ τρίγωνον πρὸς τὸ $A\Delta E$ τρίγωνον. ἀλλ' ὡς μὲν τὸ $B\Delta E$ τρίγωνον πρὸς τὸ $A\Delta E$, οὕτως ἡ $B\Delta$ πρὸς τὴν ΔA · ὑπὸ γὰρ τὸ αὐτὸ ὕψος ὄντα τὴν ἀπὸ τοῦ E ἐπὶ τὴν AB κάθετον ἀγομένην πρὸς ἄλληλά εἰσιν ὡς αἱ βάσεις. διὰ τὰ αὐτὰ δὴ ὡς τὸ $\Gamma\Delta E$ τρίγωνον πρὸς τὸ $A\Delta E$, οὕτως ἡ ΓE πρὸς τὴν EA · καὶ ὡς ἄρα ἡ $B\Delta$ πρὸς τὴν ΔA , οὕτως ἡ ΓE πρὸς τὴν EA .

Ἀλλὰ δὴ αἱ τοῦ $AB\Gamma$ τριγώνου πλευραὶ αἱ AB , $A\Gamma$ ἀνάλογον τετμήσθωσαν, ὡς ἡ $B\Delta$ πρὸς τὴν ΔA , οὕτως ἡ ΓE πρὸς τὴν EA , καὶ ἐπεζεύχθω ἡ ΔE · λέγω, ὅτι παράλληλός ἐστὶν ἡ ΔE τῇ $B\Gamma$.

Τῶν γὰρ αὐτῶν κατασκευασθέντων, ἐπεὶ ἐστὶν ὡς ἡ $B\Delta$ πρὸς τὴν ΔA , οὕτως ἡ ΓE πρὸς τὴν EA , ἀλλ' ὡς μὲν ἡ $B\Delta$ πρὸς τὴν ΔA , οὕτως τὸ $B\Delta E$ τρίγωνον πρὸς τὸ $A\Delta E$ τρίγωνον, ὡς δὲ ἡ ΓE πρὸς τὴν EA , οὕτως τὸ $\Gamma\Delta E$ τρίγωνον πρὸς τὸ $A\Delta E$ τρίγωνον, καὶ ὡς ἄρα τὸ $B\Delta E$ τρίγωνον πρὸς τὸ $A\Delta E$ τρίγωνον, οὕτως τὸ $\Gamma\Delta E$ τρίγωνον πρὸς τὸ $A\Delta E$ τρίγωνον. ἐκάτερον ἄρα τῶν $B\Delta E$, $\Gamma\Delta E$ τριγώνων πρὸς τὸ $A\Delta E$ τὸν αὐτὸν ἔχει λόγον. ἴσον ἄρα ἐστὶ τὸ $B\Delta E$ τρίγωνον τῷ $\Gamma\Delta E$ τριγώνῳ· καὶ εἰσιν ἐπὶ τῆς αὐτῆς βάσεως τῆς ΔE . τὰ δὲ ἴσα τρίγωνα καὶ ἐπὶ τῆς αὐτῆς βάσεως ὄντα καὶ ἐν ταῖς αὐταῖς παραλλήλοις ἐστίν. παράλληλος ἄρα ἐστὶν ἡ ΔE τῇ $B\Gamma$.

Ἐὰν ἄρα τριγώνου παρὰ μίαν τῶν πλευρῶν ἀχθῆ τις εὐθεῖα, ἀνάλογον τεμεῖ τὰς τοῦ τριγώνου πλευράς· καὶ ἐὰν αἱ τοῦ τριγώνου πλευραὶ ἀνάλογον τμηθῶσιν, ἡ ἐπὶ τὰς τομὰς ἐπιζευγυμένη εὐθεῖα παρὰ τὴν λοιπὴν ἔσται τοῦ τριγώνου πλευράν· ὅπερ ἔδει δεῖξαι.

γ'.

Ἐὰν τριγώνου ἡ γωνία δίχα τμηθῆ, ἡ δὲ τέμνουσα τὴν γωνίαν εὐθεῖα τέμνη καὶ τὴν βάσιν, τὰ τῆς βάσεως τμήματα τὸν αὐτὸν ἔξει λόγον ταῖς λοιπαῖς τοῦ τριγώνου πλευραῖς· καὶ ἐὰν τὰ τῆς βάσεως τμήματα τὸν αὐτὸν ἔχη λόγον ταῖς λοιπαῖς τοῦ τριγώνου πλευραῖς, ἡ ἀπὸ τῆς κορυφῆς ἐπὶ τὴν τομὴν ἐπιζευγυμένη εὐθεῖα δίχα τεμεῖ τὴν τοῦ τριγώνου γωνίαν.

Ἐστω τρίγωνον τὸ $AB\Gamma$, καὶ τετμήσθω ἡ ὑπὸ $BA\Gamma$ γωνία δίχα ὑπὸ τῆς $A\Delta$ εὐθείας· λέγω, ὅτι ἐστὶν ὡς ἡ $B\Delta$ πρὸς τὴν $\Gamma\Delta$, οὕτως ἡ BA πρὸς τὴν $A\Gamma$.

Ἦχθω γὰρ διὰ τοῦ Γ τῇ ΔA παράλληλος ἡ ΓE , καὶ διαχθεῖσα ἡ BA συμπίπττω αὐτῇ κατὰ τὸ E .

For let BE and CD have been joined.

Thus, triangle BDE is equal to triangle CDE . For they are on the same base DE and between the same parallels DE and BC [Prop. 1.38]. And ADE is some other triangle. And equal (magnitudes) have the same ratio to the same (magnitude) [Prop. 5.7]. Thus, as triangle BDE is to [triangle] ADE , so triangle CDE (is) to triangle ADE . But, as triangle BDE (is) to triangle ADE , so (is) BD to DA . For, having the same height—(namely), the (straight-line) drawn from E perpendicular to AB —they are to one another as their bases [Prop. 6.1]. So, for the same (reasons), as triangle CDE (is) to ADE , so CE (is) to EA . And, thus, as BD (is) to DA , so CE (is) to EA [Prop. 5.11].

And so, let the sides AB and AC of triangle ABC have been cut proportionally (such that) as BD (is) to DA , so CE (is) to EA . And let DE have been joined. I say that DE is parallel to BC .

For, by the same construction, since as BD is to DA , so CE (is) to EA , but as BD (is) to DA , so triangle BDE (is) to triangle ADE , and as CE (is) to EA , so triangle CDE (is) to triangle ADE [Prop. 6.1], thus, also, as triangle BDE (is) to triangle ADE , so triangle CDE (is) to triangle ADE [Prop. 5.11]. Thus, triangles BDE and CDE each have the same ratio to ADE . Thus, triangle BDE is equal to triangle CDE [Prop. 5.9]. And they are on the same base DE . And equal triangles, which are also on the same base, are also between the same parallels [Prop. 1.39]. Thus, DE is parallel to BC .


Thus, if some straight-line is drawn parallel to one of the sides of a triangle, then it will cut the (other) sides of the triangle proportionally. And if (two of) the sides of a triangle are cut proportionally, then the straight-line joining the cutting (points) will be parallel to the remaining side of the triangle. (Which is) the very thing it was required to show.

Proposition 3

If an angle of a triangle is cut in half, and the straight-line cutting the angle also cuts the base, then the segments of the base will have the same ratio as the remaining sides of the triangle. And if the segments of the base have the same ratio as the remaining sides of the triangle, then the straight-line joining the vertex to the cutting (point) will cut the angle of the triangle in half.

Let ABC be a triangle. And let the angle BAC have been cut in half by the straight-line AD . I say that as BD is to CD , so BA (is) to AC .

For let CE have been drawn through (point) C parallel to DA . And, BA being drawn through, let it meet




Καὶ ἐπεὶ εἰς παραλλήλους τὰς AD , EG εὐθεῖα ἐνέπεσεν ἡ AG , ἡ ἄρα ὑπὸ AGE γωνία ἴση ἐστὶ τῇ ὑπὸ $ΓAD$. ἀλλ' ἡ ὑπὸ $ΓAD$ τῇ ὑπὸ $BAΔ$ ὑπόκειται ἴση· καὶ ἡ ὑπὸ $BAΔ$ ἄρα τῇ ὑπὸ AGE ἐστὶν ἴση. πάλιν, ἐπεὶ εἰς παραλλήλους τὰς AD , EG εὐθεῖα ἐνέπεσεν ἡ BAE , ἡ ἐκτὸς γωνία ἡ ὑπὸ $BAΔ$ ἴση ἐστὶ τῇ ἐντὸς τῇ ὑπὸ AEG . ἐδείχθη δὲ καὶ ἡ ὑπὸ AGE τῇ ὑπὸ $BAΔ$ ἴση· καὶ ἡ ὑπὸ AGE ἄρα γωνία τῇ ὑπὸ AEG ἐστὶν ἴση· ὥστε καὶ πλευρὰ ἡ AE πλευρᾶ τῇ AG ἐστὶν ἴση. καὶ ἐπεὶ τριγώνου τοῦ BGE παρὰ μίαν τῶν πλευρῶν τὴν EG ἤχται ἡ AD , ἀνάλογον ἄρα ἐστὶν ὡς ἡ BD πρὸς τὴν $ΔΓ$, οὕτως ἡ BA πρὸς τὴν AE . ἴση δὲ ἡ AE τῇ AG · ὡς ἄρα ἡ BD πρὸς τὴν $ΔΓ$, οὕτως ἡ BA πρὸς τὴν AG .

Ἀλλὰ δὴ ἔστω ὡς ἡ BD πρὸς τὴν $ΔΓ$, οὕτως ἡ BA πρὸς τὴν AG , καὶ ἐπεζεύχθω ἡ AD · λέγω, ὅτι δίχα τέτμηται ἡ ὑπὸ $BAΓ$ γωνία ὑπὸ τῆς AD εὐθείας.

Τῶν γὰρ αὐτῶν κατασκευασθέντων, ἐπεὶ ἐστὶν ὡς ἡ BD πρὸς τὴν $ΔΓ$, οὕτως ἡ BA πρὸς τὴν AG , ἀλλὰ καὶ ὡς ἡ BD πρὸς τὴν $ΔΓ$, οὕτως ἐστὶν ἡ BA πρὸς τὴν AE · τριγώνου γὰρ τοῦ BGE παρὰ μίαν τὴν EG ἤχται ἡ AD · καὶ ὡς ἄρα ἡ BA πρὸς τὴν AG , οὕτως ἡ BA πρὸς τὴν AE . ἴση ἄρα ἡ AG τῇ AE · ὥστε καὶ γωνία ἡ ὑπὸ AEG τῇ ὑπὸ AGE ἐστὶν ἴση. ἀλλ' ἡ μὲν ὑπὸ AEG τῇ ἐκτὸς τῇ ὑπὸ $BAΔ$ [ἐστὶν] ἴση, ἡ δὲ ὑπὸ AGE τῇ ἐναλλάξ τῇ ὑπὸ $ΓAD$ ἐστὶν ἴση· καὶ ἡ ὑπὸ $BAΔ$ ἄρα τῇ ὑπὸ $ΓAD$ ἐστὶν ἴση. ἡ ἄρα ὑπὸ $BAΓ$ γωνία δίχα τέτμηται ὑπὸ τῆς AD εὐθείας.

Ἐὰν ἄρα τριγώνου ἡ γωνία δίχα τμηθῇ, ἡ δὲ τέμνουσα τὴν γωνίαν εὐθεῖα τέμνη καὶ τὴν βάσιν, τὰ τῆς βάσεως τμήματα τὸν αὐτὸν ἔξει λόγον ταῖς λοιπαῖς τοῦ τριγώνου πλευραῖς· καὶ ἐὰν τὰ τῆς βάσεως τμήματα τὸν αὐτὸν ἔχη λόγον ταῖς λοιπαῖς τοῦ τριγώνου πλευραῖς, ἡ ἀπὸ τῆς κορυφῆς ἐπὶ τὴν τομὴν ἐπιζευγνυμένη εὐθεῖα δίχα τέμνει τὴν τοῦ τριγώνου γωνίαν· ὅπερ ἔδει δεῖξαι.

(CE) at (point) E .[†]



And since the straight-line AC falls across the parallel (straight-lines) AD and EC , angle ACE is thus equal to CAD [Prop. 1.29]. But, (angle) CAD is assumed (to be) equal to BAD . Thus, (angle) BAD is also equal to ACE . Again, since the straight-line BAE falls across the parallel (straight-lines) AD and EC , the external angle BAD is equal to the internal (angle) AEC [Prop. 1.29]. And (angle) ACE was also shown (to be) equal to BAD . Thus, angle ACE is also equal to AEC . And, hence, side AE is equal to side AC [Prop. 1.6]. And since AD has been drawn parallel to one of the sides EC of triangle BCE , thus, proportionally, as BD is to DC , so BA (is) to AE [Prop. 6.2]. And AE (is) equal to AC . Thus, as BD (is) to DC , so BA (is) to AC .

And so, let BD be to DC , as BA (is) to AC . And let AD have been joined. I say that angle BAC has been cut in half by the straight-line AD .

For, by the same construction, since as BD is to DC , so BA (is) to AC , then also as BD (is) to DC , so BA is to AE . For AD has been drawn parallel to one (of the sides) EC of triangle BCE [Prop. 6.2]. Thus, also, as BA (is) to AC , so BA (is) to AE [Prop. 5.11]. Thus, AC (is) equal to AE [Prop. 5.9]. And, hence, angle AEC is equal to ACE [Prop. 1.5]. But, AEC [is] equal to the external (angle) BAD , and ACE is equal to the alternate (angle) CAD [Prop. 1.29]. Thus, (angle) BAD is also equal to CAD . Thus, angle BAC has been cut in half by the straight-line AD .


Thus, if an angle of a triangle is cut in half, and the straight-line cutting the angle also cuts the base, then the segments of the base will have the same ratio as the remaining sides of the triangle. And if the segments of the base have the same ratio as the remaining sides of the triangle, then the straight-line joining the vertex to the cutting (point) will cut the angle of the triangle in half. (Which is) the very thing it was required to show.

[†] The fact that the two straight-lines meet follows because the sum of ACE and CAE is less than two right-angles, as can easily be demonstrated.

See Post. 5.

δ'.

Τῶν ἰσογωνίων τριγώνων ἀνάλογόν εἰσιν αἱ πλευραὶ αἱ περὶ τὰς ἴσας γωνίας καὶ ὁμόλογοι αἱ ὑπὸ τὰς ἴσας γωνίας ὑποτείνουσαι.



Ἐστω ἰσογώνια τρίγωνα τὰ $ABΓ$, $ΔΓΕ$ ἴσην ἔχοντα τὴν μὲν ὑπὸ $ABΓ$ γωνίαν τῇ ὑπὸ $ΔΓΕ$, τὴν δὲ ὑπὸ $BAΓ$ τῇ ὑπὸ $ΓΔΕ$ καὶ ἔτι τὴν ὑπὸ $ΑΓΒ$ τῇ ὑπὸ $ΓΕΔ$. λέγω, ὅτι τῶν $ABΓ$, $ΔΓΕ$ τριγώνων ἀνάλογόν εἰσιν αἱ πλευραὶ αἱ περὶ τὰς ἴσας γωνίας καὶ ὁμόλογοι αἱ ὑπὸ τὰς ἴσας γωνίας ὑποτείνουσαι.


Κείσθω γὰρ ἐπ' εὐθείας ἡ $BΓ$ τῇ $ΓΕ$. καὶ ἐπεὶ αἱ ὑπὸ $ABΓ$, $ΑΓΒ$ γωνίαι δύο ὀρθῶν ἐλάττονές εἰσιν, ἴση δὲ ἡ ὑπὸ $ΑΓΒ$ τῇ ὑπὸ $ΔΕΓ$, αἱ ἄρα ὑπὸ $ABΓ$, $ΔΕΓ$ δύο ὀρθῶν ἐλάττονές εἰσιν· αἱ BA , ED ἄρα ἐκβαλλόμενα συμπεσοῦνται. ἐκβεβλήσθωσαν καὶ συμμιπτέτωσαν κατὰ τὸ Z .

Καὶ ἐπεὶ ἴση ἐστὶν ἡ ὑπὸ $ΔΓΕ$ γωνία τῇ ὑπὸ $ABΓ$, παράλληλός ἐστὶν ἡ BZ τῇ $ΓΔ$. πάλιν, ἐπεὶ ἴση ἐστὶν ἡ ὑπὸ $ΑΓΒ$ τῇ ὑπὸ $ΔΕΓ$, παράλληλός ἐστὶν ἡ $ΑΓ$ τῇ ZE . παραλληλόγραμμον ἄρα ἐστὶ τὸ $ZΑΓΔ$. ἴση ἄρα ἡ μὲν ZA τῇ $ΔΓ$, ἡ δὲ $ΑΓ$ τῇ $ZΔ$. καὶ ἐπεὶ τριγώνου τοῦ ZBE παρὰ μίαν τὴν ZE ἤκται ἡ $ΑΓ$, ἐστὶν ἄρα ὡς ἡ BA πρὸς τὴν AZ , οὕτως ἡ $BΓ$ πρὸς τὴν $ΓΕ$. ἴση δὲ ἡ AZ τῇ $ΓΔ$. ὡς ἄρα ἡ BA πρὸς τὴν $ΓΔ$, οὕτως ἡ $BΓ$ πρὸς τὴν $ΓΕ$, καὶ ἐναλλάξ ὡς ἡ AB πρὸς τὴν $BΓ$, οὕτως ἡ $ΔΓ$ πρὸς τὴν $ΓΕ$. πάλιν, ἐπεὶ παράλληλός ἐστὶν ἡ $ΓΔ$ τῇ BZ , ἔστιν ἄρα ὡς ἡ $BΓ$ πρὸς τὴν $ΓΕ$, οὕτως ἡ $ZΔ$ πρὸς τὴν $ΔΕ$. ἴση δὲ ἡ $ZΔ$ τῇ $ΑΓ$. ὡς ἄρα ἡ $BΓ$ πρὸς τὴν $ΓΕ$, οὕτως ἡ $ΑΓ$ πρὸς τὴν $ΔΕ$, καὶ ἐναλλάξ ὡς ἡ $BΓ$ πρὸς τὴν $ΓΑ$, οὕτως ἡ $ΓΕ$ πρὸς τὴν $ΕΔ$. ἐπεὶ οὖν ἐδείχθη ὡς μὲν ἡ AB πρὸς τὴν $BΓ$, οὕτως ἡ $ΔΓ$ πρὸς τὴν $ΓΕ$, ὡς δὲ ἡ $BΓ$ πρὸς τὴν $ΓΑ$, οὕτως ἡ $ΓΕ$ πρὸς τὴν $ΕΔ$, δι' ἴσου ἄρα ὡς ἡ BA πρὸς τὴν $ΑΓ$, οὕτως ἡ $ΓΔ$ πρὸς τὴν $ΔΕ$.

Τῶν ἄρα ἰσογωνίων τριγώνων ἀνάλογόν εἰσιν αἱ πλευραὶ αἱ περὶ τὰς ἴσας γωνίας καὶ ὁμόλογοι αἱ ὑπὸ τὰς ἴσας γωνίας ὑποτείνουσαι· ὅπερ ἔδει δεῖξαι.

Proposition 4

In equiangular triangles the sides about the equal angles are proportional, and those (sides) subtending equal angles correspond.



Let ABC and DCE be equiangular triangles, having angle ABC equal to DCE , and (angle) BAC to CDE , and, further, (angle) ACB to CED . I say that in triangles ABC and DCE the sides about the equal angles are proportional, and those (sides) subtending equal angles correspond.


Let BC be placed straight-on to CE . And since angles ABC and ACB are less than two right-angles [Prop 1.17], and ACB (is) equal to DEC , thus ABC and DEC are less than two right-angles. Thus, BA and ED , being produced, will meet [C.N. 5]. Let them have been produced, and let them meet at (point) F .

And since angle DCE is equal to ABC , BF is parallel to CD [Prop. 1.28]. Again, since (angle) ACB is equal to DEC , AC is parallel to FE [Prop. 1.28]. Thus, $FACD$ is a parallelogram. Thus, FA is equal to DC , and AC to FD [Prop. 1.34]. And since AC has been drawn parallel to one (of the sides) FE of triangle FBE , thus as BA is to AF , so BC (is) to CE [Prop. 6.2]. And AF (is) equal to CD . Thus, as BA (is) to CD , so BC (is) to CE , and, alternately, as AB (is) to BC , so DC (is) to CE [Prop. 5.16]. Again, since CD is parallel to BF , thus as BC (is) to CE , so FD (is) to DE [Prop. 6.2]. And FD (is) equal to AC . Thus, as BC is to CE , so AC (is) to DE , and, alternately, as BC (is) to CA , so CE (is) to ED [Prop. 6.2]. Therefore, since it was shown that as AB (is) to BC , so DC (is) to CE , and as BC (is) to CA , so CE (is) to ED , thus, via equality, as BA (is) to AC , so CD (is) to DE [Prop. 5.22].

Thus, in equiangular triangles the sides about the equal angles are proportional, and those (sides) subtend-

ε'.

Ἐάν δύο τρίγωνα τὰς πλευρὰς ἀνάλογον ἔχη, ἰσογώνια ἔσται τὰ τρίγωνα καὶ ἴσας ἔξει τὰς γωνίας, ὅφ' ἂς αἱ ὁμόλογοι πλευραὶ ὑποτείνουσιν.



Ἐστω δύο τρίγωνα τὰ $AB\Gamma$, ΔEZ τὰς πλευρὰς ἀνάλογον ἔχοντα, ὡς μὲν τὴν AB πρὸς τὴν $B\Gamma$, οὕτως τὴν ΔE πρὸς τὴν EZ , ὡς δὲ τὴν $B\Gamma$ πρὸς τὴν ΓA , οὕτως τὴν EZ πρὸς τὴν $Z\Delta$, καὶ ἔτι ὡς τὴν BA πρὸς τὴν $A\Gamma$, οὕτως τὴν $E\Delta$ πρὸς τὴν ΔZ . λέγω, ὅτι ἰσογώνιον ἔστι τὸ $AB\Gamma$ τρίγωνον τῷ ΔEZ τριγώνῳ καὶ ἴσας ἔξουσι τὰς γωνίας, ὅφ' ἂς αἱ ὁμόλογοι πλευραὶ ὑποτείνουσιν, τὴν μὲν ὑπὸ $AB\Gamma$ τῆ ὑπὸ ΔEZ , τὴν δὲ ὑπὸ $B\Gamma A$ τῆ ὑπὸ $EZ\Delta$ καὶ ἔτι τὴν ὑπὸ $B A \Gamma$ τῆ ὑπὸ $E \Delta Z$.


Συνεστάτω γὰρ πρὸς τῆ EZ εὐθείᾳ καὶ τοῖς πρὸς αὐτῆ σημείοις τοῖς E, Z τῆ μὲν ὑπὸ $AB\Gamma$ γωνία ἴση ἢ ὑπὸ $Z E H$, τῆ δὲ ὑπὸ $A \Gamma B$ ἴση ἢ ὑπὸ $E Z H$. λοιπὴ ἄρα ἢ πρὸς τῷ A λοιπῇ τῆ πρὸς τῷ H ἔστιν ἴση.

Ἰσογώνιον ἄρα ἔστι τὸ $AB\Gamma$ τρίγωνον τῷ $E H Z$ [τριγώνῳ]. τῶν ἄρα $AB\Gamma, E H Z$ τριγώνων ἀνάλογόν εἰσιν αἱ πλευραὶ αἱ περὶ τὰς ἴσας γωνίας καὶ ὁμόλογοι αἱ ὑπὸ τὰς ἴσας γωνίας ὑποτείνουσαι· ἔστιν ἄρα ὡς ἡ AB πρὸς τὴν $B\Gamma$, [οὕτως] ἢ HE πρὸς τὴν EZ . ἀλλ' ὡς ἡ AB πρὸς τὴν $B\Gamma$, οὕτως ὑπόκειται ἢ ΔE πρὸς τὴν EZ . ὡς ἄρα ἢ ΔE πρὸς τὴν EZ , οὕτως ἢ HE πρὸς τὴν EZ . ἑκατέρα ἄρα τῶν $\Delta E, HE$ πρὸς τὴν EZ τὸν αὐτὸν ἔχει λόγον· ἴση ἄρα ἔστιν ἢ ΔE τῆ HE . διὰ τὰ αὐτὰ δὴ καὶ ἢ ΔZ τῆ HZ ἔστιν ἴση. ἐπεὶ οὖν ἴση ἔστιν ἢ ΔE τῆ EH , κοινὴ δὲ ἢ EZ , δύο δὴ αἱ $\Delta E, EZ$ δυοὶ ταῖς HE, EZ ἴσαι εἰσίν· καὶ βάσις ἢ ΔZ βάσει τῆ ZH [ἔστιν] ἴση· γωνία ἄρα ἢ ὑπὸ ΔEZ γωνία τῆ ὑπὸ HEZ ἔστιν ἴση, καὶ τὸ ΔEZ τρίγωνον τῷ HEZ τριγώνῳ ἴσον, καὶ αἱ λοιπαὶ γωνίαι ταῖς λοιπαῖς γωνίαις ἴσαι, ὅφ' ἂς αἱ ἴσαι πλευραὶ ὑποτείνουσιν. ἴση ἄρα ἔστι καὶ ἢ μὲν ὑπὸ ΔZE γωνία τῆ ὑπὸ HZE , ἢ δὲ ὑπὸ $E \Delta Z$ τῆ ὑπὸ $E H Z$. καὶ ἐπεὶ ἢ μὲν ὑπὸ $Z E \Delta$ τῆ ὑπὸ $H E Z$ ἔστιν ἴση, ἀλλ' ἢ ὑπὸ $H E Z$ τῆ ὑπὸ $A B \Gamma$, καὶ ἢ ὑπὸ

ing equal angles correspond. (Which is) the very thing it was required to show.

Proposition 5

If two triangles have proportional sides then the triangles will be equiangular, and will have the angles which corresponding sides subtend equal.



Let ABC and DEF be two triangles having proportional sides, (so that) as AB (is) to BC , so DE (is) to EF , and as BC (is) to CA , so EF (is) to FD , and, further, as BA (is) to AC , so ED (is) to DF . I say that triangle ABC is equiangular to triangle DEF , and (that the triangles) will have the angles which corresponding sides subtend equal. (That is), (angle) ABC (equal) to DEF , BCA to EFD , and, further, BAC to EDF .

For let (angle) FEG , equal to angle ABC , and (angle) EFG , equal to ACB , have been constructed on the straight-line EF at the points E and F on it (respectively) [Prop. 1.23]. Thus, the remaining (angle) at A is equal to the remaining (angle) at G [Prop. 1.32].


Thus, triangle ABC is equiangular to [triangle] EGF . Thus, for triangles ABC and EGF , the sides about the equal angles are proportional, and (those) sides subtending equal angles correspond [Prop. 6.4]. Thus, as AB is to BC , [so] GE (is) to EF . But, as AB (is) to BC , so, it was assumed, (is) DE to EF . Thus, as DE (is) to EF , so GE (is) to EF [Prop. 5.11]. Thus, DE and GE each have the same ratio to EF . Thus, DE is equal to GE [Prop. 5.9]. So, for the same (reasons), DF is also equal to GF . Therefore, since DE is equal to EG , and EF (is) common, the two (sides) DE, EF are equal to the two (sides) GE, EF (respectively). And base DF [is] equal to base FG . Thus, angle DEF is equal to angle GEF [Prop. 1.8], and triangle DEF (is) equal to triangle GEF , and the remaining angles (are) equal to the remaining angles which the equal sides subtend [Prop. 1.4]. Thus, angle DFE is also equal to GFE , and

ΑΒΓ ἄρα γωνία τῆ ὑπὸ ΔΕΖ ἐστὶν ἴση. διὰ τὰ αὐτὰ δὴ καὶ ἡ ὑπὸ ΑΓΒ τῆ ὑπὸ ΔΖΕ ἐστὶν ἴση, καὶ ἔτι ἡ πρὸς τῷ Α τῆ πρὸς τῷ Δ· ἰσογώνιον ἄρα ἐστὶ τὸ ΑΒΓ τρίγωνον τῷ ΔΕΖ τριγώνῳ.

Ἐὰν ἄρα δύο τρίγωνα τὰς πλευρὰς ἀνάλογον ἔχῃ, ἰσογώνια ἔσται τὰ τρίγωνα καὶ ἴσας ἔξει τὰς γωνίας, ὅφ' ἂς αἱ ὁμόλογοι πλευραὶ ὑποτείνουσιν· ὅπερ ἔδει δεῖξαι.

Գ'.

Ἐὰν δύο τρίγωνα μίαν γωνίαν μιᾶ γωνία ἴσην ἔχῃ, περὶ δὲ τὰς ἴσας γωνίας τὰς πλευρὰς ἀνάλογον, ἰσογώνια ἔσται τὰ τρίγωνα καὶ ἴσας ἔξει τὰς γωνίας, ὅφ' ἂς αἱ ὁμόλογοι πλευραὶ ὑποτείνουσιν.



Ἐστω δύο τρίγωνα τὰ ΑΒΓ, ΔΕΖ μίαν γωνίαν τὴν ὑπὸ ΒΑΓ μιᾶ γωνία τῆ ὑπὸ ΕΔΖ ἴσην ἔχοντα, περὶ δὲ τὰς ἴσας γωνίας τὰς πλευρὰς ἀνάλογον, ὡς τὴν ΒΑ πρὸς τὴν ΑΓ, οὕτως τὴν ΕΔ πρὸς τὴν ΔΖ· λέγω, ὅτι ἰσογώνιον ἐστὶ τὸ ΑΒΓ τρίγωνον τῷ ΔΕΖ τριγώνῳ καὶ ἴσην ἔξει τὴν ὑπὸ ΑΒΓ γωνίαν τῆ ὑπὸ ΔΕΖ, τὴν δὲ ὑπὸ ΑΓΒ τῆ ὑπὸ ΔΖΕ.

Συνεστάτω γὰρ πρὸς τῆ ΔΖ εὐθείᾳ καὶ τοῖς πρὸς αὐτῆ σημείοις τοῖς Δ, Ζ ὀποτέρῃ μὲν τῶν ὑπὸ ΒΑΓ, ΕΔΖ ἴση ἡ ὑπὸ ΖΔΗ, τῆ δὲ ὑπὸ ΑΓΒ ἴση ἡ ὑπὸ ΔΖΗ· λοιπὴ ἄρα ἡ πρὸς τῷ Β γωνία λοιπῆ τῆ πρὸς τῷ Η ἴση ἐστίν.


Ἰσογώνιον ἄρα ἐστὶ τὸ ΑΒΓ τρίγωνον τῷ ΔΗΖ τριγώνῳ. ἀνάλογον ἄρα ἐστὶν ὡς ἡ ΒΑ πρὸς τὴν ΑΓ, οὕτως ἡ ΗΔ πρὸς τὴν ΔΖ. ὑπόκειται δὲ καὶ ὡς ἡ ΒΑ πρὸς τὴν ΑΓ, οὕτως ἡ ΕΔ πρὸς τὴν ΔΖ· καὶ ὡς ἄρα ἡ ΕΔ πρὸς τὴν ΔΖ, οὕτως ἡ ΗΔ πρὸς τὴν ΔΖ. ἴση ἄρα ἡ ΕΔ τῆ ΔΗ· καὶ κοινὴ ἡ ΔΖ· δύο δὴ αἱ ΕΔ, ΔΖ δυσὶ ταῖς ΗΔ, ΔΖ ἴσας εἰσίν· καὶ γωνία ἡ ὑπὸ ΕΔΖ γωνία τῆ ὑπὸ ΗΔΖ [ἐστὶν] ἴση· βάσις ἄρα ἡ ΕΖ βάσει τῆ ΗΖ ἐστὶν ἴση, καὶ τὸ ΔΕΖ τρίγωνον τῷ ΗΔΖ τριγώνῳ ἴσον ἐστίν, καὶ αἱ λοιπαὶ γωνίαὶ ταῖς λοιπαῖς γωνίαις ἴσας ἔσονται, ὅφ' ἂς ἴσας πλευραὶ ὑποτείνουσιν. ἴση ἄρα ἐστὶν ἡ μὲν ὑπὸ ΔΖΗ τῆ ὑπο ΔΖΕ, ἡ δὲ ὑπὸ ΔΗΖ

(angle) EDF to EGF . And since (angle) FED is equal to GEF , and (angle) GEF to ABC , angle ABC is thus also equal to DEF . So, for the same (reasons), (angle) ACB is also equal to DFE , and, further, the (angle) at A to the (angle) at D . Thus, triangle ABC is equiangular to triangle DEF .

Thus, if two triangles have proportional sides then the triangles will be equiangular, and will have the angles which corresponding sides subtend equal. (Which is) the very thing it was required to show.

Proposition 6

If two triangles have one angle equal to one angle, and the sides about the equal angles proportional, then the triangles will be equiangular, and will have the angles which corresponding sides subtend equal.



Let ABC and DEF be two triangles having one angle, BAC , equal to one angle, EDF (respectively), and the sides about the equal angles proportional, (so that) as BA (is) to AC , so ED (is) to DF . I say that triangle ABC is equiangular to triangle DEF , and will have angle ABC equal to DEF , and (angle) ACB to DFE .

For let (angle) FDG , equal to each of BAC and EDF , and (angle) DFG , equal to ACB , have been constructed on the straight-line AF at the points D and F on it (respectively) [Prop. 1.23]. Thus, the remaining angle at B is equal to the remaining angle at G [Prop. 1.32].


Thus, triangle ABC is equiangular to triangle DGF . Thus, proportionally, as BA (is) to AC , so GD (is) to DF [Prop. 6.4]. And it was also assumed that as BA (is) to AC , so ED (is) to DF . And, thus, as ED (is) to DF , so GD (is) to DF [Prop. 5.11]. Thus, ED (is) equal to DG [Prop. 5.9]. And DF (is) common. So, the two (sides) ED , DF are equal to the two (sides) GD , DF (respectively). And angle EDF [is] equal to angle GDF . Thus, base EF is equal to base GF , and triangle DEF is equal to triangle GDF , and the remaining angles

τῆ ὑπὸ ΔΕΖ. ἀλλ' ἡ ὑπὸ ΔΖΗ τῆ ὑπὸ ΑΓΒ ἐστὶν ἴση· καὶ ἡ ὑπὸ ΑΓΒ ἄρα τῆ ὑπὸ ΔΖΕ ἐστὶν ἴση. ὑπόκειται δὲ καὶ ἡ ὑπὸ ΒΑΓ τῆ ὑπὸ ΕΔΖ ἴση· καὶ λοιπὴ ἄρα ἡ πρὸς τῷ Β λοιπὴ τῆ πρὸς τῷ Ε ἴση ἐστίν· ἰσογώνιον ἄρα ἐστὶ τὸ ΑΒΓ τρίγωνον τῷ ΔΕΖ τριγώνῳ.

Ἐὰν ἄρα δύο τρίγωνα μίαν γωνίαν μιᾶ γωνία ἴσην ἔχῃ, περὶ δὲ τὰς ἴσας γωνίας τὰς πλευρὰς ἀνάλογον, ἰσογώνια ἔσται τὰ τρίγωνα καὶ ἴσας ἔξει τὰς γωνίας, ὅφ' ἂς αἱ ὁμόλογοι πλευραὶ ὑποτείνουσιν· ὅπερ ἔδει δεῖξαι.

ζ'.

Ἐὰν δύο τρίγωνα μίαν γωνίαν μιᾶ γωνία ἴσην ἔχῃ, περὶ δὲ ἄλλας γωνίας τὰς πλευρὰς ἀνάλογον, τῶν δὲ λοιπῶν ἑκατέραν ἅμα ἤτοι ἐλάσσονα ἢ μὴ ἐλάσσονα ὀρθῆς, ἰσογώνια ἔσται τὰ τρίγωνα καὶ ἴσας ἔξει τὰς γωνίας, περὶ ἂς ἀνάλογόν εἰσιν αἱ πλευραί.



Ἐστω δύο τρίγωνα τὰ ΑΒΓ, ΔΕΖ μίαν γωνίαν μιᾶ γωνία ἴσην ἔχοντα τὴν ὑπὸ ΒΑΓ τῆ ὑπὸ ΕΔΖ, περὶ δὲ ἄλλας γωνίας τὰς ὑπὸ ΑΒΓ, ΔΕΖ τὰς πλευρὰς ἀνάλογον, ὡς τὴν ΑΒ πρὸς τὴν ΒΓ, οὕτως τὴν ΔΕ πρὸς τὴν ΕΖ, τῶν δὲ λοιπῶν τῶν πρὸς τοῖς Γ, Ζ πρότερον ἑκατέραν ἅμα ἐλάσσονα ὀρθῆς· λέγω, ὅτι ἰσογώνιον ἐστὶ τὸ ΑΒΓ τρίγωνον τῷ ΔΕΖ τριγώνῳ, καὶ ἴση ἔσται ἡ ὑπὸ ΑΒΓ γωνία τῆ ὑπὸ ΔΕΖ, καὶ λοιπὴ δηλονότι ἢ πρὸς τῷ Γ λοιπὴ τῆ πρὸς τῷ Ζ ἴση.

Εἰ γὰρ ἄνισός ἐστὶν ἡ ὑπὸ ΑΒΓ γωνία τῆ ὑπὸ ΔΕΖ, μία αὐτῶν μείζων ἐστίν. ἔστω μείζων ἡ ὑπὸ ΑΒΓ. καὶ συνεστάτω πρὸς τῆ ΑΒ εὐθείᾳ καὶ τῷ πρὸς αὐτῇ σημείῳ τῷ Β τῆ ὑπὸ ΔΕΖ γωνία ἴση ἡ ὑπὸ ΑΒΗ.


Καὶ ἐπεὶ ἴση ἐστὶν ἡ μὲν Α γωνία τῆ Δ, ἡ δὲ ὑπὸ ΑΒΗ τῆ ὑπὸ ΔΕΖ, λοιπὴ ἄρα ἡ ὑπὸ ΑΗΒ λοιπὴ τῆ ὑπὸ ΔΖΕ ἐστὶν ἴση. ἰσογώνιον ἄρα ἐστὶ τὸ ΑΒΗ τρίγωνον τῷ ΔΕΖ

will be equal to the remaining angles which the equal sides subtend [Prop. 1.4]. Thus, (angle) DFG is equal to DFE , and (angle) DGF to DEF . But, (angle) DFG is equal to ACB . Thus, (angle) ACB is also equal to DFE . And (angle) BAC was also assumed (to be) equal to EDF . Thus, the remaining (angle) at B is equal to the remaining (angle) at E [Prop. 1.32]. Thus, triangle ABC is equiangular to triangle DEF .

Thus, if two triangles have one angle equal to one angle, and the sides about the equal angles proportional, then the triangles will be equiangular, and will have the angles which corresponding sides subtend equal. (Which is) the very thing it was required to show.

Proposition 7

If two triangles have one angle equal to one angle, and the sides about other angles proportional, and the remaining angles either both less than, or both not less than, right-angles, then the triangles will be equiangular, and will have the angles about which the sides are proportional equal.



Let ABC and DEF be two triangles having one angle, BAC , equal to one angle, EDF (respectively), and the sides about (some) other angles, ABC and DEF (respectively), proportional, (so that) as AB (is) to BC , so DE (is) to EF , and the remaining (angles) at C and F , first of all, both less than right-angles. I say that triangle ABC is equiangular to triangle DEF , and (that) angle ABC will be equal to DEF , and (that) the remaining (angle) at C (will be) manifestly equal to the remaining (angle) at F .

For if angle ABC is not equal to (angle) DEF then one of them is greater. Let ABC be greater. And let (angle) ABG , equal to (angle) DEF , have been constructed on the straight-line AB at the point B on it [Prop. 1.23].

And since angle A is equal to (angle) D , and (angle) ABG to DEF , the remaining (angle) AGB is thus equal

τριγώνω. ἔστιν ἄρα ὡς ἡ AB πρὸς τὴν BH , οὕτως ἡ ΔE πρὸς τὴν EZ . ὡς δὲ ἡ ΔE πρὸς τὴν EZ , [οὕτως] ὑπόκειται ἡ AB πρὸς τὴν BG . ἡ AB ἄρα πρὸς ἑκατέραν τῶν BG , BH τὸν αὐτὸν ἔχει λόγον· ἴση ἄρα ἡ BG τῆ BH . ὥστε καὶ γωνία ἡ πρὸς τῷ Γ γωνία τῆ ὑπὸ BHG ἔστιν ἴση. ἐλάττων δὲ ὀρθῆς ὑπόκειται ἡ πρὸς τῷ Γ . ἐλάττων ἄρα ἔστιν ὀρθῆς καὶ ὑπὸ BHG . ὥστε ἡ ἐφεξῆς αὐτῆ γωνία ἡ ὑπὸ AHB μείζων ἔστιν ὀρθῆς. καὶ ἐδείχθη ἴση οὖσα τῆ πρὸς τῷ Z · καὶ ἡ πρὸς τῷ Z ἄρα μείζων ἔστιν ὀρθῆς. ὑπόκειται δὲ ἐλάσσων ὀρθῆς· ὅπερ ἔστιν ἀτοπον. οὐκ ἄρα ἄνισός ἐστιν ἡ ὑπὸ ABG γωνία τῆ ὑπὸ ΔEZ . ἴση ἄρα. ἔστι δὲ καὶ ἡ πρὸς τῷ A ἴση τῆ πρὸς τῷ Δ · καὶ λοιπὴ ἄρα ἡ πρὸς τῷ Γ λοιπῆ τῆ πρὸς τῷ Z ἴση ἔστιν. ἰσογώνιον ἄρα ἔστι τὸ ABG τρίγωνον τῷ ΔEZ τριγώνω.

Ἄλλὰ δὴ πάλιν ὑποκείσθω ἑκατέρα τῶν πρὸς τοῖς Γ , Z μὴ ἐλάσσων ὀρθῆς· λέγω πάλιν, ὅτι καὶ οὕτως ἔστιν ἰσογώνιον τὸ ABG τρίγωνον τῷ ΔEZ τριγώνω.

Τῶν γὰρ αὐτῶν κατασκευασθέντων ὁμοίως δεῖξομεν, ὅτι ἴση ἔστιν ἡ BG τῆ BH . ὥστε καὶ γωνία ἡ πρὸς τῷ Γ τῆ ὑπὸ BHG ἴση ἔστιν. οὐκ ἐλάττων δὲ ὀρθῆς ἡ πρὸς τῷ Γ . οὐκ ἐλάττων ἄρα ὀρθῆς οὐδὲ ἡ ὑπὸ BHG . τριγώνου δὲ τοῦ BHG αἱ δύο γωνίαι δύο ὀρθῶν οὐκ εἰσιν ἐλάττονες· ὅπερ ἔστιν ἀδύνατον. οὐκ ἄρα πάλιν ἄνισός ἐστιν ἡ ὑπὸ ABG γωνία τῆ ὑπὸ ΔEZ . ἴση ἄρα. ἔστι δὲ καὶ ἡ πρὸς τῷ A τῆ πρὸς τῷ Δ ἴση· λοιπὴ ἄρα ἡ πρὸς τῷ Γ λοιπῆ τῆ πρὸς τῷ Z ἴση ἔστιν. ἰσογώνιον ἄρα ἔστι τὸ ABG τρίγωνον τῷ ΔEZ τριγώνω.

Ἐὰν ἄρα δύο τρίγωνα μίαν γωνίαν μιᾶ γωνία ἴσην ἔχῃ, περὶ δὲ ἄλλας γωνίας τὰς πλευρὰς ἀνάλογον, τῶν δὲ λοιπῶν ἑκατέραν ἅμα ἐλάττονα ἢ μὴ ἐλάττονα ὀρθῆς, ἰσογώνια ἔσται τὰ τρίγωνα καὶ ἴσας ἔξει τὰς γωνίας, περὶ ἃς ἀνάλογόν εἰσιν αἱ πλευραὶ· ὅπερ ἔδει δεῖξαι.

η'.

Ἐὰν ἐν ὀρθογώνιῳ τριγώνω ἀπὸ τῆς ὀρθῆς γωνίας ἐπὶ τὴν βάσιν κάθετος ἀχθῆ, τὰ πρὸς τῆ καθέτω τρίγωνα ὁμοία ἔστι τῷ τε ὅλῳ καὶ ἀλλήλοισ.

Ἐστω τρίγωνον ὀρθογώνιον τὸ ABG ὀρθὴν ἔχον τὴν ὑπὸ BAG γωνίαν, καὶ ἤχθω ἀπὸ τοῦ A ἐπὶ τὴν BG κάθετος ἡ AD . λέγω, ὅτι ὁμοίον ἔστιν ἑκάτερον τῶν ABD , ADG

to the remaining (angle) DFE [Prop. 1.32]. Thus, triangle ABG is equiangular to triangle DEF . Thus, as AB is to BG , so DE (is) to EF [Prop. 6.4]. And as DE (is) to EF , [so] it was assumed (is) AB to BC . Thus, AB has the same ratio to each of BC and BG [Prop. 5.11]. Thus, BC (is) equal to BG [Prop. 5.9]. And, hence, the angle at C is equal to angle BGC [Prop. 1.5]. And the angle at C was assumed (to be) less than a right-angle. Thus, (angle) BGC is also less than a right-angle. Hence, the adjacent angle to it, AGB , is greater than a right-angle [Prop. 1.13]. And (AGB) was shown to be equal to the (angle) at F . Thus, the (angle) at F is also greater than a right-angle. But it was assumed (to be) less than a right-angle. The very thing is absurd. Thus, angle ABC is not unequal to (angle) DEF . Thus, (it is) equal. And the (angle) at A is also equal to the (angle) at D . And thus the remaining (angle) at C is equal to the remaining (angle) at F [Prop. 1.32]. Thus, triangle ABC is equiangular to triangle DEF .

But, again, let each of the (angles) at C and F be assumed (to be) not less than a right-angle. I say, again, that triangle ABC is equiangular to triangle DEF in this case also.

For, with the same construction, we can similarly show that BC is equal to BG . Hence, also, the angle at C is equal to (angle) BGC . And the (angle) at C (is) not less than a right-angle. Thus, BGC (is) not less than a right-angle either. So, in triangle BGC the (sum of) two angles is not less than two right-angles. The very thing is impossible [Prop. 1.17]. Thus, again, angle ABC is not unequal to DEF . Thus, (it is) equal. And the (angle) at A is also equal to the (angle) at D . Thus, the remaining (angle) at C is equal to the remaining (angle) at F [Prop. 1.32]. Thus, triangle ABC is equiangular to triangle DEF .


Thus, if two triangles have one angle equal to one angle, and the sides about other angles proportional, and the remaining angles both less than, or both not less than, right-angles, then the triangles will be equiangular, and will have the angles about which the sides (are) proportional equal. (Which is) the very thing it was required to show.

Proposition 8

If, in a right-angled triangle, a (straight-line) is drawn from the right-angle perpendicular to the base then the triangles around the perpendicular are similar to the whole (triangle), and to one another.

Let ABC be a right-angled triangle having the angle BAC a right-angle, and let AD have been drawn from

τριγώνων ὅλων τῶν $AB\Gamma$ καὶ ἔτι ἀλλήλοις.




Ἐπεὶ γὰρ ἴση ἐστὶν ἡ ὑπὸ $BA\Gamma$ τῆς ὑπὸ $A\Delta B$: ὀρθὴ γὰρ ἑκατέρα· καὶ κοινὴ τῶν δύο τριγώνων τοῦ τε $AB\Gamma$ καὶ τοῦ $AB\Delta$ ἢ πρὸς τῶν B , λοιπὴ ἄρα ἡ ὑπὸ AGB λοιπὴ τῆς ὑπὸ $BA\Delta$ ἐστὶν ἴση· ἰσογώνιον ἄρα ἐστὶ τὸ $AB\Gamma$ τρίγωνον τῶν $AB\Delta$ τριγώνων. ἔστιν ἄρα ὡς ἡ $B\Gamma$ ὑποτείνουσα τὴν ὀρθὴν τοῦ $AB\Gamma$ τριγώνου πρὸς τὴν BA ὑποτείνουσαν τὴν ὀρθὴν τοῦ $AB\Delta$ τριγώνου, οὕτως αὐτὴ ἡ AB ὑποτείνουσα τὴν πρὸς τῶν Γ γωνίαν τοῦ $AB\Gamma$ τριγώνου πρὸς τὴν $B\Delta$ ὑποτείνουσαν τὴν ἴσην τὴν ὑπὸ $BA\Delta$ τοῦ $AB\Delta$ τριγώνου, καὶ ἔτι ἡ AG πρὸς τὴν $A\Delta$ ὑποτείνουσαν τὴν πρὸς τῶν B γωνίαν κοινὴν τῶν δύο τριγώνων. τὸ $AB\Gamma$ ἄρα τρίγωνον τῶν $AB\Delta$ τριγώνων ἰσογώνιον τέ ἐστι καὶ τὰς περὶ τὰς ἴσας γωνίας πλευρὰς ἀνάλογον ἔχει. ὅμοιον ἄρα [ἐστὶ] τὸ $AB\Gamma$ τρίγωνον τῶν $AB\Delta$ τριγώνων. ὁμοίως δὲ δείξομεν, ὅτι καὶ τῶν $A\Delta\Gamma$ τριγώνων ὅμοιον ἐστὶ τὸ $AB\Gamma$ τρίγωνον· ἑκάτερον ἄρα τῶν $AB\Delta$, $A\Delta\Gamma$ [τριγώνων] ὅμοιον ἐστὶν ὅλων τῶν $AB\Gamma$.

Λέγω δὴ, ὅτι καὶ ἀλλήλοις ἐστὶν ὅμοια τὰ $AB\Delta$, $A\Delta\Gamma$ τρίγωνα.

Ἐπεὶ γὰρ ὀρθὴ ἡ ὑπὸ $B\Delta A$ ὀρθὴ τῆς ὑπὸ $A\Delta\Gamma$ ἐστὶν ἴση, ἀλλὰ μὴν καὶ ἡ ὑπὸ $BA\Delta$ τῆς πρὸς τῶν Γ ἐδείχθη ἴση, καὶ λοιπὴ ἄρα ἡ πρὸς τῶν B λοιπὴ τῆς ὑπὸ $\Delta A\Gamma$ ἐστὶν ἴση· ἰσογώνιον ἄρα ἐστὶ τὸ $AB\Delta$ τρίγωνον τῶν $A\Delta\Gamma$ τριγώνων. ἔστιν ἄρα ὡς ἡ $B\Delta$ τοῦ $AB\Delta$ τριγώνου ὑποτείνουσα τὴν ὑπὸ $BA\Delta$ πρὸς τὴν ΔA τοῦ $A\Delta\Gamma$ τριγώνου ὑποτείνουσαν τὴν πρὸς τῶν Γ ἴσην τῆς ὑπὸ $BA\Delta$, οὕτως αὐτὴ ἡ $A\Delta$ τοῦ $AB\Delta$ τριγώνου ὑποτείνουσα τὴν πρὸς τῶν B γωνίαν πρὸς τὴν $\Delta\Gamma$ ὑποτείνουσαν τὴν ὑπὸ $\Delta A\Gamma$ τοῦ $A\Delta\Gamma$ τριγώνου ἴσην τῆς πρὸς τῶν B , καὶ ἔτι ἡ BA πρὸς τὴν AG ὑποτείνουσαι τὰς ὀρθὰς· ὅμοιον ἄρα ἐστὶ τὸ $AB\Delta$ τρίγωνον τῶν $A\Delta\Gamma$ τριγώνων.

Ἐὰν ἄρα ἐν ὀρθογώνιῳ τριγώνῳ ἀπὸ τῆς ὀρθῆς γωνίας ἐπὶ τὴν βᾶσιν κάθετος ἀχθῆ, τὰ πρὸς τῆς καθέτου τρίγωνα ὁμοία ἐστὶ τῶν τε ὅλων καὶ ἀλλήλοις [ὅπερ ἔδει δεῖξαι].

A , perpendicular to BC [Prop. 1.12]. I say that triangles ABD and ADC are each similar to the whole (triangle) ABC and, further, to one another.



For since (angle) BAC is equal to ADB —for each (are) right-angles—and the (angle) at B (is) common to the two triangles ABC and ABD , the remaining (angle) ACB is thus equal to the remaining (angle) BAD [Prop. 1.32]. Thus, triangle ABC is equiangular to triangle ABD . Thus, as BC , subtending the right-angle in triangle ABC , is to BA , subtending the right-angle in triangle ABD , so the same AB , subtending the angle at C in triangle ABC , (is) to BD , subtending the equal (angle) BAD in triangle ABD , and, further, (so is) AC to AD , (both) subtending the angle at B common to the two triangles [Prop. 6.4]. Thus, triangle ABC is equiangular to triangle ABD , and has the sides about the equal angles proportional. Thus, triangle ABC [is] similar to triangle ABD [Def. 6.1]. So, similarly, we can show that triangle ABC is also similar to triangle ADC . Thus, [triangles] ABD and ADC are each similar to the whole (triangle) ABC .

So I say that triangles ABD and ADC are also similar to one another.

For since the right-angle BDA is equal to the right-angle ADC , and, indeed, (angle) BAD was also shown (to be) equal to the (angle) at C , thus the remaining (angle) at B is also equal to the remaining (angle) DAC [Prop. 1.32]. Thus, triangle ABD is equiangular to triangle ADC . Thus, as BD , subtending (angle) BAD in triangle ABD , is to DA , subtending the (angle) at C in triangle ADC , (which is) equal to (angle) BAD , so (is) the same AD , subtending the angle at B in triangle ABD , to DC , subtending (angle) DAC in triangle ADC , (which is) equal to the (angle) at B , and, further, (so is) BA to AC , (each) subtending right-angles [Prop. 6.4]. Thus, triangle ABD is similar to triangle ADC [Def. 6.1].

Thus, if, in a right-angled triangle, a (straight-line) is drawn from the right-angle perpendicular to the base


Πόρισμα.

Ἐκ δὴ τούτου φανερόν, ὅτι ἐὰν ἐν ὀρθογωνίῳ τριγώνῳ ἀπὸ τῆς ὀρθῆς γωνίας ἐπὶ τὴν βάσιν κάθετος ἀχθῆ, ἡ ἀχθεῖσα τῶν τῆς βάσεως τμημάτων μέση ἀνάλογόν ἐστίν· ὅπερ ἔδει δεῖξαι.

† In other words, the perpendicular is the geometric mean of the pieces.

θ'.

Τῆς δοθείσης εὐθείας τὸ προσταχθὲν μέρος ἀφελεῖν.



Ἐστω ἡ δοθεῖσα εὐθεῖα ἡ AB . δεῖ δὴ τῆς AB τὸ προσταχθὲν μέρος ἀφελεῖν.

Ἐπιτετάχτω δὴ τὸ τρίτον. [καὶ] διήθχω τις ἀπὸ τοῦ A εὐθεῖα ἡ AG γωνίαν περιέχουσα μετὰ τῆς AB τυχούσαν· καὶ εἰλήφθω τυχὸν σημεῖον ἐπὶ τῆς AG τὸ Δ , καὶ κείσθωσαν τῇ $A\Delta$ ἴσαι αἱ ΔE , $E\Gamma$. καὶ ἐπεζεύχθω ἡ $B\Gamma$, καὶ διὰ τοῦ Δ παράλληλος αὐτῇ ἦχθω ἡ ΔZ .

Ἐπεὶ οὖν τριγώνου τοῦ $AB\Gamma$ παρὰ μίαν τῶν πλευρῶν τὴν $B\Gamma$ ἦκται ἡ $Z\Delta$, ἀνάλογον ἄρα ἐστὶν ὡς ἡ $\Gamma\Delta$ πρὸς τὴν ΔA , οὕτως ἡ BZ πρὸς τὴν ZA . διπλῆ δὲ ἡ $\Gamma\Delta$ τῆς ΔA · διπλῆ ἄρα καὶ ἡ BZ τῆς ZA · τριπλῆ ἄρα ἡ BA τῆς AZ .

Τῆς ἄρα δοθείσης εὐθείας τῆς AB τὸ ἐπιταχθὲν τρίτον μέρος ἀφῆρηται τὸ AZ · ὅπερ ἔδει ποιῆσαι.

ι'.

Τὴν δοθεῖσαν εὐθεῖαν ἄτμητον τῇ δοθείσῃ τετμημένη ὁμοίως τεμεῖν.


then the triangles around the perpendicular are similar to the whole (triangle), and to one another. [(Which is) the very thing it was required to show.]

Corollary

So (it is) clear, from this, that if, in a right-angled triangle, a (straight-line) is drawn from the right-angle perpendicular to the base then the (straight-line so) drawn is in mean proportion to the pieces of the base.† (Which is) the very thing it was required to show.

Proposition 9

To cut off a prescribed part from a given straight-line.



Let AB be the given straight-line. So it is required to cut off a prescribed part from AB .


So let a third (part) have been prescribed. [And] let some straight-line AC have been drawn from (point) A , encompassing a random angle with AB . And let a random point D have been taken on AC . And let DE and EC be made equal to AD [Prop. 1.3]. And let BC have been joined. And let DF have been drawn through D parallel to it [Prop. 1.31].

Therefore, since FD has been drawn parallel to one of the sides, BC , of triangle ABC , then, proportionally, as CD is to DA , so BF (is) to FA [Prop. 6.2]. And CD (is) double DA . Thus, BF (is) also double FA . Thus, BA (is) triple AF .

Thus, the prescribed third part, AF , has been cut off from the given straight-line, AB . (Which is) the very thing it was required to do.

Proposition 10

To cut a given uncut straight-line similarly to a given cut (straight-line).



Ἐστω ἡ μὲν δοθεῖσα εὐθεῖα ἀτμητος ἡ AB , ἡ δὲ τετμημένη ἡ $A\Gamma$ κατὰ τὰ Δ , E σημεία, καὶ κείσθωσαν ὥστε γωνίαν τυχοῦσαν περιέχειν, καὶ ἐπεζεύχθω ἡ GB , καὶ διὰ τῶν Δ , E τῆ $B\Gamma$ παράλληλοι ἦχθωσαν αἱ ΔZ , EH , διὰ δὲ τοῦ Δ τῆ AB παράλληλος ἦχθω ἡ $\Delta\Theta K$.

Παράλληλόγραμμον ἄρα ἐστὶν ἐκάτερον τῶν $Z\Theta$, ΘB . ἴση ἄρα ἡ μὲν $\Delta\Theta$ τῆ ZH , ἡ δὲ ΘK τῆ HB . καὶ ἐπεὶ τριγώνου τοῦ $\Delta K\Gamma$ παρὰ μίαν τῶν πλευρῶν τὴν $K\Gamma$ εὐθεῖα ἦχται ἡ ΘE , ἀνάλογον ἄρα ἐστὶν ὡς ἡ GE πρὸς τὴν $E\Delta$, οὕτως ἡ $K\Theta$ πρὸς τὴν $\Theta\Delta$. ἴση δὲ ἡ μὲν $K\Theta$ τῆ BH , ἡ δὲ $\Theta\Delta$ τῆ HZ . ἔστιν ἄρα ὡς ἡ GE πρὸς τὴν $E\Delta$, οὕτως ἡ BH πρὸς τὴν HZ . πάλιν, ἐπεὶ τριγώνου τοῦ AHE παρὰ μίαν τῶν πλευρῶν τὴν HE ἦχται ἡ $Z\Delta$, ἀνάλογον ἄρα ἐστὶν ὡς ἡ $E\Delta$ πρὸς τὴν ΔA , οὕτως ἡ HZ πρὸς τὴν ZA . ἐδείχθη δὲ καὶ ὡς ἡ GE πρὸς τὴν $E\Delta$, οὕτως ἡ BH πρὸς τὴν HZ . ἔστιν ἄρα ὡς μὲν ἡ GE πρὸς τὴν $E\Delta$, οὕτως ἡ BH πρὸς τὴν HZ , ὡς δὲ ἡ $E\Delta$ πρὸς τὴν ΔA , οὕτως ἡ HZ πρὸς τὴν ZA .


Ἡ ἄρα δοθεῖσα εὐθεῖα ἀτμητος ἡ AB τῆ δοθείσῃ εὐθείᾳ τετμημένη τῆ $A\Gamma$ ὁμοίως τέτμηται· ὅπερ ἔδει ποιῆσαι.

ια'.

Δύο δοθεισῶν εὐθειῶν τρίτην ἀνάλογον προσευρεῖν.

Ἐστώσαν αἱ δοθεῖσαι [δύο εὐθεῖαι] αἱ BA , AC καὶ κείσθωσαν γωνίαν περιέχουσαι τυχοῦσαν. δεῖ δὴ τῶν BA , AC τρίτην ἀνάλογον προσευρεῖν. ἐκβεβλήσθωσαν γὰρ ἐπὶ τὰ Δ , E σημεία, καὶ κείσθω τῆ $A\Gamma$ ἴση ἡ $B\Delta$, καὶ ἐπεζεύχθω ἡ $B\Gamma$, καὶ διὰ τοῦ Δ παράλληλος αὐτῆ ἦχθω ἡ ΔE .

Ἐπεὶ οὖν τριγώνου τοῦ $A\Delta E$ παρὰ μίαν τῶν πλευρῶν τὴν ΔE ἦχται ἡ $B\Gamma$, ἀνάλογόν ἐστὶν ὡς ἡ AB πρὸς τὴν $B\Delta$, οὕτως ἡ $A\Gamma$ πρὸς τὴν GE . ἴση δὲ ἡ $B\Delta$ τῆ $A\Gamma$. ἔστιν ἄρα ὡς ἡ AB πρὸς τὴν $A\Gamma$, οὕτως ἡ $A\Gamma$ πρὸς τὴν GE .



Let AB be the given uncut straight-line, and AC a (straight-line) cut at points D and E , and let (AC) be laid down so as to encompass a random angle (with AB). And let CB have been joined. And let DF and EG have been drawn through (points) D and E (respectively), parallel to BC , and let DHK have been drawn through (point) D , parallel to AB [Prop. 1.31].

Thus, FH and HB are each parallelograms. Thus, DH (is) equal to FG , and HK to GB [Prop. 1.34]. And since the straight-line HE has been drawn parallel to one of the sides, KC , of triangle DKC , thus, proportionally, as CE is to ED , so KH (is) to HD [Prop. 6.2]. And KH (is) equal to BG , and HD to GF . Thus, as CE is to ED , so BG (is) to GF . Again, since FD has been drawn parallel to one of the sides, GE , of triangle AGE , thus, proportionally, as ED is to DA , so GF (is) to FA [Prop. 6.2]. And it was also shown that as CE (is) to ED , so BG (is) to GF . Thus, as CE is to ED , so BG (is) to GF , and as ED (is) to DA , so GF (is) to FA .


Thus, the given uncut straight-line, AB , has been cut similarly to the given cut straight-line, AC . (Which is) the very thing it was required to do.

Proposition 11

To find a third (straight-line) proportional to two given straight-lines.


Let BA and AC be the [two] given [straight-lines], and let them be laid down encompassing a random angle. So it is required to find a third (straight-line) proportional to BA and AC . For let (BA and AC) have been produced to points D and E (respectively), and let BD be made equal to AC [Prop. 1.3]. And let BC have been joined. And let DE have been drawn through (point) D parallel to it [Prop. 1.31].

Therefore, since BC has been drawn parallel to one of the sides DE of triangle ADE , proportionally, as AB is to BD , so AC (is) to CE [Prop. 6.2]. And BD (is) equal



Δύο ἄρα δοθεισῶν εὐθειῶν τῶν AB, AG τρίτη ἀνάλογον αὐταῖς προσεύρηται ἡ ΓΕ· ὅπερ ἔδει ποιῆσαι.


to AC. Thus, as AB is to AC, so AC (is) to CE.



Thus, a third (straight-line), CE, has been found (which is) proportional to the two given straight-lines, AB and AC. (Which is) the very thing it was required to do.

ιβ'.

Τριῶν δοθεισῶν εὐθειῶν τετάρτην ἀνάλογον προσευρεῖν.



Ἐστωσαν αἱ δοθεῖσαι τρεῖς εὐθεῖαι αἱ A, B, Γ· δεῖ δὴ τῶν A, B, Γ τετάρτην ἀνάλογον προσευρεῖν.


Ἐκκείσθωσαν δύο εὐθεῖαι αἱ ΔE, ΔZ γωνίαν περιέχουσαι [τυχοῦσαν] τὴν ὑπὸ EΔZ· καὶ κείσθω τῇ μὲν A ἴση ἡ ΔH, τῇ δὲ B ἴση ἡ HE, καὶ ἔτι τῇ Γ ἴση ἡ ΔΘ· καὶ ἐπιζευχθείσης τῆς HΘ παράλληλος αὐτῇ ἦχθω διὰ τοῦ E ἡ EZ.

Ἐπεὶ οὖν τριγώνου τοῦ ΔEZ παρὰ μίαν τὴν EZ ἦχται ἡ HΘ, ἔστιν ἄρα ὡς ἡ ΔH πρὸς τὴν HE, οὕτως ἡ ΔΘ πρὸς τὴν ΘZ. ἴση δὲ ἡ μὲν ΔH τῇ A, ἡ δὲ HE τῇ B, ἡ δὲ ΔΘ τῇ Γ· ἔστιν ἄρα ὡς ἡ A πρὸς τὴν B, οὕτως ἡ Γ πρὸς τὴν ΘZ.

Τριῶν ἄρα δοθεισῶν εὐθειῶν τῶν A, B, Γ τετάρτη ἀνάλογον προσεύρηται ἡ ΘZ· ὅπερ ἔδει ποιῆσαι.

Proposition 12

To find a fourth (straight-line) proportional to three given straight-lines.




Let A, B, and C be the three given straight-lines. So it is required to find a fourth (straight-line) proportional to A, B, and C.

Let the two straight-lines DE and DF be set out encompassing the [random] angle EDF. And let DG be made equal to A, and GE to B, and, further, DH to C [Prop. 1.3]. And GH being joined, let EF have been drawn through (point) E parallel to it [Prop. 1.31].

Therefore, since GH has been drawn parallel to one of the sides EF of triangle DEF, thus as DG is to GE, so DH (is) to HF [Prop. 6.2]. And DG (is) equal to A, and GE to B, and DH to C. Thus, as A is to B, so C (is)

ιγ'.

Δύο δοθεισῶν εὐθειῶν μέσην ἀνάλογον προσευρεῖν.



Ἐστωσαν αἱ δοθεῖσαι δύο εὐθεῖαι αἱ AB , $BΓ$. δεῖ δὴ τῶν AB , $BΓ$ μέσην ἀνάλογον προσευρεῖν.


Κείσθωσαν ἐπ' εὐθείας, καὶ γεγράφθω ἐπὶ τῆς AG ἡμικύκλιον τὸ $AΔΓ$, καὶ ἤχθω ἀπὸ τοῦ B σημείου τῆς AG εὐθείας πρὸς ὀρθὰς ἢ BA , καὶ ἐπεξεύχθωσαν αἱ $AΔ$, $ΔΓ$.

Ἐπεὶ ἐν ἡμικυκλίῳ γωνία ἐστὶν ἡ ὑπὸ $AΔΓ$, ὀρθή ἐστίν. καὶ ἐπεὶ ἐν ὀρθογωνίῳ τριγώνῳ τῷ $AΔΓ$ ἀπὸ τῆς ὀρθῆς γωνίας ἐπὶ τὴν βάσιν κάθετος ἤκται ἡ $ΔB$, ἡ $ΔB$ ἄρα τῶν τῆς βάσεως τμημάτων τῶν AB , $BΓ$ μέση ἀνάλογόν ἐστίν.

Δύο ἄρα δοθεισῶν εὐθειῶν τῶν AB , $BΓ$ μέση ἀνάλογον προσεύρηται ἡ $ΔB$. ὅπερ ἔδει ποιῆσαι.

Proposition 13

To find the (straight-line) in mean proportion to two given straight-lines.†



Let AB and BC be the two given straight-lines. So it is required to find the (straight-line) in mean proportion to AB and BC .

Let (AB and BC) be laid down straight-on (with respect to one another), and let the semi-circle ADC have been drawn on AC [Prop. 1.10]. And let BD have been drawn from (point) B , at right-angles to AC [Prop. 1.11]. And let AD and DC have been joined.

And since ADC is an angle in a semi-circle, it is a right-angle [Prop. 3.31]. And since, in the right-angled triangle ADC , the (straight-line) DB has been drawn from the right-angle perpendicular to the base, DB is thus the mean proportional to the pieces of the base, AB and BC [Prop. 6.8 corr.].

Thus, DB has been found (which is) in mean proportion to the two given straight-lines, AB and BC . (Which is) the very thing it was required to do.

† In other words, to find the geometric mean of two given straight-lines.

ιδ'.


Τῶν ἴσων τε καὶ ἰσογωνίων παραλληλογράμμων ἀντιπεπόνθασιν αἱ πλευραὶ αἱ περὶ τὰς ἴσας γωνίας· καὶ ὧν ἰσογωνίων παραλληλογράμμων ἀντιπεπόνθασιν αἱ πλευραὶ αἱ περὶ τὰς ἴσας γωνίας, ἴσα ἐστὶν ἐκεῖνα.

Ἐστω ἴσα τε καὶ ἰσογώνια παραλληλόγραμμα τὰ AB , $BΓ$ ἴσας ἔχοντα τὰς πρὸς τῷ B γωνίας, καὶ κείσθωσαν ἐπ' εὐθείας αἱ $ΔB$, BE . ἐπ' εὐθείας ἄρα εἰσὶ καὶ αἱ ZB , BH . λέγω, ὅτι τῶν AB , $BΓ$ ἀντιπεπόνθασιν αἱ πλευραὶ αἱ περὶ τὰς ἴσας γωνίας, τουτέστιν, ὅτι ἐστὶν ὡς ἡ $ΔB$ πρὸς τὴν BE , οὕτως ἡ HB πρὸς τὴν BZ .

Proposition 14

In equal and equiangular parallelograms the sides about the equal angles are reciprocally proportional. And those equiangular parallelograms in which the sides about the equal angles are reciprocally proportional are equal.

Let AB and BC be equal and equiangular parallelograms having the angles at B equal. And let DB and BE be laid down straight-on (with respect to one another). Thus, FB and BG are also straight-on (with respect to one another) [Prop. 1.14]. I say that the sides of AB and



Συμπεπληρώσω γὰρ τὸ ZE παραλληλόγραμμον. ἐπεὶ οὖν ἴσον ἐστὶ τὸ AB παραλληλόγραμμον τῷ BG παραλληλόγραμμῳ, ἄλλο δὲ τι τὸ ZE, ἔστιν ἄρα ὡς τὸ AB πρὸς τὸ ZE, οὕτως τὸ BG πρὸς τὸ ZE. ἀλλ' ὡς μὲν τὸ AB πρὸς τὸ ZE, οὕτως ἡ ΔB πρὸς τὴν BE, ὡς δὲ τὸ BG πρὸς τὸ ZE, οὕτως ἡ HB πρὸς τὴν BZ· καὶ ὡς ἄρα ἡ ΔB πρὸς τὴν BE, οὕτως ἡ HB πρὸς τὴν BZ. τῶν ἄρα AB, BG παραλληλογράμμων ἀντιπεπόνθασιν αἱ πλευραὶ αἱ περὶ τὰς ἴσας γωνίας.

Ἀλλὰ δὴ ἔστω ὡς ἡ ΔB πρὸς τὴν BE, οὕτως ἡ HB πρὸς τὴν BZ· λέγω, ὅτι ἴσον ἐστὶ τὸ AB παραλληλόγραμμον τῷ BG παραλληλογράμμῳ.

Ἐπεὶ γάρ ἐστιν ὡς ἡ ΔB πρὸς τὴν BE, οὕτως ἡ HB πρὸς τὴν BZ, ἀλλ' ὡς μὲν ἡ ΔB πρὸς τὴν BE, οὕτως τὸ AB παραλληλόγραμμον πρὸς τὸ ZE παραλληλόγραμμον, ὡς δὲ ἡ HB πρὸς τὴν BZ, οὕτως τὸ BG παραλληλόγραμμον πρὸς τὸ ZE παραλληλόγραμμον, καὶ ὡς ἄρα τὸ AB πρὸς τὸ ZE, οὕτως τὸ BG πρὸς τὸ ZE· ἴσον ἄρα ἐστὶ τὸ AB παραλληλόγραμμον τῷ BG παραλληλογράμμῳ.


Τῶν ἄρα ἴσων τε καὶ ἰσογωνίων παραλληλογράμμων ἀντιπεπόνθασιν αἱ πλευραὶ αἱ περὶ τὰς ἴσας γωνίας· καὶ ὡν ἰσογωνίων παραλληλογράμμων ἀντιπεπόνθασιν αἱ πλευραὶ αἱ περὶ τὰς ἴσας γωνίας, ἴσα ἐστὶν ἐκεῖνα· ὅπερ ἔδει δεῖξαι.

ιε'.

Τῶν ἴσων καὶ μίαν μᾶ ἴσην ἐχόντων γωνίαν τριγῶνων ἀντιπεπόνθασιν αἱ πλευραὶ αἱ περὶ τὰς ἴσας γωνίας· καὶ ὡν μίαν μᾶ ἴσην ἐχόντων γωνίαν τριγῶνων ἀντιπεπόνθασιν αἱ πλευραὶ αἱ περὶ τὰς ἴσας γωνίας, ἴσα ἐστὶν ἐκεῖνα.

Ἐστω ἴσα τρίγωνα τὰ ABΓ, AΔE μίαν μᾶ ἴσην ἐχοντα γωνίαν τὴν ὑπὸ ΒΑΓ τῇ ὑπὸ ΔΑΕ· λέγω, ὅτι τῶν ABΓ, AΔE τριγῶνων ἀντιπεπόνθασιν αἱ πλευραὶ αἱ περὶ τὰς ἴσας γωνίας, τουτέστιν, ὅτι ἐστὶν ὡς ἡ ΓΑ πρὸς τὴν ΑΔ, οὕτως

BC about the equal angles are reciprocally proportional, that is to say, that as *DB* is to *BE*, so *GB* (is) to *BF*.



For let the parallelogram *FE* have been completed. Therefore, since parallelogram *AB* is equal to parallelogram *BC*, and *FE* (is) some other (parallelogram), thus as (parallelogram) *AB* is to *FE*, so (parallelogram) *BC* (is) to *FE* [Prop. 5.7]. But, as (parallelogram) *AB* (is) to *FE*, so *DB* (is) to *BE*, and as (parallelogram) *BC* (is) to *FE*, so *GB* (is) to *BF* [Prop. 6.1]. Thus, also, as *DB* (is) to *BE*, so *GB* (is) to *BF*. Thus, in parallelograms *AB* and *BC* the sides about the equal angles are reciprocally proportional.

And so, let *DB* be to *BE*, as *GB* (is) to *BF*. I say that parallelogram *AB* is equal to parallelogram *BC*.

For since as *DB* is to *BE*, so *GB* (is) to *BF*, but as *DB* (is) to *BE*, so parallelogram *AB* (is) to parallelogram *FE*, and as *GB* (is) to *BF*, so parallelogram *BC* (is) to parallelogram *FE* [Prop. 6.1], thus, also, as (parallelogram) *AB* (is) to *FE*, so (parallelogram) *BC* (is) to *FE* [Prop. 5.11]. Thus, parallelogram *AB* is equal to parallelogram *BC* [Prop. 5.9].


Thus, in equal and equiangular parallelograms the sides about the equal angles are reciprocally proportional. And those equiangular parallelograms in which the sides about the equal angles are reciprocally proportional are equal. (Which is) the very thing it was required to show.

Proposition 15

In equal triangles also having one angle equal to one (angle) the sides about the equal angles are reciprocally proportional. And those triangles having one angle equal to one angle for which the sides about the equal angles (are) reciprocally proportional are equal.

Let *ABC* and *ADE* be equal triangles having one angle equal to one (angle), (namely) *BAC* (equal) to *DAE*. I say that, in triangles *ABC* and *ADE*, the sides about the

ἡ EA πρὸς τὴν AB .



Κεῖσθω γὰρ ὥστε ἐπ' εὐθείας εἶναι τὴν GA τῆ AD · ἐπ' εὐθείας ἄρα ἐστὶ καὶ ἡ EA τῆ AB . καὶ ἐπεζεύχθω ἡ BD .

Ἐπεὶ οὖν ἴσον ἐστὶ τὸ ABG τρίγωνον τῷ ADE τριγώνῳ, ἄλλο δὲ τι τὸ BAD , ἔστιν ἄρα ὡς τὸ GAB τρίγωνον πρὸς τὸ BAD τρίγωνον, οὕτως τὸ EAD τρίγωνον πρὸς τὸ BAD τρίγωνον. ἀλλ' ὡς μὲν τὸ GAB πρὸς τὸ BAD , οὕτως ἡ GA πρὸς τὴν AD , ὡς δὲ τὸ EAD πρὸς τὸ BAD , οὕτως ἡ EA πρὸς τὴν AB . καὶ ὡς ἄρα ἡ GA πρὸς τὴν AD , οὕτως ἡ EA πρὸς τὴν AB . τῶν ABG , ADE ἄρα τριγώνων ἀντιπεπόνθασιν αἱ πλευραὶ αἱ περὶ τὰς ἴσας γωνίας.

Ἀλλὰ δὴ ἀντιπεπονηθέντων αἱ πλευραὶ τῶν ABG , ADE τριγώνων, καὶ ἔστω ὡς ἡ GA πρὸς τὴν AD , οὕτως ἡ EA πρὸς τὴν AB . λέγω, ὅτι ἴσον ἐστὶ τὸ ABG τρίγωνον τῷ ADE τριγώνῳ.


Ἐπιζευχθείσης γὰρ πάλιν τῆς BD , ἐπεὶ ἐστὶν ὡς ἡ GA πρὸς τὴν AD , οὕτως ἡ EA πρὸς τὴν AB , ἀλλ' ὡς μὲν ἡ GA πρὸς τὴν AD , οὕτως τὸ ABG τρίγωνον πρὸς τὸ BAD τρίγωνον, ὡς δὲ ἡ EA πρὸς τὴν AB , οὕτως τὸ EAD τρίγωνον πρὸς τὸ BAD τρίγωνον, ὡς ἄρα τὸ ABG τρίγωνον πρὸς τὸ BAD τρίγωνον, οὕτως τὸ EAD τρίγωνον πρὸς τὸ BAD τρίγωνον. ἐκάτερον ἄρα τῶν ABG , EAD πρὸς τὸ BAD τὸν αὐτὸν ἔχει λόγον. ἴσον ἄρα ἐστὶ τὸ ABG [τρίγωνον] τῷ EAD τριγώνῳ.

Τῶν ἄρα ἴσων καὶ μίαν μιᾶ ἴσην ἐχόντων γωνίαν τριγώνων ἀντιπεπόνθασιν αἱ πλευραὶ αἱ περὶ τὰς ἴσας γωνίας· καὶ ὡς μίαν μιᾶ ἴσην ἐχόντων γωνίαν τριγώνων ἀντιπεπόνθασιν αἱ πλευραὶ αἱ περὶ τὰς ἴσας γωνίας, ἐκεῖνα ἴσα ἐστὶν· ὅπερ ἔδει δεῖξαι.

15'.

Ἐὰν τέσσαρες εὐθεῖαι ἀνάλογον ᾖσιν, τὸ ὑπὸ τῶν ἄκρων περιεχόμενον ὀρθογώνιον ἴσον ἐστὶ τῷ ὑπὸ τῶν μέσων περιεχομένῳ ὀρθογώνιῳ· καὶ τὸ ὑπὸ τῶν ἄκρων

equal angles are reciprocally proportional, that is to say, that as CA is to AD , so EA (is) to AB .



For let CA be laid down so as to be straight-on (with respect) to AD . Thus, EA is also straight-on (with respect) to AB [Prop. 1.14]. And let BD have been joined.

Therefore, since triangle ABC is equal to triangle ADE , and BAD (is) some other (triangle), thus as triangle CAB is to triangle BAD , so triangle EAD (is) to triangle BAD [Prop. 5.7]. But, as (triangle) CAB (is) to BAD , so CA (is) to AD , and as (triangle) EAD (is) to BAD , so EA (is) to AB [Prop. 6.1]. And thus, as CA (is) to AD , so EA (is) to AB . Thus, in triangles ABC and ADE the sides about the equal angles (are) reciprocally proportional.

And so, let the sides of triangles ABC and ADE be reciprocally proportional, and (thus) let CA be to AD , as EA (is) to AB . I say that triangle ABC is equal to triangle ADE .


For, BD again being joined, since as CA is to AD , so EA (is) to AB , but as CA (is) to AD , so triangle ABC (is) to triangle BAD , and as EA (is) to AB , so triangle EAD (is) to triangle BAD [Prop. 6.1], thus as triangle ABC (is) to triangle BAD , so triangle EAD (is) to triangle BAD . Thus, (triangles) ABC and EAD each have the same ratio to BAD . Thus, [triangle] ABC is equal to triangle EAD [Prop. 5.9].

Thus, in equal triangles also having one angle equal to one (angle) the sides about the equal angles (are) reciprocally proportional. And those triangles having one angle equal to one angle for which the sides about the equal angles (are) reciprocally proportional are equal. (Which is) the very thing it was required to show.

Proposition 16

If four straight-lines are proportional then the rectangle contained by the (two) outermost is equal to the rectangle contained by the middle (two). And if the rect-

περιεχόμενον ὀρθογώνιον ἴσον ἢ τῶ ὑπὸ τῶν μέσων περιεχομένῳ ὀρθογώνιῳ, αἱ τέσσαρες εὐθεῖαι ἀνάλογον ἔσσονται.



Ἐστωσαν τέσσαρες εὐθεῖαι ἀνάλογον αἱ AB , $\Gamma\Delta$, E , Z , ὡς ἡ AB πρὸς τὴν $\Gamma\Delta$, οὕτως ἡ E πρὸς τὴν Z : λέγω, ὅτι τὸ ὑπὸ τῶν AB , Z περιεχόμενον ὀρθογώνιον ἴσον ἐστὶ τῶ ὑπὸ τῶν $\Gamma\Delta$, E περιεχομένῳ ὀρθογώνιῳ.

Ἦχθωσαν [γὰρ] ἀπὸ τῶν A , Γ σημείων ταῖς AB , $\Gamma\Delta$ εὐθείαις πρὸς ὀρθὰς αἱ AH , $\Gamma\Theta$, καὶ κείσθω τῇ μὲν Z ἴση ἡ AH , τῇ δὲ E ἴση ἡ $\Gamma\Theta$. καὶ συμπληρώσω τὰ BH , $\Delta\Theta$ παραλληλόγραμμα.


Καὶ ἐπεὶ ἐστὶν ὡς ἡ AB πρὸς τὴν $\Gamma\Delta$, οὕτως ἡ E πρὸς τὴν Z , ἴση δὲ ἡ μὲν E τῇ $\Gamma\Theta$, ἡ δὲ Z τῇ AH , ἐστὶν ἄρα ὡς ἡ AB πρὸς τὴν $\Gamma\Delta$, οὕτως ἡ $\Gamma\Theta$ πρὸς τὴν AH . τῶν BH , $\Delta\Theta$ ἄρα παραλληλογράμμων ἀντιπεπόνθασιν αἱ πλευραὶ αἱ περὶ τὰς ἴσας γωνίας. ὦν δὲ ἰσογωνίων παραλληλογράμμων ἀντιπεπόνθασιν αἱ πλευραὶ αἱ περὶ τὰς ἴσας γωνίας, ἴσα ἐστὶν ἐκεῖνα: ἴσον ἄρα ἐστὶ τὸ BH παραλληλόγραμμον τῶ $\Delta\Theta$ παραλληλογράμμῳ. καὶ ἐστὶ τὸ μὲν BH τὸ ὑπὸ τῶν AB , Z : ἴση γὰρ ἡ AH τῇ Z : τὸ δὲ $\Delta\Theta$ τὸ ὑπὸ τῶν $\Gamma\Delta$, E : ἴση γὰρ ἡ E τῇ $\Gamma\Theta$: τὸ ἄρα ὑπὸ τῶν AB , Z περιεχόμενον ὀρθογώνιον ἴσον ἐστὶ τῶ ὑπὸ τῶν $\Gamma\Delta$, E περιεχομένῳ ὀρθογώνιῳ.

Ἄλλὰ δὴ τὸ ὑπὸ τῶν AB , Z περιεχόμενον ὀρθογώνιον ἴσον ἔστω τῶ ὑπὸ τῶν $\Gamma\Delta$, E περιεχομένῳ ὀρθογώνιῳ. λέγω, ὅτι αἱ τέσσαρες εὐθεῖαι ἀνάλογον ἔσσονται, ὡς ἡ AB πρὸς τὴν $\Gamma\Delta$, οὕτως ἡ E πρὸς τὴν Z .

Τῶν γὰρ αὐτῶν κατασκευασθέντων, ἐπεὶ τὸ ὑπὸ τῶν AB , Z ἴσον ἐστὶ τῶ ὑπὸ τῶν $\Gamma\Delta$, E , καὶ ἐστὶ τὸ μὲν ὑπὸ τῶν AB , Z τὸ BH : ἴση γὰρ ἐστὶν ἡ AH τῇ Z : τὸ δὲ ὑπὸ τῶν $\Gamma\Delta$, E τὸ $\Delta\Theta$: ἴση γὰρ ἡ $\Gamma\Theta$ τῇ E : τὸ ἄρα BH ἴσον ἐστὶ τῶ $\Delta\Theta$. καὶ ἐστὶν ἰσογώνια. τῶν δὲ ἴσων καὶ ἰσογωνίων παραλληλογράμμων ἀντιπεπόνθασιν αἱ πλευραὶ αἱ περὶ τὰς ἴσας γωνίας. ἐστὶν ἄρα ὡς ἡ AB πρὸς τὴν $\Gamma\Delta$, οὕτως ἡ $\Gamma\Theta$ πρὸς τὴν AH . ἴση δὲ ἡ μὲν $\Gamma\Theta$ τῇ E , ἡ δὲ AH τῇ Z : ἐστὶν ἄρα ὡς ἡ AB πρὸς τὴν $\Gamma\Delta$, οὕτως ἡ E πρὸς τὴν Z .

Ἐὰν ἄρα τέσσαρες εὐθεῖαι ἀνάλογον ᾖσιν, τὸ ὑπὸ τῶν ἄκρων περιεχόμενον ὀρθογώνιον ἴσον ἐστὶ τῶ ὑπὸ τῶν μέσων περιεχομένῳ ὀρθογώνιῳ: καὶ τὸ ὑπὸ τῶν ἄκρων περιεχόμενον ὀρθογώνιον ἴσον ἢ τῶ ὑπὸ τῶν μέσων περιεχομένῳ ὀρθογώνιῳ, αἱ τέσσαρες εὐθεῖαι ἀνάλογον ἔσσονται: ὅπερ ἔδει δεῖξαι.

angle contained by the (two) outermost is equal to the rectangle contained by the middle (two) then the four straight-lines will be proportional.



Let AB , CD , E , and F be four proportional straight-lines, (such that) as AB (is) to CD , so E (is) to F . I say that the rectangle contained by AB and F is equal to the rectangle contained by CD and E .

[For] let AG and CH have been drawn from points A and C at right-angles to the straight-lines AB and CD (respectively) [Prop. 1.11]. And let AG be made equal to F , and CH to E [Prop. 1.3]. And let the parallelograms BG and DH have been completed.

And since as AB is to CD , so E (is) to F , and E (is) equal CH , and F to AG , thus as AB is to CD , so CH (is) to AG . Thus, in the parallelograms BG and DH the sides about the equal angles are reciprocally proportional. And those equiangular parallelograms in which the sides about the equal angles are reciprocally proportional are equal [Prop. 6.14]. Thus, parallelogram BG is equal to parallelogram DH . And BG is the (rectangle contained) by AB and F . For AG (is) equal to F . And DH (is) the (rectangle contained) by CD and E . For E (is) equal to CH . Thus, the rectangle contained by AB and F is equal to the rectangle contained by CD and E .


And so, let the rectangle contained by AB and F be equal to the rectangle contained by CD and E . I say that the four straight-lines will be proportional, (so that) as AB (is) to CD , so E (is) to F .

For, with the same construction, since the (rectangle contained) by AB and F is equal to the (rectangle contained) by CD and E . And BG is the (rectangle contained) by AB and F . For AG is equal to F . And DH (is) the (rectangle contained) by CD and E . For CH (is) equal to E . BG is thus equal to DH . And they are equiangular. And in equal and equiangular parallelograms the sides about the equal angles are reciprocally proportional [Prop. 6.14]. Thus, as AB is to CD , so CH (is) to AG . And CH (is) equal to E , and AG to F . Thus, as AB is to CD , so E (is) to F .

Thus, if four straight-lines are proportional then the rectangle contained by the (two) outermost is equal to the rectangle contained by the middle (two). And if the rectangle contained by the (two) outermost is equal to

ιζ'.

Ἐάν τρεῖς εὐθεῖαι ἀνάλογον ὦσιν, τὸ ὑπὸ τῶν ἄκρων περιεχόμενον ὀρθογώνιον ἴσον ἐστὶ τῷ ἀπὸ τῆς μέσης τετραγώνῳ· καὶ τὸ ὑπὸ τῶν ἄκρων περιεχόμενον ὀρθογώνιον ἴσον ἢ τῷ ἀπὸ τῆς μέσης τετραγώνῳ, αἱ τρεῖς εὐθεῖαι ἀνάλογον ἔσσονται.



Ἐστωσαν τρεῖς εὐθεῖαι ἀνάλογον αἱ A, B, Γ , ὡς ἡ A πρὸς τὴν B , οὕτως ἡ B πρὸς τὴν Γ . λέγω, ὅτι τὸ ὑπὸ τῶν A, Γ περιεχόμενον ὀρθογώνιον ἴσον ἐστὶ τῷ ἀπὸ τῆς B τετραγώνῳ.

Κείσθω τῆ B ἴση ἡ Δ .

Καὶ ἐπεὶ ἐστὶν ὡς ἡ A πρὸς τὴν B , οὕτως ἡ B πρὸς τὴν Γ , ἴση δὲ ἡ B τῆ Δ , ἔστιν ἄρα ὡς ἡ A πρὸς τὴν B , ἡ Δ πρὸς τὴν Γ . ἐὰν δὲ τέσσαρες εὐθεῖαι ἀνάλογον ὦσιν, τὸ ὑπὸ τῶν ἄκρων περιεχόμενον [ὀρθογώνιον] ἴσον ἐστὶ τῷ ὑπὸ τῶν μέσων περιεχομένῳ ὀρθογωνίῳ. τὸ ἄρα ὑπὸ τῶν A, Γ ἴσον ἐστὶ τῷ ὑπὸ τῶν B, Δ . ἀλλὰ τὸ ὑπὸ τῶν B, Δ τὸ ἀπὸ τῆς B ἐστὶν ἴση γὰρ ἡ B τῆ Δ . τὸ ἄρα ὑπὸ τῶν A, Γ περιεχόμενον ὀρθογώνιον ἴσον ἐστὶ τῷ ἀπὸ τῆς B τετραγώνῳ.

Ἀλλὰ δὴ τὸ ὑπὸ τῶν A, Γ ἴσον ἔστω τῷ ἀπὸ τῆς B . λέγω, ὅτι ἐστὶν ὡς ἡ A πρὸς τὴν B , οὕτως ἡ B πρὸς τὴν Γ .


Τῶν γὰρ αὐτῶν κατασκευασθέντων, ἐπεὶ τὸ ὑπὸ τῶν A, Γ ἴσον ἐστὶ τῷ ἀπὸ τῆς B , ἀλλὰ τὸ ἀπὸ τῆς B τὸ ὑπὸ τῶν B, Δ ἐστὶν ἴση γὰρ ἡ B τῆ Δ . τὸ ἄρα ὑπὸ τῶν A, Γ ἴσον ἐστὶ τῷ ὑπὸ τῶν B, Δ . ἐὰν δὲ τὸ ὑπὸ τῶν ἄκρων ἴσον ἢ τῷ ὑπὸ τῶν μέσων, αἱ τέσσαρες εὐθεῖαι ἀνάλογον εἰσιν. ἔστιν ἄρα ὡς ἡ A πρὸς τὴν B , οὕτως ἡ Δ πρὸς τὴν Γ . ἴση δὲ ἡ B τῆ Δ . ὡς ἄρα ἡ A πρὸς τὴν B , οὕτως ἡ B πρὸς τὴν Γ .

Ἐὰν ἄρα τρεῖς εὐθεῖαι ἀνάλογον ὦσιν, τὸ ὑπὸ τῶν ἄκρων περιεχόμενον ὀρθογώνιον ἴσον ἐστὶ τῷ ἀπὸ τῆς μέσης τετραγώνῳ· καὶ τὸ ὑπὸ τῶν ἄκρων περιεχόμενον ὀρθογώνιον ἴσον ἢ τῷ ἀπὸ τῆς μέσης τετραγώνῳ, αἱ τρεῖς εὐθεῖαι ἀνάλογον ἔσσονται· ὅπερ ἔδει δεῖξαι.

the rectangle contained by the middle (two) then the four straight-lines will be proportional. (Which is) the very thing it was required to show.

Proposition 17

If three straight-lines are proportional then the rectangle contained by the (two) outermost is equal to the square on the middle (one). And if the rectangle contained by the (two) outermost is equal to the square on the middle (one) then the three straight-lines will be proportional.



Let A, B and C be three proportional straight-lines, (such that) as A (is) to B , so B (is) to C . I say that the rectangle contained by A and C is equal to the square on B .

Let D be made equal to B [Prop. 1.3].

And since as A is to B , so B (is) to C , and B (is) equal to D , thus as A is to B , (so) D (is) to C . And if four straight-lines are proportional then the [rectangle] contained by the (two) outermost is equal to the rectangle contained by the middle (two) [Prop. 6.16]. Thus, the (rectangle contained) by A and C is equal to the (rectangle contained) by B and D . But, the (rectangle contained) by B and D is the (square) on B . For B (is) equal to D . Thus, the rectangle contained by A and C is equal to the square on B .


And so, let the (rectangle contained) by A and C be equal to the (square) on B . I say that as A is to B , so B (is) to C .

For, with the same construction, since the (rectangle contained) by A and C is equal to the (square) on B . But, the (square) on B is the (rectangle contained) by B and D . For B (is) equal to D . The (rectangle contained) by A and C is thus equal to the (rectangle contained) by B and D . And if the (rectangle contained) by the (two) outermost is equal to the (rectangle contained) by the middle (two) then the four straight-lines are proportional [Prop. 6.16]. Thus, as A is to B , so D (is) to C . And B (is) equal to D . Thus, as A (is) to B , so B (is) to C .

Thus, if three straight-lines are proportional then the rectangle contained by the (two) outermost is equal to the square on the middle (one). And if the rectangle contained by the (two) outermost is equal to the square on the middle (one) then the three straight-lines will be proportional. (Which is) the very thing it was required to

ιη'.

Ἀπὸ τῆς δοθείσης εὐθείας τῷ δοθέντι εὐθυγράμμῳ ὁμοίον τε καὶ ὁμοίως κείμενον εὐθυγράμμον ἀναγράψαι.



Ἐστω ἡ μὲν δοθεῖσα εὐθεῖα ἡ ΑΒ, τὸ δὲ δοθὲν εὐθυγράμμον τὸ ΓΕ· δεῖ δὴ ἀπὸ τῆς ΑΒ εὐθείας τῷ ΓΕ εὐθυγράμμῳ ὁμοίον τε καὶ ὁμοίως κείμενον εὐθυγράμμον ἀναγράψαι.


Ἐπεζεύχθω ἡ ΔΖ, καὶ συνεστάτω πρὸς τῇ ΑΒ εὐθείᾳ καὶ τοῖς πρὸς αὐτῇ σημείοις τοῖς Α, Β τῇ μὲν πρὸς τῷ Γ γωνία ἴση ἢ ὑπὸ ΗΑΒ, τῇ δὲ ὑπὸ ΓΔΖ ἴση ἢ ὑπὸ ΑΒΗ. λοιπὴ ἄρα ἢ ὑπὸ ΓΖΔ τῇ ὑπὸ ΑΗΒ ἐστὶν ἴση· ἰσογώνιον ἄρα ἐστὶ τὸ ΖΓΔ τρίγωνον τῷ ΗΑΒ τριγώνῳ. ἀνάλογον ἄρα ἐστὶν ὡς ἡ ΖΔ πρὸς τὴν ΗΒ, οὕτως ἡ ΖΓ πρὸς τὴν ΗΑ, καὶ ἡ ΓΔ πρὸς τὴν ΑΒ. πάλιν συνεστάτω πρὸς τῇ ΒΗ εὐθείᾳ καὶ τοῖς πρὸς αὐτῇ σημείοις τοῖς Β, Η τῇ μὲν ὑπὸ ΔΖΕ γωνία ἴση ἢ ὑπὸ ΒΗΘ, τῇ δὲ ὑπὸ ΖΔΕ ἴση ἢ ὑπὸ ΗΒΘ. λοιπὴ ἄρα ἢ πρὸς τῷ Ε λοιπῇ τῇ πρὸς τῷ Θ ἐστὶν ἴση· ἰσογώνιον ἄρα ἐστὶ τὸ ΖΔΕ τρίγωνον τῷ ΗΘΒ τριγώνῳ· ἀνάλογον ἄρα ἐστὶν ὡς ἡ ΖΔ πρὸς τὴν ΗΒ, οὕτως ἡ ΖΕ πρὸς τὴν ΗΘ καὶ ἡ ΕΔ πρὸς τὴν ΘΒ. ἐδείχθη δὲ καὶ ὡς ἡ ΖΔ πρὸς τὴν ΗΒ, οὕτως ἡ ΖΓ πρὸς τὴν ΗΑ καὶ ἡ ΓΔ πρὸς τὴν ΑΒ· καὶ ὡς ἄρα ἡ ΖΓ πρὸς τὴν ΗΑ, οὕτως ἡ τε ΓΔ πρὸς τὴν ΑΒ καὶ ἡ ΖΕ πρὸς τὴν ΗΘ καὶ ἔτι ἡ ΕΑ πρὸς τὴν ΘΒ. καὶ ἐπεὶ ἴση ἐστὶν ἡ μὲν ὑπὸ ΓΖΔ γωνία τῇ ὑπὸ ΑΗΒ, ἡ δὲ ὑπὸ ΔΖΕ τῇ ὑπὸ ΒΗΘ, ὅλη ἄρα ἢ ὑπὸ ΓΖΕ ὅλη τῇ ὑπὸ ΑΗΘ ἐστὶν ἴση. διὰ τὰ αὐτὰ δὴ καὶ ἡ ὑπὸ ΓΔΕ τῇ ὑπὸ ΑΒΘ ἐστὶν ἴση. ἔστι δὲ καὶ ἡ μὲν πρὸς τῷ Γ τῇ πρὸς τῷ Α ἴση, ἡ δὲ πρὸς τῷ Ε τῇ πρὸς τῷ Θ. ἰσογώνιον ἄρα ἐστὶ τὸ ΑΘ τῷ ΓΕ· καὶ τὰς περὶ τὰς ἴσας γωνίας αὐτῶν πλευρὰς ἀνάλογον ἔχει· ὁμοιον ἄρα ἐστὶ τὸ ΑΘ εὐθυγράμμον τῷ ΓΕ εὐθυγράμμῳ.

Ἀπὸ τῆς δοθείσης ἄρα εὐθείας τῆς ΑΒ τῷ δοθέντι εὐθυγράμμῳ τῷ ΓΕ ὁμοίον τε καὶ ὁμοίως κείμενον εὐθυγράμμον ἀναγράφεται τὸ ΑΘ· ὅπερ ἔδει ποιῆσαι.

show.

Proposition 18

To describe a rectilinear figure similar, and similarly laid down, to a given rectilinear figure on a given straight-line.




Let ΑΒ be the given straight-line, and CΕ the given rectilinear figure. So it is required to describe a rectilinear figure similar, and similarly laid down, to the rectilinear figure CΕ on the straight-line ΑΒ.

Let DF have been joined, and let GAB, equal to the angle at C, and ABG, equal to (angle) CDF, have been constructed on the straight-line ΑΒ at the points Α and Β on it (respectively) [Prop. 1.23]. Thus, the remaining (angle) CFD is equal to AGB [Prop. 1.32]. Thus, triangle FCD is equiangular to triangle GAB. Thus, proportionally, as FD is to GB, so FC (is) to GA, and CD to AB [Prop. 6.4]. Again, let BGH, equal to angle DFE, and GBH equal to (angle) FDE, have been constructed on the straight-line ΒΓ at the points G and Β on it (respectively) [Prop. 1.23]. Thus, the remaining (angle) at E is equal to the remaining (angle) at H [Prop. 1.32]. Thus, triangle FDE is equiangular to triangle GHB. Thus, proportionally, as FD is to GB, so FE (is) to GH, and ED to HB [Prop. 6.4]. And it was also shown (that) as FD (is) to GB, so FC (is) to GA, and CD to AB. Thus, also, as FC (is) to AG, so CD (is) to AB, and FE to GH, and, further, ED to HB. And since angle CFD is equal to AGB, and DFE to BGH, thus the whole (angle) CFE is equal to the whole (angle) AGH. So, for the same (reasons), (angle) CDE is also equal to ABH. And the (angle) at C is also equal to the (angle) at Α, and the (angle) at E to the (angle) at Η. Thus, (figure) ΑΗ is equiangular to CΕ. And (the two figures) have the sides about their equal angles proportional. Thus, the rectilinear figure ΑΗ is similar to the rectilinear figure CΕ [Def. 6.1].

Thus, the rectilinear figure ΑΗ, similar, and similarly laid down, to the given rectilinear figure CΕ has been constructed on the given straight-line ΑΒ. (Which is) the

ιθ'.

Τὰ ὅμοια τρίγωνα πρὸς ἀλλήλα ἐν διπλασίονι λόγῳ ἐστὶ τῶν ὁμολόγων πλευρῶν.



Ἐστω ὅμοια τρίγωνα τὰ ABG , ΔEZ ἴσην ἔχοντα τὴν πρὸς τῷ B γωνίαν τῇ πρὸς τῷ E , ὡς δὲ τὴν AB πρὸς τὴν BG , οὕτως τὴν ΔE πρὸς τὴν EZ , ὥστε ὁμόλογον εἶναι τὴν BG τῇ EZ : λέγω, ὅτι τὸ ABG τρίγωνον πρὸς τὸ ΔEZ τρίγωνον διπλασίονα λόγον ἔχει ἢ περὶ ἢ BG πρὸς τὴν EZ .

Εἰλήφθω γὰρ τῶν BG , EZ τρίτη ἀνάλογον ἢ BH , ὥστε εἶναι ὡς τὴν BG πρὸς τὴν EZ , οὕτως τὴν EZ πρὸς τὴν BH : καὶ ἐπεζεύχθω ἢ AH .

Ἐπεὶ οὖν ἐστὶν ὡς ἢ AB πρὸς τὴν BG , οὕτως ἢ ΔE πρὸς τὴν EZ , ἐναλλάξ ἄρα ἐστὶν ὡς ἢ AB πρὸς τὴν ΔE , οὕτως ἢ BG πρὸς τὴν EZ . ἀλλ' ὡς ἢ BG πρὸς τὴν EZ , οὕτως ἐστὶν ἢ EZ πρὸς BH . καὶ ὡς ἄρα ἢ AB πρὸς ΔE , οὕτως ἢ EZ πρὸς BH : τῶν ABH , ΔEZ ἄρα τριγώνων ἀντιπεπόνθησιν αἱ πλευραὶ αἱ περὶ τὰς ἴσας γωνίας. ὣν δὲ μίαν μὲν ἴσην ἔχοντων γωνίαν τριγώνων ἀντιπεπόνθησιν αἱ πλευραὶ αἱ περὶ τὰς ἴσας γωνίας, ἴσα ἐστὶν ἐκεῖνα. ἴσον ἄρα ἐστὶ τὸ ABH τρίγωνον τῷ ΔEZ τριγώνῳ. καὶ ἐπεὶ ἐστὶν ὡς ἢ BG πρὸς τὴν EZ , οὕτως ἢ EZ πρὸς τὴν BH , ἐὰν δὲ τρεῖς εὐθεῖαι ἀνάλογον ὦσιν, ἢ πρώτη πρὸς τὴν τρίτην διπλασίονα λόγον ἔχει ἢ περὶ πρὸς τὴν δευτέραν, ἢ BG ἄρα πρὸς τὴν BH διπλασίονα λόγον ἔχει ἢ περὶ ἢ GB πρὸς τὴν EZ . ὡς δὲ ἢ GB πρὸς τὴν BH , οὕτως τὸ ABG τρίγωνον πρὸς τὸ ABH τρίγωνον: καὶ τὸ ABG ἄρα τρίγωνον πρὸς τὸ ABH διπλασίονα λόγον ἔχει ἢ περὶ ἢ BG πρὸς τὴν EZ . ἴσον δὲ τὸ ABH τρίγωνον τῷ ΔEZ τριγώνῳ. καὶ τὸ ABG ἄρα τρίγωνον πρὸς τὸ ΔEZ τρίγωνον διπλασίονα λόγον ἔχει ἢ περὶ ἢ BG πρὸς τὴν EZ .

Τὰ ἄρα ὅμοια τρίγωνα πρὸς ἀλλήλα ἐν διπλασίονι λόγῳ ἐστὶ τῶν ὁμολόγων πλευρῶν. [ἔπερ ἔδει δεῖξαι.]


Πόρισμα.

Ἐκ δὴ τούτου φανερόν, ὅτι, ἐὰν τρεῖς εὐθεῖαι ἀνάλογον ὦσιν, ἐστὶν ὡς ἢ πρώτη πρὸς τὴν τρίτην, οὕτως τὸ ἀπό

very thing it was required to do.

Proposition 19

Similar triangles are to one another in the squared ratio of (their) corresponding sides.



Let ABC and DEF be similar triangles having the angle at B equal to the (angle) at E , and AB to BC , as DE (is) to EF , such that BC corresponds to EF . I say that triangle ABC has a squared ratio to triangle DEF with respect to (that side) BC (has) to EF .

For let a third (straight-line), BG , have been taken (which is) proportional to BC and EF , so that as BC (is) to EF , so EF (is) to BG [Prop. 6.11]. And let AG have been joined.

Therefore, since as AB is to BC , so DE (is) to EF , thus, alternately, as AB is to DE , so BC (is) to EF [Prop. 5.16]. But, as BC (is) to EF , so EF is to BG . And, thus, as AB (is) to DE , so EF (is) to BG . Thus, for triangles ABG and DEF , the sides about the equal angles are reciprocally proportional. And those triangles having one (angle) equal to one (angle) for which the sides about the equal angles are reciprocally proportional are equal [Prop. 6.15]. Thus, triangle ABG is equal to triangle DEF . And since as BC (is) to EF , so EF (is) to BG , and if three straight-lines are proportional then the first has a squared ratio to the third with respect to the second [Def. 5.9], BC thus has a squared ratio to BG with respect to (that) CB (has) to EF . And as CB (is) to BG , so triangle ABC (is) to triangle ABG [Prop. 6.1]. Thus, triangle ABC also has a squared ratio to (triangle) ABG with respect to (that side) BC (has) to EF . And triangle ABG (is) equal to triangle DEF . Thus, triangle ABC also has a squared ratio to triangle DEF with respect to (that side) BC (has) to EF .

Thus, similar triangles are to one another in the squared ratio of (their) corresponding sides. [(Which is) the very thing it was required to show].

Corollary

So it is clear, from this, that if three straight-lines are proportional, then as the first is to the third, so the figure

τῆς πρώτης εἶδος πρὸς τὸ ἀπὸ τῆς δευτέρας τὸ ὅμοιον καὶ ὁμοίως ἀναγραφόμενον. ὅπερ εἶδει δεῖξαι.

(described) on the first (is) to the similar, and similarly described, (figure) on the second. (Which is) the very thing it was required to show.





† Literally, "double".

κ'.

Proposition 20

Τὰ ὅμοια πολύγωνα εἰς τε ὅμοια τρίγωνα διαιρεῖται καὶ εἰς ἴσα τὸ πλῆθος καὶ ὁμόλογα τοῖς ὅλοις, καὶ τὸ πολύγωνον πρὸς τὸ πολύγωνον διπλασίονα λόγον ἔχει ἤπερ ἡ ὁμόλογος πλευρὰ πρὸς τὴν ὁμόλογον πλευράν.

Similar polygons can be divided into equal numbers of similar triangles corresponding (in proportion) to the wholes, and one polygon has to the (other) polygon a squared ratio with respect to (that) a corresponding side (has) to a corresponding side.



Ἐστω ὅμοια πολύγωνα τὰ $ABΓΔΕ$, $ZHΘΚΛ$, ὁμόλογος δὲ ἔστω ἡ AB τῆ ZH . λέγω, ὅτι τὰ $ABΓΔΕ$, $ZHΘΚΛ$ πολύγωνα εἰς τε ὅμοια τρίγωνα διαιρεῖται καὶ εἰς ἴσα τὸ πλῆθος καὶ ὁμόλογα τοῖς ὅλοις, καὶ τὸ $ABΓΔΕ$ πολύγωνον πρὸς τὸ $ZHΘΚΛ$ πολύγωνον διπλασίονα λόγον ἔχει ἤπερ ἡ AB πρὸς τὴν ZH .

Let $ABCDE$ and $FGHKL$ be similar polygons, and let AB correspond to FG . I say that polygons $ABCDE$ and $FGHKL$ can be divided into equal numbers of similar triangles corresponding (in proportion) to the wholes, and (that) polygon $ABCDE$ has a squared ratio to polygon $FGHKL$ with respect to that AB (has) to FG .

Ἐπεξεύχθωσαν αἱ BE , $ΕΓ$, $ΗΛ$, $ΛΘ$.

Let BE , EC , GL , and LH have been joined.

Καὶ ἐπεὶ ὁμοίον ἔστι τὸ $ABΓΔΕ$ πολύγωνον τῷ $ZHΘΚΛ$ πολυγώνῳ, ἴση ἔστιν ἡ ὑπὸ BAE γωνία τῆ ὑπὸ $HZΛ$. καὶ ἔστιν ὡς ἡ BA πρὸς AE , οὕτως ἡ HZ πρὸς $ZΛ$. ἐπεὶ οὖν δύο τρίγωνά ἐστι τὰ ABE , ZHA μίαν γωνίαν μιᾶ γωνία ἴσην ἔχοντα, περὶ δὲ τὰς ἴσας γωνίας τὰς πλευρὰς ἀνάλογον, ἰσογώνιον ἄρα ἔστι τὸ ABE τρίγωνον τῷ ZHA τριγώνῳ· ὥστε καὶ ὁμοίον· ἴση ἄρα ἔστιν ἡ ὑπὸ ABE γωνία τῆ ὑπὸ ZHA . ἔστι δὲ καὶ ὅλη ἡ ὑπὸ $ABΓ$ ὅλη τῆ ὑπὸ $ZHΘ$ ἴση διὰ τὴν ὁμοιότητα τῶν πολυγώνων· λοιπὴ ἄρα ἡ ὑπὸ $EBΓ$ γωνία τῆ ὑπὸ $ΛΗΘ$ ἔστιν ἴση. καὶ ἐπεὶ διὰ τὴν ὁμοιότητα τῶν ABE , ZHA τριγώνων ἔστιν ὡς ἡ EB πρὸς BA , οὕτως ἡ $ΛΗ$ πρὸς HZ , ἀλλὰ μὴν καὶ διὰ τὴν ὁμοιότητα τῶν πολυγώνων ἔστιν ὡς ἡ AB πρὸς $ΒΓ$, οὕτως ἡ ZH πρὸς $HΘ$, δι' ἴσου ἄρα ἔστιν ὡς ἡ EB πρὸς $ΒΓ$, οὕτως ἡ $ΛΗ$ πρὸς $HΘ$, καὶ περὶ τὰς ἴσας γωνίας τὰς ὑπὸ $EBΓ$, $ΛΗΘ$ αἱ πλευραὶ ἀνάλογόν εἰσιν· ἰσογώνιον ἄρα ἔστι τὸ $EBΓ$ τρίγωνον τῷ $ΛΗΘ$ τριγώνῳ· ὥστε καὶ ὁμοίον ἔστι τὸ $EBΓ$ τρίγωνον τῷ $ΛΗΘ$ τριγώνῳ. διὰ τὰ αὐτὰ δὴ καὶ τὸ $ΕΓΔ$ τρίγωνον ὁμοίον ἔστι τῷ $ΛΘΚ$ τριγώνῳ. τὰ ἄρα ὅμοια πολύγωνα τὰ $ABΓΔΕ$, $ZHΘΚΛ$ εἰς τε ὅμοια τρίγωνα διήρηται καὶ εἰς ἴσα

And since polygon $ABCDE$ is similar to polygon $FGHKL$, angle BAE is equal to angle GFL , and as BA is to AE , so GF (is) to FL [Def. 6.1]. Therefore, since ABE and FGL are two triangles having one angle equal to one angle and the sides about the equal angles proportional, triangle ABE is thus equiangular to triangle FGL [Prop. 6.6]. Hence, (they are) also similar [Prop. 6.4, Def. 6.1]. Thus, angle ABE is equal to (angle) FGL . And the whole (angle) ABC is equal to the whole (angle) FGH , on account of the similarity of the polygons. Thus, the remaining angle EBC is equal to LGH . And since, on account of the similarity of triangles ABE and FGL , as EB is to BA , so LG (is) to GF , but also, on account of the similarity of the polygons, as AB is to BC , so FG (is) to GH , thus, via equality, as EB is to BC , so LG (is) to GH [Prop. 5.22], and the sides about the equal angles, EBC and LGH , are proportional. Thus, triangle EBC is equiangular to triangle LGH [Prop. 6.6]. Hence, triangle EBC is also similar to triangle LGH [Prop. 6.4, Def. 6.1]. So, for the same (reasons), triangle ECD is also similar

τὸ πλῆθος.

Λέγω, ὅτι καὶ ὁμόλογα τοῖς ὅλοις, τούτέστιν ὥστε ἀνάλογον εἶναι τὰ τρίγωνα, καὶ ἡγούμενα μὲν εἶναι τὰ ABE , $EBΓ$, $ΕΓΔ$, ἐπόμενα δὲ αὐτῶν τὰ ZHA , $ΛΗΘ$, $ΛΘΚ$, καὶ ὅτι τὸ $ABΓΔΕ$ πολύγωνον πρὸς τὸ $ZHΘΚΛ$ πολύγωνον διπλασίονα λόγον ἔχει ἥπερ ἡ ὁμόλογος πλευρὰ πρὸς τὴν ὁμόλογον πλευράν, τούτέστιν ἡ AB πρὸς τὴν ZH .

Ἐπεζύχθωσαν γὰρ αἱ $ΑΓ$, $ΖΘ$. καὶ ἐπεὶ διὰ τὴν ὁμοιότητα τῶν πολυγώνων ἴση ἐστὶν ἡ ὑπὸ $ABΓ$ γωνία τῇ ὑπὸ $ZHΘ$, καὶ ἐστὶν ὡς ἡ AB πρὸς $BΓ$, οὕτως ἡ ZH πρὸς $HΘ$, ἰσογώνιον ἐστὶ τὸ $ABΓ$ τρίγωνον τῷ $ZHΘ$ τριγώνῳ· ἴση ἄρα ἐστὶν ἡ μὲν ὑπὸ BAG γωνία τῇ ὑπὸ $HZΘ$, ἡ δὲ ὑπὸ BGA τῇ ὑπὸ $HΘZ$. καὶ ἐπεὶ ἴση ἐστὶν ἡ ὑπὸ BAM γωνία τῇ ὑπὸ HZN , ἐστὶ δὲ καὶ ἡ ὑπὸ ABM τῇ ὑπὸ ZHN ἴση, καὶ λοιπὴ ἄρα ἡ ὑπὸ AMB λοιπὴ τῇ ὑπὸ ZNH ἴση ἐστὶν· ἰσογώνιον ἄρα ἐστὶ τὸ ABM τρίγωνον τῷ ZHN τριγώνῳ. ὁμοίως δὴ δεῖξομεν, ὅτι καὶ τὸ $BΜΓ$ τρίγωνον ἰσογώνιον ἐστὶ τῷ $HNΘ$ τριγώνῳ. ἀνάλογον ἄρα ἐστὶν, ὡς μὲν ἡ AM πρὸς MB , οὕτως ἡ ZN πρὸς NH , ὡς δὲ ἡ BM πρὸς $ΜΓ$, οὕτως ἡ HN πρὸς $NΘ$. ὥστε καὶ δι' ἴσου, ὡς ἡ AM πρὸς $ΜΓ$, οὕτως ἡ ZN πρὸς $NΘ$. ἀλλ' ὡς ἡ AM πρὸς $ΜΓ$, οὕτως τὸ ABM [τρίγωνον] πρὸς τὸ $MBΓ$, καὶ τὸ AME πρὸς τὸ $EMΓ$. πρὸς ἀλληλα γὰρ εἰσιν ὡς αἱ βάσεις. καὶ ὡς ἄρα ἐν τῶν ἡγουμένων πρὸς ἐν τῶν ἐπόμενων, οὕτως ἅπαντα τὰ ἡγούμενα πρὸς ἅπαντα τὰ ἐπόμενα· ὡς ἄρα τὸ AMB τρίγωνον πρὸς τὸ $BΜΓ$, οὕτως τὸ ABE πρὸς τὸ $ΓBE$. ἀλλ' ὡς τὸ AMB πρὸς τὸ $BΜΓ$, οὕτως ἡ AM πρὸς $ΜΓ$. καὶ ὡς ἄρα ἡ AM πρὸς $ΜΓ$, οὕτως τὸ ABE τρίγωνον πρὸς τὸ $EBΓ$ τρίγωνον. διὰ τὰ αὐτὰ δὴ καὶ ὡς ἡ ZN πρὸς $NΘ$, οὕτως τὸ ZHA τρίγωνον πρὸς τὸ $HAΘ$ τρίγωνον. καὶ ἐστὶν ὡς ἡ AM πρὸς $ΜΓ$, οὕτως ἡ ZN πρὸς $NΘ$. καὶ ὡς ἄρα τὸ ABE τρίγωνον πρὸς τὸ $BEΓ$ τρίγωνον, οὕτως τὸ ZHA τρίγωνον πρὸς τὸ $HAΘ$ τρίγωνον, καὶ ἐναλλάξ ὡς τὸ ABE τρίγωνον πρὸς τὸ ZHA τρίγωνον, οὕτως τὸ $BEΓ$ τρίγωνον πρὸς τὸ $HAΘ$ τρίγωνον. ὁμοίως δὴ δεῖξομεν ἐπιζευχθεισῶν τῶν $BΔ$, HK , ὅτι καὶ ὡς τὸ $BEΓ$ τρίγωνον πρὸς τὸ $ΛΗΘ$ τρίγωνον, οὕτως τὸ $ΕΓΔ$ τρίγωνον πρὸς τὸ $ΛΘΚ$ τρίγωνον. καὶ ἐπεὶ ἐστὶν ὡς τὸ ABE τρίγωνον πρὸς τὸ ZHA τρίγωνον, οὕτως τὸ $EBΓ$ πρὸς τὸ $ΛΗΘ$, καὶ ἔτι τὸ $ΕΓΔ$ πρὸς τὸ $ΛΘΚ$, καὶ ὡς ἄρα ἐν τῶν ἡγουμένων πρὸς ἐν τῶν ἐπομένων, οὕτως ἅπαντα τὰ ἡγούμενα πρὸς ἅπαντα τὰ ἐπόμενα· ἐστὶν ἄρα ὡς τὸ ABE τρίγωνον πρὸς τὸ ZHA τρίγωνον, οὕτως τὸ $ABΓΔΕ$ πολύγωνον πρὸς τὸ $ZHΘΚΛ$ πολύγωνον. ἀλλὰ τὸ ABE τρίγωνον πρὸς τὸ ZHA τρίγωνον διπλασίονα λόγον ἔχει ἥπερ ἡ AB ὁμόλογος πλευρὰ πρὸς τὴν ZH ὁμόλογον πλευράν· τὰ γὰρ ὅμοια τρίγωνα ἐν διπλασίονι λόγῳ ἐστὶ τῶν ὁμολόγων πλευρῶν. καὶ τὸ $ABΓΔΕ$ ἄρα πολύγωνον πρὸς τὸ $ZHΘΚΛ$ πολύγωνον διπλασίονα λόγον ἔχει ἥπερ ἡ AB ὁμόλογος πλευρὰ πρὸς τὴν ZH ὁμόλογον πλευράν.

Τὰ ἄρα ὅμοια πολύγωνα εἰς τε ὅμοια τρίγωνα διαιρεῖται καὶ εἰς ἴσα τὸ πλῆθος καὶ ὁμόλογα τοῖς ὅλοις, καὶ τὸ

to triangle LHK . Thus, the similar polygons $ABCDE$ and $FGHKL$ have been divided into equal numbers of similar triangles.

I also say that (the triangles) correspond (in proportion) to the wholes. That is to say, the triangles are proportional: ABE , EBC , and ECD are the leading (magnitudes), and their (associated) following (magnitudes are) FGL , LGH , and LHK (respectively). (I) also (say) that polygon $ABCDE$ has a squared ratio to polygon $FGHKL$ with respect to (that) a corresponding side (has) to a corresponding side—that is to say, (side) AB to FG .

For let AC and FH have been joined. And since angle ABC is equal to FGH , and as AB is to BC , so FG (is) to GH , on account of the similarity of the polygons, triangle ABC is equiangular to triangle FGH [Prop. 6.6]. Thus, angle BAC is equal to GFH , and (angle) BCA to GHF . And since angle BAM is equal to GFN , and (angle) ABM is also equal to FGN (see earlier), the remaining (angle) AMB is thus also equal to the remaining (angle) FNG [Prop. 1.32]. Thus, triangle ABM is equiangular to triangle FGN . So, similarly, we can show that triangle BMC is also equiangular to triangle GNH . Thus, proportionally, as AM is to MB , so FN (is) to NG , and as BM (is) to MC , so GN (is) to NH [Prop. 6.4]. Hence, also, via equality, as AM (is) to MC , so FN (is) to NH [Prop. 5.22]. But, as AM (is) to MC , so [triangle] ABM is to MBC , and AME to EMC . For they are to one another as their bases [Prop. 6.1]. And as one of the leading (magnitudes) is to one of the following (magnitudes), so (the sum of) all the leading (magnitudes) is to (the sum of) all the following (magnitudes) [Prop. 5.12]. Thus, as triangle AMB (is) to BMC , so (triangle) ABE (is) to CBE . But, as (triangle) AMB (is) to BMC , so AM (is) to MC . Thus, also, as AM (is) to MC , so triangle ABE (is) to triangle EBC . And so, for the same (reasons), as FN (is) to NH , so triangle FGL (is) to triangle GLH . And as AM is to MC , so FN (is) to NH . Thus, also, as triangle ABE (is) to triangle BEC , so triangle FGL (is) to triangle GLH , and, alternately, as triangle ABE (is) to triangle FGL , so triangle BEC (is) to triangle GLH [Prop. 5.16]. So, similarly, we can also show, by joining BD and GK , that as triangle BEC (is) to triangle LGH , so triangle ECD (is) to triangle LHK . And since as triangle ABE is to triangle FGL , so (triangle) EBC (is) to LGH , and, further, (triangle) ECD to LHK , and also as one of the leading (magnitudes) is to one of the following, so (the sum of) all the leading (magnitudes) is to (the sum of) all the following [Prop. 5.12], thus as triangle ABE is to triangle FGL , so polygon $ABCDE$ (is) to polygon $FGHKL$. But, triangle ABE has a squared ratio

πολύγωνον πρὸς τὸ πολύγωνον διπλασίονα λόγον ἔχει ἢπερ ἡ ὁμόλογος πλευρὰ πρὸς τὴν ὁμόλογον πλευρὰν [ὅπερ ἔδει δεῖξαι].

to triangle FGL with respect to (that) the corresponding side AB (has) to the corresponding side FG . For, similar triangles are in the squared ratio of corresponding sides [Prop. 6.14]. Thus, polygon $ABCDE$ also has a squared ratio to polygon $FGHKL$ with respect to (that) the corresponding side AB (has) to the corresponding side FG .

Thus, similar polygons can be divided into equal numbers of similar triangles corresponding (in proportion) to the wholes, and one polygon has to the (other) polygon a squared ratio with respect to (that) a corresponding side (has) to a corresponding side. [(Which is) the very thing it was required to show].

Πόρισμα.


Ἦσαύτως δὲ καὶ ἐπὶ τῶν [ὁμοίων] τετραπλεύρων δειχθήσεται, ὅτι ἐν διπλασίονι λόγῳ εἰσὶ τῶν ὁμολόγων πλευρῶν. ἐδείχθη δὲ καὶ ἐπὶ τῶν τριγώνων· ὥστε καὶ καθόλου τὰ ὅμοια εὐθύγραμμα σχήματα πρὸς ἄλληλα ἐν διπλασίονι λόγῳ εἰσὶ τῶν ὁμολόγων πλευρῶν. ὅπερ ἔδει δεῖξαι.

Corollary

And, in the same manner, it can also be shown for [similar] quadrilaterals that they are in the squared ratio of (their) corresponding sides. And it was also shown for triangles. Hence, in general, similar rectilinear figures are also to one another in the squared ratio of (their) corresponding sides. (Which is) the very thing it was required to show.

κα'.

Τὰ τῶν αὐτῶν εὐθύγραμμω ὅμοια καὶ ἀλλήλοις ἐστὶν ὅμοια.




Ἐστω γὰρ ἑκάτερον τῶν A , B εὐθύγραμμων τῶν Γ ὁμοιον· λέγω, ὅτι καὶ τὸ A τῶν B ἐστὶν ὅμοιον.

Ἐπεὶ γὰρ ὁμοιον ἐστὶ τὸ A τῶν Γ , ἰσογώνιον τέ ἐστὶν αὐτῶν καὶ τὰς περὶ τὰς ἴσας γωνίας πλευρὰς ἀνάλογον ἔχει. πάλιν, ἐπεὶ ὁμοιον ἐστὶ τὸ B τῶν Γ , ἰσογώνιον τέ ἐστὶν αὐτῶν καὶ τὰς περὶ τὰς ἴσας γωνίας πλευρὰς ἀνάλογον ἔχει. ἑκάτερον ἄρα τῶν A , B τῶν Γ ἰσογώνιον τέ ἐστὶ καὶ τὰς περὶ τὰς ἴσας γωνίας πλευρὰς ἀνάλογον ἔχει [ὥστε καὶ τὸ A τῶν B ἰσογώνιον τέ ἐστὶ καὶ τὰς περὶ τὰς ἴσας γωνίας

Proposition 21

(Rectilinear figures) similar to the same rectilinear figure are also similar to one another.



Let each of the rectilinear figures A and B be similar to (the rectilinear figure) C . I say that A is also similar to B .

For since A is similar to C , (A) is equiangular to (C), and has the sides about the equal angles proportional [Def. 6.1]. Again, since B is similar to C , (B) is equiangular to (C), and has the sides about the equal angles proportional [Def. 6.1]. Thus, A and B are each equiangular to C , and have the sides about the equal angles

πλευράς ἀνάλογον ἔχει]. ὁμοιον ἄρα ἐστὶ τὸ A τῷ B · ὅπερ ἔδει δεῖξαι.



proportional [hence, A is also equiangular to B , and has the sides about the equal angles proportional]. Thus, A is similar to B [Def. 6.1]. (Which is) the very thing it was required to show.

κβ'.

Proposition 22

Ἐὰν τέσσαρες εὐθεῖαι ἀνάλογον ᾧσιν, καὶ τὰ ἀπ' αὐτῶν εὐθύγραμμα ὁμοιά τε καὶ ὁμοίως ἀναγεγραμμένα ἀνάλογον ἔσται· καὶ τὰ ἀπ' αὐτῶν εὐθύγραμμα ὁμοιά τε καὶ ὁμοίως ἀναγεγραμμένα ἀνάλογον ἦ, καὶ αὐτὰ αἱ εὐθεῖαι ἀνάλογον ἔσονται.

If four straight-lines are proportional then similar, and similarly described, rectilinear figures (drawn) on them will also be proportional. And if similar, and similarly described, rectilinear figures (drawn) on them are proportional then the straight-lines themselves will also be proportional.



Ἐστωσαν τέσσαρες εὐθεῖαι ἀνάλογον αἱ AB , $\Gamma\Delta$, EZ , $H\Theta$, ὡς ἡ AB πρὸς τὴν $\Gamma\Delta$, οὕτως ἡ EZ πρὸς τὴν $H\Theta$, καὶ ἀναγεγράφωσαν ἀπὸ μὲν τῶν AB , $\Gamma\Delta$ ὁμοιά τε καὶ ὁμοίως κείμενα εὐθύγραμμα τὰ KAB , $\Lambda\Gamma\Delta$, ἀπὸ δὲ τῶν EZ , $H\Theta$ ὁμοιά τε καὶ ὁμοίως κείμενα εὐθύγραμμα τὰ MZ , $N\Theta$ · λέγω, ὅτι ἐστὶν ὡς τὸ KAB πρὸς τὸ $\Lambda\Gamma\Delta$, οὕτως τὸ MZ πρὸς τὸ $N\Theta$.

Let AB , CD , EF , and GH be four proportional straight-lines, (such that) as AB (is) to CD , so EF (is) to GH . And let the similar, and similarly laid out, rectilinear figures KAB and LCD have been described on AB and CD (respectively), and the similar, and similarly laid out, rectilinear figures MF and NH on EF and GH (respectively). I say that as KAB is to LCD , so MF (is) to NH .

Εἰλήφθω γὰρ τῶν μὲν AB , $\Gamma\Delta$ τρίτη ἀνάλογον ἡ Ξ , τῶν δὲ EZ , $H\Theta$ τρίτη ἀνάλογον ἡ O . καὶ ἐπεὶ ἐστὶν ὡς μὲν ἡ AB πρὸς τὴν $\Gamma\Delta$, οὕτως ἡ EZ πρὸς τὴν $H\Theta$, ὡς δὲ ἡ $\Gamma\Delta$ πρὸς τὴν Ξ , οὕτως ἡ $H\Theta$ πρὸς τὴν O , δι' ἴσου ἄρα ἐστὶν ὡς ἡ AB πρὸς τὴν Ξ , οὕτως ἡ EZ πρὸς τὴν O . ἀλλ' ὡς μὲν ἡ AB πρὸς τὴν Ξ , οὕτως [καὶ] τὸ KAB πρὸς τὸ $\Lambda\Gamma\Delta$, ὡς δὲ ἡ EZ πρὸς τὴν O , οὕτως τὸ MZ πρὸς τὸ $N\Theta$ · καὶ ὡς ἄρα τὸ KAB πρὸς τὸ $\Lambda\Gamma\Delta$, οὕτως τὸ MZ πρὸς τὸ $N\Theta$.

For let a third (straight-line) O have been taken (which is) proportional to AB and CD , and a third (straight-line) P proportional to EF and GH [Prop. 6.11]. And since as AB is to CD , so EF (is) to GH , and as CD (is) to O , so GH (is) to P , thus, via equality, as AB is to O , so EF (is) to P [Prop. 5.22]. But, as AB (is) to O , so [also] KAB (is) to LCD , and as EF (is) to P , so MF (is) to NH [Prop. 5.19 corr.]. And, thus, as KAB (is) to LCD , so MF (is) to NH .

Ἀλλὰ δὴ ἔστω ὡς τὸ KAB πρὸς τὸ $\Lambda\Gamma\Delta$, οὕτως τὸ MZ πρὸς τὸ $N\Theta$ · λέγω, ὅτι ἐστὶ καὶ ὡς ἡ AB πρὸς τὴν $\Gamma\Delta$, οὕτως ἡ EZ πρὸς τὴν $H\Theta$. εἰ γὰρ μὴ ἐστὶν, ὡς ἡ AB πρὸς τὴν $\Gamma\Delta$, οὕτως ἡ EZ πρὸς τὴν $H\Theta$, ἔστω ὡς ἡ AB πρὸς τὴν $\Gamma\Delta$, οὕτως ἡ EZ πρὸς τὴν $\Pi\Gamma$, καὶ ἀναγεγράφω ἀπὸ τῆς

And so let KAB be to LCD , as MF (is) to NH . I say also that as AB is to CD , so EF (is) to GH . For if as AB is to CD , so EF (is) not to GH , let AB be to CD , as EF

ΠΡ ὁποτέρῳ τῶν ΜΖ, ΝΘ ὁμοίον τε καὶ ὁμοίως κείμενον εὐθύγραμμον τὸ ΣΡ.

Ἐπεὶ οὖν ἐστὶν ὡς ἡ ΑΒ πρὸς τὴν ΓΔ, οὕτως ἡ ΕΖ πρὸς τὴν ΠΡ, καὶ ἀναγέγραπται ἀπὸ μὲν τῶν ΑΒ, ΓΔ ὁμοία τε καὶ ὁμοίως κείμενα τὰ ΚΑΒ, ΛΓΔ, ἀπὸ δὲ τῶν ΕΖ, ΠΡ ὁμοία τε καὶ ὁμοίως κείμενα τὰ ΜΖ, ΣΡ, ἔστιν ἄρα ὡς τὸ ΚΑΒ πρὸς τὸ ΛΓΔ, οὕτως τὸ ΜΖ πρὸς τὸ ΣΡ. ὑπόκειται δὲ καὶ ὡς τὸ ΚΑΒ πρὸς τὸ ΛΓΔ, οὕτως τὸ ΜΖ πρὸς τὸ ΝΘ· καὶ ὡς ἄρα τὸ ΜΖ πρὸς τὸ ΣΡ, οὕτως τὸ ΜΖ πρὸς τὸ ΝΘ. τὸ ΜΖ ἄρα πρὸς ἐκάτερον τῶν ΝΘ, ΣΡ τὸν αὐτὸν ἔχει λόγον· ἴσον ἄρα ἐστὶ τὸ ΝΘ τῷ ΣΡ. ἔστι δὲ αὐτῶ καὶ ὁμοιον καὶ ὁμοίως κείμενον· ἴση ἄρα ἡ ΗΘ τῇ ΠΡ. καὶ ἐπεὶ ἐστὶν ὡς ἡ ΑΒ πρὸς τὴν ΓΔ, οὕτως ἡ ΕΖ πρὸς τὴν ΠΡ, ἴση δὲ ἡ ΠΡ τῇ ΗΘ, ἔστιν ἄρα ὡς ἡ ΑΒ πρὸς τὴν ΓΔ, οὕτως ἡ ΕΖ πρὸς τὴν ΗΘ.

Ἐάν ἄρα τέσσαρες εὐθεῖαι ἀνάλογον ὦσιν, καὶ τὰ ἀπ' αὐτῶν εὐθύγραμμα ὁμοία τε καὶ ὁμοίως ἀναγεγραμμένα ἀνάλογον ἔσται· καθ' ἃ τὰ ἀπ' αὐτῶν εὐθύγραμμα ὁμοία τε καὶ ὁμοίως ἀναγεγραμμένα ἀνάλογον ἦ, καὶ αὐτὰ αἱ εὐθεῖαι ἀνάλογον ἔσονται· ὅπερ ἔδει δεῖξαι.

(is) to QR [Prop. 6.12]. And let the rectilinear figure SR , similar, and similarly laid down, to either of MF or NH , have been described on QR [Props. 6.18, 6.21].

Therefore, since as AB is to CD , so EF (is) to QR , and the similar, and similarly laid out, (rectilinear figures) KAB and LCD have been described on AB and CD (respectively), and the similar, and similarly laid out, (rectilinear figures) MF and SR on EF and QR (respectively), thus as KAB is to LCD , so MF (is) to SR (see above). And it was also assumed that as KAB (is) to LCD , so MF (is) to NH . Thus, also, as MF (is) to SR , so MF (is) to NH [Prop. 5.11]. Thus, MF has the same ratio to each of NH and SR . Thus, NH is equal to SR [Prop. 5.9]. And it is also similar, and similarly laid out, to it. Thus, GH (is) equal to QR .[†] And since AB is to CD , as EF (is) to QR , and QR (is) equal to GH , thus as AB is to CD , so EF (is) to GH .

Thus, if four straight-lines are proportional, then similar, and similarly described, rectilinear figures (drawn) on them will also be proportional. And if similar, and similarly described, rectilinear figures (drawn) on them are proportional then the straight-lines themselves will also be proportional. (Which is) the very thing it was required to show.

[†] Here, Euclid assumes, without proof, that if two similar figures are equal then any pair of corresponding sides is also equal.

κγ'.

Τὰ ἰσογώνια παραλληλόγραμμα πρὸς ἄλληλα λόγον ἔχει τὸν συγκείμενον ἐκ τῶν πλευρῶν.

Ἐστω ἰσογώνια παραλληλόγραμμα τὰ ΑΓ, ΓΖ ἴσην ἔχοντα τὴν ὑπὸ ΒΓΔ γωνίαν τῇ ὑπὸ ΕΓΗ· λέγω, ὅτι τὸ ΑΓ παραλληλόγραμμον πρὸς τὸ ΓΖ παραλληλόγραμμον λόγον ἔχει τὸν συγκείμενον ἐκ τῶν πλευρῶν.

Κείσθω γὰρ ὥστε ἐπ' εὐθείας εἶναι τὴν ΒΓ τῇ ΓΗ· ἐπ' εὐθείας ἄρα ἐστὶ καὶ ἡ ΔΓ τῇ ΓΕ. καὶ συμπληρώσθω τὸ ΔΗ παραλληλόγραμμον, καὶ ἐκκείσθω τις εὐθεῖα ἡ Κ, καὶ γερονέτω ὡς μὲν ἡ ΒΓ πρὸς τὴν ΓΗ, οὕτως ἡ Κ πρὸς τὴν Λ, ὡς δὲ ἡ ΔΓ πρὸς τὴν ΓΕ, οὕτως ἡ Λ πρὸς τὴν Μ.

Οἱ ἄρα λόγοι τῆς τε Κ πρὸς τὴν Λ καὶ τῆς Λ πρὸς τὴν Μ οἱ αὐτοὶ εἰσι τοῖς λόγοις τῶν πλευρῶν, τῆς τε ΒΓ πρὸς τὴν ΓΗ καὶ τῆς ΔΓ πρὸς τὴν ΓΕ. ἀλλ' ὁ τῆς Κ πρὸς Μ λόγος σύγκειται ἐκ τε τοῦ τῆς Κ πρὸς Λ λόγου καὶ τοῦ τῆς Λ πρὸς Μ· ὥστε καὶ ἡ Κ πρὸς τὴν Μ λόγον ἔχει τὸν συγκείμενον ἐκ τῶν πλευρῶν. καὶ ἐπεὶ ἐστὶν ὡς ἡ ΒΓ πρὸς τὴν ΓΗ, οὕτως τὸ ΑΓ παραλληλόγραμμον πρὸς τὸ ΓΘ, ἀλλ' ὡς ἡ ΒΓ πρὸς τὴν ΓΗ, οὕτως ἡ Κ πρὸς τὴν Λ, καὶ ὡς ἄρα ἡ Κ πρὸς τὴν Λ, οὕτως τὸ ΑΓ πρὸς τὸ ΓΘ. πάλιν, ἐπεὶ ἐστὶν ὡς ἡ ΔΓ πρὸς τὴν ΓΕ, οὕτως τὸ ΓΘ παραλληλόγραμμον πρὸς τὸ ΓΖ, ἀλλ' ὡς ἡ ΔΓ πρὸς τὴν ΓΕ,

Proposition 23

Equiangular parallelograms have to one another the ratio compounded[†] out of (the ratios of) their sides.


Let AC and CF be equiangular parallelograms having angle BCD equal to ECG . I say that parallelogram AC has to parallelogram CF the ratio compounded out of (the ratios of) their sides.

For let BC be laid down so as to be straight-on to CG . Thus, DC is also straight-on to CE [Prop. 1.14]. And let the parallelogram DG have been completed. And let some straight-line K have been laid down. And let it be contrived that as BC (is) to CG , so K (is) to L , and as DC (is) to CE , so L (is) to M [Prop. 6.12].


Thus, the ratios of K to L and of L to M are the same as the ratios of the sides, (namely), BC to CG and DC to CE (respectively). But, the ratio of K to M is compounded out of the ratio of K to L and (the ratio) of L to M . Hence, K also has to M the ratio compounded out of (the ratios of) the sides (of the parallelograms). And since as BC is to CG , so parallelogram AC (is) to CH [Prop. 6.1], but as BC (is) to CG , so K (is) to L , thus, also, as K (is) to L , so (parallelogram) AC (is) to CH . Again, since as DC (is) to CE , so parallelogram

οὕτως ἡ Λ πρὸς τὴν M , καὶ ὡς ἄρα ἡ Λ πρὸς τὴν M , οὕτως τὸ $\Gamma\Theta$ παραλληλόγραμμον πρὸς τὸ ΓZ παραλληλόγραμμον. ἐπεὶ οὖν ἐδείχθη, ὡς μὲν ἡ K πρὸς τὴν Λ , οὕτως τὸ $ΑΓ$ παραλληλόγραμμον πρὸς τὸ $\Gamma\Theta$ παραλληλόγραμμον, ὡς δὲ ἡ Λ πρὸς τὴν M , οὕτως τὸ $\Gamma\Theta$ παραλληλόγραμμον πρὸς τὸ ΓZ παραλληλόγραμμον, δι' ἴσου ἄρα ἐστὶν ὡς ἡ K πρὸς τὴν M , οὕτως τὸ $ΑΓ$ πρὸς τὸ ΓZ παραλληλόγραμμον. ἡ δὲ K πρὸς τὴν M λόγον ἔχει τὸν συγκείμενον ἐκ τῶν πλευρῶν· καὶ τὸ $ΑΓ$ ἄρα πρὸς τὸ ΓZ λόγον ἔχει τὸν συγκείμενον ἐκ τῶν πλευρῶν.

CH (is) to CF [Prop. 6.1], but as DC (is) to CE , so L (is) to M , thus, also, as L (is) to M , so parallelogram CH (is) to parallelogram CF . Therefore, since it was shown that as K (is) to L , so parallelogram AC (is) to parallelogram CH , and as L (is) to M , so parallelogram CH (is) to parallelogram CF , thus, via equality, as K is to M , so (parallelogram) AC (is) to parallelogram CF [Prop. 5.22]. And K has to M the ratio compounded out of (the ratios of) the sides (of the parallelograms). Thus, (parallelogram) AC also has to (parallelogram) CF the ratio compounded out of (the ratio of) their sides.



Τὰ ἄρα ἰσογώνια παραλληλόγραμματα πρὸς ἀλλήλα λόγον ἔχει τὸν συγκείμενον ἐκ τῶν πλευρῶν· ὅπερ ἔδει δεῖξαι.



Thus, equiangular parallelograms have to one another the ratio compounded out of (the ratio of) their sides. (Which is) the very thing it was required to show.

† In modern terminology, if two ratios are “compounded” then they are multiplied together.

κδ'.

Proposition 24

Παντὸς παραλληλογράμμου τὰ περὶ τὴν διάμετρον παραλληλόγραμματα ὁμοία ἐστὶ τῷ τε ὅλῳ καὶ ἀλλήλοις.

Ἐστω παραλληλόγραμμον τὸ $ΑΒΓΔ$, διάμετρος δὲ αὐτοῦ ἡ $ΑΓ$, περὶ δὲ τὴν $ΑΓ$ παραλληλόγραμματα ἔστω τὰ $ΕΗ$, ΘK . λέγω, ὅτι ἐκάτερον τῶν $ΕΗ$, ΘK παραλληλογράμμων ὁμοιὸν ἐστὶ ὅλῳ τῷ $ΑΒΓΔ$ καὶ ἀλλήλοις.

Ἐπεὶ γὰρ τριγώνου τοῦ $ΑΒΓ$ παρὰ μίαν τῶν πλευρῶν τὴν $ΒΓ$ ἦρται ἡ $ΕΖ$, ἀνάλογόν ἐστὶν ὡς ἡ $ΒΕ$ πρὸς τὴν $ΕΑ$, οὕτως ἡ ΓZ πρὸς τὴν $ΖΑ$. πάλιν, ἐπεὶ τριγώνου τοῦ $ΑΓΔ$ παρὰ μίαν τὴν $\Gamma Δ$ ἦρται ἡ $ΖΗ$, ἀνάλογόν ἐστὶν ὡς ἡ ΓZ πρὸς τὴν $ΖΑ$, οὕτως ἡ ΔH πρὸς τὴν $ΗΑ$. ἀλλ' ὡς ἡ ΓZ πρὸς τὴν $ΖΑ$, οὕτως ἐδείχθη καὶ ἡ $ΒΕ$ πρὸς τὴν $ΕΑ$ · καὶ ὡς ἄρα ἡ $ΒΕ$ πρὸς τὴν $ΕΑ$, οὕτως ἡ ΔH πρὸς τὴν $ΗΑ$, καὶ συνθέντι ἄρα ὡς ἡ $ΒΑ$ πρὸς $ΑΕ$, οὕτως ἡ $\Delta Α$ πρὸς $ΑΗ$, καὶ ἐναλλάξ ὡς ἡ $ΒΑ$ πρὸς τὴν $ΑΔ$, οὕτως ἡ $ΕΑ$ πρὸς τὴν $ΑΗ$. τῶν ἄρα $ΑΒΓΔ$, $ΕΗ$ παραλληλογράμμων ἀνάλογόν εἰσιν αἱ πλευραὶ αἱ περὶ τὴν κοινὴν γωνίαν τὴν ὑπὸ $ΒΑΔ$. καὶ ἐπεὶ παράλληλός ἐστὶν ἡ $ΗΖ$ τῇ $\Delta Γ$, ἴση ἐστὶν ἡ μὲν ὑπὸ $ΑΖΗ$ γωνία τῇ ὑπὸ $\Delta ΓΑ$ · καὶ κοινὴ τῶν δύο


In any parallelogram the parallelograms about the diagonal are similar to the whole, and to one another.

Let $ABCD$ be a parallelogram, and AC its diagonal. And let EG and HK be parallelograms about AC . I say that the parallelograms EG and HK are each similar to the whole (parallelogram) $ABCD$, and to one another.

For since EF has been drawn parallel to one of the sides BC of triangle ABC , proportionally, as BE is to EA , so CF (is) to FA [Prop. 6.2]. Again, since FG has been drawn parallel to one (of the sides) CD of triangle ACD , proportionally, as CF is to FA , so DG (is) to GA [Prop. 6.2]. But, as CF (is) to FA , so it was also shown (is) BE to EA . And thus as BE (is) to EA , so DG (is) to GA . And, thus, compounding, as BA (is) to AE , so DA (is) to AG [Prop. 5.18]. And, alternately, as BA (is) to AD , so EA (is) to AG [Prop. 5.16]. Thus, in parallelograms $ABCD$ and EG the sides about the common angle BAD are proportional. And since GF is parallel to DC , angle AFG is equal to DCA [Prop. 1.29].

τριγώνων τῶν $\Delta\Gamma$, AHZ ἡ ὑπὸ $\Delta\Gamma$ γωνία· ἰσογώνιον ἄρα ἐστὶ τὸ $\Delta\Gamma$ τρίγωνον τῷ AHZ τριγώνῳ. διὰ τὰ αὐτὰ δὴ καὶ τὸ $ΑΓΒ$ τρίγωνον ἰσογώνιον ἐστὶ τῷ AZE τριγώνῳ, καὶ ὅλον τὸ $ΑΒΓΔ$ παραλληλόγραμμον τῷ EH παραλληλογράμμῳ ἰσογώνιον ἐστίν. ἀνάλογον ἄρα ἐστὶν ὡς ἡ AD πρὸς τὴν $\Delta\Gamma$, οὕτως ἡ AH πρὸς τὴν HZ , ὡς δὲ ἡ $\Delta\Gamma$ πρὸς τὴν ΓA , οὕτως ἡ HZ πρὸς τὴν ZA , ὡς δὲ ἡ $\Delta\Gamma$ πρὸς τὴν ΓB , οὕτως ἡ AZ πρὸς τὴν ZE , καὶ ἔτι ὡς ἡ ΓB πρὸς τὴν BA , οὕτως ἡ ZE πρὸς τὴν EA . καὶ ἐπεὶ ἐδείχθη ὡς μὲν ἡ $\Delta\Gamma$ πρὸς τὴν ΓA , οὕτως ἡ HZ πρὸς τὴν ZA , ὡς δὲ ἡ $\Delta\Gamma$ πρὸς τὴν ΓB , οὕτως ἡ AZ πρὸς τὴν ZE , δι' ἴσου ἄρα ἐστὶν ὡς ἡ $\Delta\Gamma$ πρὸς τὴν ΓB , οὕτως ἡ HZ πρὸς τὴν ZE . τῶν ἄρα $ΑΒΓΔ$, EH παραλληλογράμμων ἀνάλογόν εἰσιν αἱ πλευραὶ αἱ περὶ τὰς ἴσας γωνίας· ὁμοίον ἄρα ἐστὶ τὸ $ΑΒΓΔ$ παραλληλόγραμμον τῷ EH παραλληλογράμμῳ. διὰ τὰ αὐτὰ δὴ τὸ $ΑΒΓΔ$ παραλληλόγραμμον καὶ τῷ $K\Theta$ παραλληλογράμμῳ ὁμοίον ἐστίν· ἐκάτερον ἄρα τῶν EH , ΘK παραλληλογράμμων τῷ $ΑΒΓΔ$ [παραλληλογράμμῳ] ὁμοίον ἐστίν. τὰ δὲ τῶν αὐτῶν εὐθυγράμμῳ ὁμοία καὶ ἀλλήλοις ἐστὶν ὁμοία· καὶ τὸ EH ἄρα παραλληλόγραμμον τῷ ΘK παραλληλογράμμῳ ὁμοίον ἐστίν.


And angle DAC (is) common to the two triangles ADC and AGF . Thus, triangle ADC is equiangular to triangle AGF [Prop. 1.32]. So, for the same (reasons), triangle ACB is equiangular to triangle AFE , and the whole parallelogram $ABCD$ is equiangular to parallelogram EG . Thus, proportionally, as AD (is) to DC , so AG (is) to GF , and as DC (is) to CA , so GF (is) to FA , and as AC (is) to CB , so AF (is) to FE , and, further, as CB (is) to BA , so FE (is) to EA [Prop. 6.4]. And since it was shown that as DC is to CA , so GF (is) to FA , and as AC (is) to CB , so AF (is) to FE , thus, via equality, as DC is to CB , so GF (is) to FE [Prop. 5.22]. Thus, in parallelograms $ABCD$ and EG the sides about the equal angles are proportional. Thus, parallelogram $ABCD$ is similar to parallelogram EG [Def. 6.1]. So, for the same (reasons), parallelogram $ABCD$ is also similar to parallelogram KH . Thus, parallelograms EG and HK are each similar to [parallelogram] $ABCD$. And (rectilinear figures) similar to the same rectilinear figure are also similar to one another [Prop. 6.21]. Thus, parallelogram EG is also similar to parallelogram HK .



Παντὸς ἄρα παραλληλογράμμου τὰ περὶ τὴν διάμετρον παραλληλόγραμμά ὁμοία ἐστὶ τῷ τε ὅλῳ καὶ ἀλλήλοις· ὅπερ ἔδει δεῖξαι.

κε'.


Τῷ δοθέντι εὐθυγράμμῳ ὁμοίον καὶ ἄλλῳ τῷ δοθέντι ἴσον τὸ αὐτὸ συστήσασθαι.



Thus, in any parallelogram the parallelograms about the diagonal are similar to the whole, and to one another. (Which is) the very thing it was required to show.

Proposition 25

To construct a single (rectilinear figure) similar to a given rectilinear figure, and equal to a different given rectilinear figure.



Ἐστω τὸ μὲν δοθὲν εὐθύγραμμον, ᾧ δεῖ ὅμοιον συστήσασθαι, τὸ $ABΓ$, ᾧ δὲ δεῖ ἴσον, τὸ Δ . δεῖ δὴ τῶ μὲν $ABΓ$ ὅμοιον, τῶ δὲ Δ ἴσον τὸ αὐτὸ συστήσασθαι.

Παραβεβλήσθω γὰρ παρὰ μὲν τὴν $BΓ$ τῶ $ABΓ$ τριγώνω ἴσον παραλληλόγραμμον τὸ BE , παρὰ δὲ τὴν $ΓΕ$ τῶ Δ ἴσον παραλληλόγραμμον τὸ $ΓΜ$ ἐν γωνίᾳ τῇ ὑπὸ $ZΓΕ$, ἣ ἔστιν ἴση τῇ ὑπὸ $ΓΒΛ$. ἐπ' εὐθείας ἄρα ἔστιν ἡ μὲν $BΓ$ τῇ $ΓΖ$, ἣ δὲ $ΛΕ$ τῇ $ΕΜ$. καὶ εἰλήφθω τῶν $BΓ$, $ΓΖ$ μέση ἀνάλογον ἡ $HΘ$, καὶ ἀναγεγράφθω ἀπὸ τῆς $HΘ$ τῶ $ABΓ$ ὁμοίον τε καὶ ὁμοίως κείμενον τὸ $KHΘ$.


Καὶ ἐπεὶ ἔστιν ὡς ἡ $BΓ$ πρὸς τὴν $HΘ$, οὕτως ἡ $HΘ$ πρὸς τὴν $ΓΖ$, ἐὰν δὲ τρεῖς εὐθεῖαι ἀνάλογον ὦσιν, ἔστιν ὡς ἡ πρώτη πρὸς τὴν τρίτην, οὕτως τὸ ἀπὸ τῆς πρώτης εἶδος πρὸς τὸ ἀπὸ τῆς δευτέρας τὸ ὅμοιον καὶ ὁμοίως ἀναγεγόμενον, ἔστιν ἄρα ὡς ἡ $BΓ$ πρὸς τὴν $ΓΖ$, οὕτως τὸ $ABΓ$ τρίγωνον πρὸς τὸ $KHΘ$ τρίγωνον. ἀλλὰ καὶ ὡς ἡ $BΓ$ πρὸς τὴν $ΓΖ$, οὕτως τὸ BE παραλληλόγραμμον πρὸς τὸ EZ παραλληλόγραμμον. καὶ ὡς ἄρα τὸ $ABΓ$ τρίγωνον πρὸς τὸ $KHΘ$ τρίγωνον, οὕτως τὸ BE παραλληλόγραμμον πρὸς τὸ EZ παραλληλόγραμμον. ἐναλλάξ ἄρα ὡς τὸ $ABΓ$ τρίγωνον πρὸς τὸ BE παραλληλόγραμμον, οὕτως τὸ $KHΘ$ τρίγωνον πρὸς τὸ EZ παραλληλόγραμμον. ἴσον δὲ τὸ $ABΓ$ τρίγωνον τῶ BE παραλληλογράμμῳ· ἴσον ἄρα καὶ τὸ $KHΘ$ τρίγωνον τῶ EZ παραλληλογράμμῳ. ἀλλὰ τὸ EZ παραλληλόγραμμον τῶ Δ ἔστιν ἴσον· καὶ τὸ $KHΘ$ ἄρα τῶ Δ ἔστιν ἴσον. ἔστι δὲ τὸ $KHΘ$ καὶ τῶ $ABΓ$ ὅμοιον.

Τῶ ἄρα δοθέντι εὐθυγράμμῳ τῶ $ABΓ$ ὅμοιον καὶ ἄλλω τῶ δοθέντι τῶ Δ ἴσον τὸ αὐτὸ συνέσταται τὸ $KHΘ$ ὅπερ ἔδει ποιῆσαι.

κς'.

Ἐὰν ἀπὸ παραλληλογράμμου παραλληλόγραμμον ἀφαιρεθῇ ὁμοίον τε τῶ ὅλῳ καὶ ὁμοίως κείμενον κοινὴν γωνίαν ἔχον αὐτῶ, περὶ τὴν αὐτὴν διάμετρον ἔστι τῶ ὅλῳ.

Ἀπὸ γὰρ παραλληλογράμμου τοῦ $ABΓΔ$ παραλληλόγραμμον ἀφηρήσθω τὸ AZ ὅμοιον τῶ $ABΓΔ$ καὶ ὁμοίως κείμενον κοινὴν γωνίαν ἔχον αὐτῶ τὴν ὑπὸ ΔAB . λέγω,



Let ABC be the given rectilinear figure to which it is required to construct a similar (rectilinear figure), and D the (rectilinear figure) to which (the constructed figure) is required (to be) equal. So it is required to construct a single (rectilinear figure) similar to ABC , and equal to D .

For let the parallelogram BE , equal to triangle ABC , have been applied to (the straight-line) BC [Prop. 1.44], and the parallelogram CM , equal to D , (have been applied) to (the straight-line) CE , in the angle FCE , which is equal to CBL [Prop. 1.45]. Thus, BC is straight-on to CF , and LE to EM [Prop. 1.14]. And let the mean proportion GH have been taken of BC and CF [Prop. 6.13]. And let KGH , similar, and similarly laid out, to ABC have been described on GH [Prop. 6.18].

And since as BC is to GH , so GH (is) to CF , and if three straight-lines are proportional then as the first is to the third, so the figure (described) on the first (is) to the similar, and similarly described, (figure) on the second [Prop. 6.19 corr.], thus as BC is to CF , so triangle ABC (is) to triangle KGH . But, also, as BC (is) to CF , so parallelogram BE (is) to parallelogram EF [Prop. 6.1]. And, thus, as triangle ABC (is) to triangle KGH , so parallelogram BE (is) to parallelogram EF . Thus, alternately, as triangle ABC (is) to parallelogram BE , so triangle KGH (is) to parallelogram EF [Prop. 5.16]. And triangle ABC (is) equal to parallelogram BE . Thus, triangle KGH (is) also equal to parallelogram EF . But, parallelogram EF is equal to D . Thus, KGH is also equal to D . And KGH is also similar to ABC .


Thus, a single (rectilinear figure) KGH has been constructed (which is) similar to the given rectilinear figure ABC , and equal to a different given (rectilinear figure) D . (Which is) the very thing it was required to do.

Proposition 26

If from a parallelogram a(nother) parallelogram is subtracted (which is) similar, and similarly laid out, to the whole, having a common angle with it, then (the subtracted parallelogram) is about the same diagonal as the whole.

For, from parallelogram $ABCD$, let (parallelogram)

ὅτι περί τήν αὐτήν διάμετρον ἔστι τὸ ΑΒΓΔ τῷ ΑΖ.



Μή γάρ, ἀλλ' εἰ δυνατόν, ἔστω [αὐτῶν] διάμετρος ἡ ΑΘΓ, καὶ ἐκβληθεῖσα ἡ ΗΖ διήχθω ἐπὶ τὸ Θ, καὶ ἤχθω διὰ τοῦ Θ ὀπορέρα τῶν ΑΔ, ΒΓ παράλληλος ἡ ΘΚ.

Ἐπεὶ οὖν περί τήν αὐτήν διάμετρον ἔστι τὸ ΑΒΓΔ τῷ ΚΗ, ἔστιν ἄρα ὡς ἡ ΔΑ πρὸς τήν ΑΒ, οὕτως ἡ ΗΑ πρὸς τήν ΑΚ. ἔστι δὲ καὶ διὰ τήν ὁμοιότητα τῶν ΑΒΓΔ, ΕΗ καὶ ὡς ἡ ΔΑ πρὸς τήν ΑΒ, οὕτως ἡ ΗΑ πρὸς τήν ΑΕ· καὶ ὡς ἄρα ἡ ΗΑ πρὸς τήν ΑΚ, οὕτως ἡ ΗΑ πρὸς τήν ΑΕ. ἡ ΗΑ ἄρα πρὸς ἑκατέραν τῶν ΑΚ, ΑΕ τὸν αὐτὸν ἔχει λόγον. ἴση ἄρα ἔστιν ἡ ΑΕ τῇ ΑΚ ἢ ἐλάττων τῇ μείζονι· ὅπερ ἔστιν ἀδύνατον. οὐκ ἄρα οὐκ ἔστι περί τήν αὐτήν διάμετρον τὸ ΑΒΓΔ τῷ ΑΖ· περί τήν αὐτήν ἄρα ἔστι διάμετρον τὸ ΑΒΓΔ παραλληλόγραμμον τῷ ΑΖ παραλληλογράμμου.


Ἐὰν ἄρα ἀπὸ παραλληλογράμμου παραλληλόγραμμον ἀφαιρεθῇ ὁμοίον τε τῷ ὅλῳ καὶ ὁμοίως κείμενον κοινὴν γωνίαν ἔχον αὐτῷ, περί τήν αὐτήν διάμετρον ἔστι τῷ ὅλῳ· ὅπερ ἔδει δεῖξαι.

κζ'.

Πάντων τῶν παρὰ τήν αὐτήν εὐθεῖαν παραβαλλομένων παραλληλογράμμων καὶ ἐλλειπόντων εἶδει παραλληλογράμμοις ὁμοίοις τε καὶ ὁμοίως κείμενοις τῷ ἀπὸ τῆς ἡμισείας ἀναγραφόμενῳ μέγιστόν ἐστι τὸ ἀπὸ τῆς ἡμισείας παραβαλλόμενον [παραλληλόγραμμον] ὁμοίον ὃν τῷ ἐλλείμμαντι.

Ἐστω εὐθεῖα ἡ ΑΒ καὶ τετημήσθω δίχα κατὰ τὸ Γ, καὶ παραβεβλήσθω παρὰ τήν ΑΒ εὐθεῖαν τὸ ΑΔ παραλληλόγραμμον ἐλλείπον εἶδει παραλληλογράμμου τῷ ΔΒ ἀναγραφέντι ἀπὸ τῆς ἡμισείας τῆς ΑΒ, τουτέστι τῆς ΓΒ· λέγω, ὅτι πάντων τῶν παρὰ τήν ΑΒ παραβαλλομένων παραλληλογράμμων καὶ ἐλλειπόντων εἶδει [παραλληλογράμμοις] ὁμοίοις τε καὶ ὁμοίως κείμενοις τῷ ΔΒ μέγιστόν ἐστι τὸ

AF have been subtracted (which is) similar, and similarly laid out, to *ABCD*, having the common angle *DAB* with it. I say that *ABCD* is about the same diagonal as *AF*.



For (if) not, then, if possible, let *AHC* be [*ABCD*'s] diagonal. And producing *GF*, let it have been drawn through to (point) *H*. And let *HK* have been drawn through (point) *H*, parallel to either of *AD* or *BC* [Prop. 1.31].

Therefore, since *ABCD* is about the same diagonal as *KG*, thus as *DA* is to *AB*, so *GA* (is) to *AK* [Prop. 6.24]. And, on account of the similarity of *ABCD* and *EG*, also, as *DA* (is) to *AB*, so *GA* (is) to *AE*. Thus, also, as *GA* (is) to *AK*, so *GA* (is) to *AE*. Thus, *GA* has the same ratio to each of *AK* and *AE*. Thus, *AE* is equal to *AK* [Prop. 5.9], the lesser to the greater. The very thing is impossible. Thus, *ABCD* is not about the same diagonal as *AF*. Thus, parallelogram *ABCD* is about the same diagonal as parallelogram *AF*.


Thus, if from a parallelogram a (nother) parallelogram is subtracted (which is) similar, and similarly laid out, to the whole, having a common angle with it, then (the subtracted parallelogram) is about the same diagonal as the whole. (Which is) the very thing it was required to show.

Proposition 27

Of all the parallelograms applied to the same straight-line, and falling short by parallelogrammic figures similar, and similarly laid out, to the (parallelogram) described on half (the straight-line), the greatest is the [parallelogram] applied to half (the straight-line) which (is) similar to (that parallelogram) by which it falls short.

Let *AB* be a straight-line, and let it have been cut in half at (point) *C* [Prop. 1.10]. And let the parallelogram *AD* have been applied to the straight-line *AB*, falling short by the parallelogrammic figure *DB* (which is) applied to half of *AB*—that is to say, *CB*. I say that of all the parallelograms applied to *AB*, and falling short by

ΑΔ. παραβεβλήσθω γὰρ παρὰ τὴν ΑΒ εὐθεῖαν τὸ ΑΖ παρλληλόγραμμον ἑλλείπον εἶδει παραλληλογράμμῳ τῷ ΖΒ ὁμοίῳ τε καὶ ὁμοίως κειμένῳ τῷ ΔΒ· λέγω, ὅτι μείζον ἐστὶ τὸ ΑΔ τοῦ ΑΖ.



Ἐπεὶ γὰρ ὁμοίον ἐστὶ τὸ ΔΒ παραλληλόγραμμον τῷ ΖΒ παραλληλογράμμῳ, περὶ τὴν αὐτὴν εἰσι διάμετρον. ἤχθω αὐτῶν διάμετρος ἡ ΔΒ, καὶ καταγεγράφθω τὸ σχῆμα.

Ἐπεὶ οὖν ἴσον ἐστὶ τὸ ΓΖ τῷ ΖΕ, κοινὸν δὲ τὸ ΖΒ, ὅλον ἄρα τὸ ΓΘ ὅλῳ τῷ ΚΕ ἐστὶν ἴσον. ἀλλὰ τὸ ΓΘ τῷ ΓΗ ἐστὶν ἴσον, ἐπεὶ καὶ ἡ ΑΓ τῇ ΓΒ. καὶ τὸ ΗΓ ἄρα τῷ ΕΚ ἐστὶν ἴσον. κοινὸν προσκείσθω τὸ ΓΖ· ὅλον ἄρα τὸ ΑΖ τῷ ΑΜΝ γνώμονί ἐστιν ἴσον· ὥστε τὸ ΔΒ παραλληλόγραμμον, τουτέστι τὸ ΑΔ, τοῦ ΑΖ παραλληλογράμμου μείζον ἐστὶν.


Πάντων ἄρα τῶν παρὰ τὴν αὐτὴν εὐθεῖαν παραβαλλομένων παραλληλογράμμων καὶ ἑλλειπόντων εἶδει παραλληλογράμμοις ὁμοίοις τε καὶ ὁμοίως κειμένοις τῷ ἀπὸ τῆς ἡμισείας ἀναγραφομένῳ μέγιστόν ἐστὶ τὸ ἀπὸ τῆς ἡμισείας παραβληθέν· ὅπερ εἶδει δεῖξαι.

κη'.

Παρὰ τὴν δοθεῖσαν εὐθεῖαν τῷ δοθέντι εὐθυγράμμῳ ἴσον παραλληλόγραμμον παραβαλεῖν ἑλλείπον εἶδει παραλληλογράμμῳ ὁμοίῳ τῷ δοθέντι· δεῖ δὲ τὸ διδόμενον εὐθύγραμμον [ᾧ δεῖ ἴσον παραβαλεῖν] μὴ μείζον εἶναι τοῦ ἀπὸ τῆς ἡμισείας ἀναγραφομένου ὁμοίου τῷ ἑλλείμματι [τοῦ τε ἀπὸ τῆς ἡμισείας καὶ ᾧ δεῖ ὅμοιον ἑλλείπειν].

Ἐστω ἡ μὲν δοθεῖσα εὐθεῖα ἡ ΑΒ, τὸ δὲ δοθὲν εὐθύγραμμον, ᾧ δεῖ ἴσον παρὰ τὴν ΑΒ παραβαλεῖν, τὸ Γ μὴ μείζον [δὲ] τοῦ ἀπὸ τῆς ἡμισείας τῆς ΑΒ ἀναγραφομένου ὁμοίου τῷ ἑλλείμματι, ᾧ δὲ δεῖ ὅμοιον ἑλλείπειν, τὸ Δ· δεῖ δὴ

[parallelogrammic] figures similar, and similarly laid out, to DB , the greatest is AD . For let the parallelogram AF have been applied to the straight-line AB , falling short by the parallelogrammic figure FB (which is) similar, and similarly laid out, to DB . I say that AD is greater than AF .



For since parallelogram DB is similar to parallelogram FB , they are about the same diagonal [Prop. 6.26]. Let their (common) diagonal DB have been drawn, and let the (rest of the) figure have been described.

Therefore, since (complement) CF is equal to (complement) FE [Prop. 1.43], and (parallelogram) FB is common, the whole (parallelogram) CH is thus equal to the whole (parallelogram) KE . But, (parallelogram) CH is equal to CG , since AC (is) also (equal) to CB [Prop. 6.1]. Thus, (parallelogram) GC is also equal to EK . Let (parallelogram) CF have been added to both. Thus, the whole (parallelogram) AF is equal to the gnomon LMN . Hence, parallelogram DB —that is to say, AD —is greater than parallelogram AF .


Thus, for all parallelograms applied to the same straight-line, and falling short by a parallelogrammic figure similar, and similarly laid out, to the (parallelogram) described on half (the straight-line), the greatest is the [parallelogram] applied to half (the straight-line). (Which is) the very thing it was required to show.

Proposition 28[†]

To apply a parallelogram, equal to a given rectilinear figure, to a given straight-line, (the applied parallelogram) falling short by a parallelogrammic figure similar to a given (parallelogram). It is necessary for the given rectilinear figure [to which it is required to apply an equal (parallelogram)] not to be greater than the (parallelogram) described on half (of the straight-line) and similar to the deficit.

Let AB be the given straight-line, and C the given rectilinear figure to which the (parallelogram) applied to

παρὰ τὴν δοθεῖσαν εὐθεῖαν τὴν AB τῷ δοθέντι εὐθυγράμμῳ τῷ Γ ἴσον παραλληλόγραμμον παραβαλεῖν ἔλλειπον εἶδει παραλληλογράμμῳ ὁμοίῳ ὄντι τῷ Δ .




Τετμήσθω ἡ AB δίχα κατὰ τὸ E σημεῖον, καὶ ἀναγεγράφθω ἀπὸ τῆς EB τῷ Δ ὁμοιον καὶ ὁμοίως κείμενον τὸ $EBZH$, καὶ συμπληρώσθω τὸ AH παραλληλόγραμμον.

Εἰ μὲν οὖν ἴσον ἐστὶ τὸ AH τῷ Γ , γεγονόςς ἂν εἴη τὸ ἐπιταχθέν· παραβέβληται γὰρ παρὰ τὴν δοθεῖσαν εὐθεῖαν τὴν AB τῷ δοθέντι εὐθυγράμμῳ τῷ Γ ἴσον παραλληλόγραμμον τὸ AH ἔλλειπον εἶδει παραλληλογράμμῳ τῷ HB ὁμοίῳ ὄντι τῷ Δ . εἰ δὲ οὐ, μείζον ἔστω τὸ ΘE τοῦ Γ . ἴσον δὲ τὸ ΘE τῷ HB · μείζον ἄρα καὶ τὸ HB τοῦ Γ . ὅ δὲ μείζον ἐστὶ τὸ HB τοῦ Γ , ταύτῃ τῇ ὑπεροχῇ ἴσον, τῷ δὲ Δ ὁμοιον καὶ ὁμοίως κείμενον τὸ αὐτὸ συνεστάτω τὸ $KLMN$. ἀλλὰ τὸ Δ τῷ HB [ἐστὶν] ὁμοιον· καὶ τὸ KM ἄρα τῷ HB ἐστὶν ὁμοιον. ἔστω οὖν ὁμόλογος ἡ μὲν KA τῇ HE , ἡ δὲ AM τῇ HZ . καὶ ἐπεὶ ἴσον ἐστὶ τὸ HB τοῖς Γ, KM , μείζον ἄρα ἐστὶ τὸ HB τοῦ KM · μείζων ἄρα ἐστὶ καὶ ἡ μὲν HE τῆς KA , ἡ δὲ HZ τῆς AM . κείσθω τῇ μὲν KA ἴση ἡ $HΞ$, τῇ δὲ AM ἴση ἡ HO , καὶ συμπληρώσθω τὸ $\Xi HO \Pi$ παραλληλόγραμμον· ἴσον ἄρα καὶ ὁμοιον ἐστὶ [τὸ $H\Pi$] τῷ KM [ἀλλὰ τὸ KM τῷ HB ὁμοιον ἐστὶν]. καὶ τὸ $H\Pi$ ἄρα τῷ HB ὁμοιον ἐστὶν· περὶ τὴν αὐτὴν ἄρα διάμετρον ἐστὶ τὸ $H\Pi$ τῷ HB . ἔστω αὐτῶν διάμετρος ἡ $H\Pi B$, καὶ καταγεγράφθω τὸ σχῆμα.

Ἐπεὶ οὖν ἴσον ἐστὶ τὸ BH τοῖς Γ, KM , ὧν τὸ $H\Pi$ τῷ KM ἐστὶν ἴσον, λοιπὸς ἄρα ὁ $\Upsilon X \Phi$ γνῶμων λοιπῷ τῷ Γ ἴσος ἐστίν. καὶ ἐπεὶ ἴσον ἐστὶ τὸ OP τῷ $\Xi\Sigma$, κοινὸν προσκείσθω τὸ ΠB · ὅλον ἄρα τὸ OB ὅλῳ τῷ ΞB ἴσον ἐστίν. ἀλλὰ τὸ ΞB τῷ TE ἐστὶν ἴσον, ἐπεὶ καὶ πλευρὰ ἡ AE πλευρᾶ τῇ EB ἐστὶν ἴση· καὶ τὸ TE ἄρα τῷ OB ἐστὶν ἴσον. κοινὸν προσκείσθω τὸ $\Xi\Sigma$ · ὅλον ἄρα τὸ $T\Sigma$ ὅλῳ τῷ $\Phi X \Upsilon$ γνῶμονι ἐστὶν ἴσον. ἀλλ' ὁ $\Phi X \Upsilon$ γνῶμων τῷ Γ ἔδειχθη ἴσος· καὶ τὸ $T\Sigma$ ἄρα τῷ Γ ἐστὶν ἴσον.

Παρὰ τὴν δοθεῖσαν ἄρα εὐθεῖαν τὴν AB τῷ δοθέντι εὐθυγράμμῳ τῷ Γ ἴσον παραλληλόγραμμον παραβέβληται τὸ ΣT ἔλλειπον εἶδει παραλληλογράμμῳ τῷ ΠB ὁμοίῳ ὄντι

AB is required (to be) equal, [being] not greater than the (parallelogram) described on half of AB and similar to the deficit, and D the (parallelogram) to which the deficit is required (to be) similar. So it is required to apply a parallelogram, equal to the given rectilinear figure C , to the straight-line AB , falling short by a parallelogrammic figure which is similar to D .



Let AB have been cut in half at point E [Prop. 1.10], and let (parallelogram) $EBFG$, (which is) similar, and similarly laid out, to (parallelogram) D , have been described on EB [Prop. 6.18]. And let parallelogram AG have been completed.

Therefore, if AG is equal to C then the thing prescribed has happened. For a parallelogram AG , equal to the given rectilinear figure C , has been applied to the given straight-line AB , falling short by a parallelogrammic figure GB which is similar to D . And if not, let HE be greater than C . And HE (is) equal to GB [Prop. 6.1]. Thus, GB (is) also greater than C . So, let (parallelogram) $KLMN$ have been constructed (so as to be) both similar, and similarly laid out, to D , and equal to the excess by which GB is greater than C [Prop. 6.25]. But, GB [is] similar to D . Thus, KM is also similar to GB [Prop. 6.21]. Therefore, let KL correspond to GE , and LM to GF . And since (parallelogram) GB is equal to (figure) C and (parallelogram) KM , GB is thus greater than KM . Thus, GE is also greater than KL , and GF than LM . Let GO be made equal to KL , and GP to LM [Prop. 1.3]. And let the parallelogram $OGPQ$ have been completed. Thus, [GQ] is equal and similar to KM [but, KM is similar to GB]. Thus, GQ is also similar to GB [Prop. 6.21]. Thus, GQ and GB are about the same diagonal [Prop. 6.26]. Let GQB be their (common) diagonal, and let the (remainder of the) figure have been described.

Therefore, since BG is equal to C and KM , of which GQ is equal to KM , the remaining gnomon UWV is thus equal to the remainder C . And since (the complement) PR is equal to (the complement) OS [Prop. 1.43], let (parallelogram) QB have been added to both. Thus, the whole (parallelogram) PB is equal to the whole (par-

τῷ Δ [ἐπειδὴ περ τὸ ΠΒ τῷ ΗΠ ὁμοίον ἐστίν]· ὅπερ ἔδει ποιῆσαι.

allelogram) *OB*. But, *OB* is equal to *TE*, since side *AE* is equal to side *EB* [Prop. 6.1]. Thus, *TE* is also equal to *PB*. Let (parallelogram) *OS* have been added to both. Thus, the whole (parallelogram) *TS* is equal to the gnomon *VWU*. But, gnomon *VWU* was shown (to be) equal to *C*. Therefore, (parallelogram) *TS* is also equal to (figure) *C*.

Thus, the parallelogram *ST*, equal to the given rectilinear figure *C*, has been applied to the given straight-line *AB*, falling short by the parallelogrammic figure *QB*, which is similar to *D* [inasmuch as *QB* is similar to *GQ* [Prop. 6.24]]. (Which is) the very thing it was required to do.



† This proposition is a geometric solution of the quadratic equation $x^2 - \alpha x + \beta = 0$. Here, x is the ratio of a side of the deficit to the corresponding side of figure *D*, α is the ratio of the length of *AB* to the length of that side of figure *D* which corresponds to the side of the deficit running along *AB*, and β is the ratio of the areas of figures *C* and *D*. The constraint corresponds to the condition $\beta < \alpha^2/4$ for the equation to have real roots. Only the smaller root of the equation is found. The larger root can be found by a similar method.

κθ'.

Proposition 29†

Παρά τὴν δοθεῖσαν εὐθεῖαν τῷ δοθέντι εὐθύγραμμῳ ἴσον παραλληλόγραμμον παραβαλεῖν ὑπερβάλλον εἶδει παραλληλογράμμῳ ὁμοίῳ τῷ δοθέντι.

To apply a parallelogram, equal to a given rectilinear figure, to a given straight-line, (the applied parallelogram) overshooting by a parallelogrammic figure similar to a given (parallelogram).



Ἐστω ἡ μὲν δοθεῖσα εὐθεῖα ἡ *AB*, τὸ δὲ δοθὲν εὐθύγραμμον, ᾧ δεῖ ἴσον παρὰ τὴν *AB* παραβαλεῖν, τὸ *Γ*, ᾧ δὲ δεῖ ὁμοῖον ὑπερβάλλειν, τὸ *Δ*. δεῖ δὴ παρὰ τὴν *AB* εὐθεῖαν τῷ *Γ* εὐθύγραμμῳ ἴσον παραλληλόγραμμον παραβαλεῖν ὑπερβάλλον εἶδει παραλληλογράμμῳ ὁμοίῳ τῷ *Δ*.

Let *AB* be the given straight-line, and *C* the given rectilinear figure to which the (parallelogram) applied to *AB* is required (to be) equal, and *D* the (parallelogram) to which the excess is required (to be) similar. So it is required to apply a parallelogram, equal to the given rectilinear figure *C*, to the given straight-line *AB*, overshooting by a parallelogrammic figure similar to *D*.

Τετμήσθω ἡ *AB* δίχα κατὰ τὸ *E*, καὶ ἀναγεγράθω ἀπὸ τῆς *EB* τῷ *Δ* ὁμοῖον καὶ ὁμοίως κείμενον παραλληλόγραμμον τὸ *BZ*, καὶ συναμφοτέροις μὲν τοῖς *BZ*, *Γ* ἴσον, τῷ δὲ *Δ* ὁμοῖον καὶ ὁμοίως κείμενον τὸ αὐτὸ συνεστάτω τὸ *HΘ*. ὁμόλογος δὲ ἔστω ἡ μὲν *KΘ* τῇ *ΖΛ*, ἡ δὲ *KH* τῇ *ΖΕ*. καὶ ἐπεὶ μείζον ἐστὶ τὸ *HΘ* τοῦ *ZB*, μείζων ἄρα ἐστὶ καὶ ἡ μὲν *KΘ* τῆς *ΖΛ*, ἡ δὲ *KH* τῇ *ΖΕ*. ἐκβεβλήσθωσαν αἱ *ΖΛ*, *ΖΕ*, καὶ τῇ μὲν *KΘ* ἴση ἔστω ἡ *ZAM*, τῇ δὲ *KH* ἴση ἡ *ZEN*, καὶ συμπεπληρώσθω τὸ *MN*. τὸ *MN* ἄρα τῷ *HΘ* ἴσον τέ ἐστὶ καὶ ὁμοῖον. ἀλλὰ τὸ *HΘ* τῷ *ΕΛ* ἐστὶν ὁμοῖον.

Let *AB* have been cut in half at (point) *E* [Prop. 1.10], and let the parallelogram *BF*, (which is) similar, and similarly laid out, to *D*, have been described on *EB* [Prop. 6.18]. And let (parallelogram) *GH* have been constructed (so as to be) both similar, and similarly laid out, to *D*, and equal to the sum of *BF* and *C* [Prop. 6.25]. And let *KH* correspond to *FL*, and *KG* to *FE*. And since (parallelogram) *GH* is greater than (parallelogram) *FB*,

καὶ τὸ MN ἄρα τῷ EL ὁμοίον ἐστίν· περὶ τὴν αὐτὴν ἄρα διάμετρον ἐστὶ τὸ EL τῷ MN. ἤχθω αὐτῶν διάμετρος ἡ ZE, καὶ καταγεγράφθω τὸ σχῆμα.

Ἐπεὶ ἴσον ἐστὶ τὸ HΘ τοῖς EL, Γ, ἀλλὰ τὸ HΘ τῷ MN ἴσον ἐστίν, καὶ τὸ MN ἄρα τοῖς EL, Γ ἴσον ἐστίν. κοινὸν ἀφηρήσθω τὸ EL· λοιπὸς ἄρα ὁ ΨXΦ γνώμων τῷ Γ ἐστὶν ἴσος. καὶ ἐπεὶ ἴση ἐστὶν ἡ AE τῇ EB, ἴσον ἐστὶ καὶ τὸ AN τῷ NB, τοῦτέστι τῷ ΛO. κοινὸν προσκείσθω τὸ EΞ· ὅλον ἄρα τὸ AΞ ἴσον ἐστὶ τῷ ΦXΨ γνώμονι. ἀλλὰ ὁ ΦXΨ γνώμων τῷ Γ ἴσος ἐστίν· καὶ τὸ AΞ ἄρα τῷ Γ ἴσον ἐστίν.

Παρὰ τὴν δοθεῖσαν ἄρα εὐθεῖαν τὴν AB τῷ δοθέντι εὐθυγράμμῳ τῷ Γ ἴσον παραλληλόγραμμον παραβέβληται τὸ AΞ ὑπερβάλλον εἶδει παραλληλογράμμῳ τῷ ΠO ὁμοίῳ ὄντι τῷ Δ, ἐπεὶ καὶ τῷ EL ἐστὶν ὁμοίον τὸ OΠ· ὅπερ ἔδει ποιῆσαι.

KH is thus also greater than *FL*, and *KG* than *FE*. Let *FL* and *FE* have been produced, and let *FLM* be (made) equal to *KH*, and *FEN* to *KG* [Prop. 1.3]. And let (parallelogram) *MN* have been completed. Thus, *MN* is equal and similar to *GH*. But, *GH* is similar to *EL*. Thus, *MN* is also similar to *EL* [Prop. 6.21]. *EL* is thus about the same diagonal as *MN* [Prop. 6.26]. Let their (common) diagonal *FO* have been drawn, and let the (remainder of the) figure have been described.


And since (parallelogram) *GH* is equal to (parallelogram) *EL* and (figure) *C*, but *GH* is equal to (parallelogram) *MN*, *MN* is thus also equal to *EL* and *C*. Let *EL* have been subtracted from both. Thus, the remaining gnomon *XWV* is equal to (figure) *C*. And since *AE* is equal to *EB*, (parallelogram) *AN* is also equal to (parallelogram) *NB* [Prop. 6.1], that is to say, (parallelogram) *LP* [Prop. 1.43]. Let (parallelogram) *EO* have been added to both. Thus, the whole (parallelogram) *AO* is equal to the gnomon *VWX*. But, the gnomon *VWX* is equal to (figure) *C*. Thus, (parallelogram) *AO* is also equal to (figure) *C*.

Thus, the parallelogram *AO*, equal to the given rectilinear figure *C*, has been applied to the given straight-line *AB*, overshooting by the parallelogrammic figure *QP* which is similar to *D*, since *PQ* is also similar to *EL* [Prop. 6.24]. (Which is) the very thing it was required to do.

† This proposition is a geometric solution of the quadratic equation $x^2 + \alpha x - \beta = 0$. Here, x is the ratio of a side of the excess to the corresponding side of figure *D*, α is the ratio of the length of *AB* to the length of that side of figure *D* which corresponds to the side of the excess running along *AB*, and β is the ratio of the areas of figures *C* and *D*. Only the positive root of the equation is found.

λ'.


Τὴν δοθεῖσαν εὐθεῖαν πεπερασμένην ἄκρον καὶ μέσον λόγον τεμεῖν.



Ἐστω ἡ δοθεῖσα εὐθεῖα πεπερασμένη ἡ AB· δεῖ δὴ τὴν AB εὐθεῖαν ἄκρον καὶ μέσον λόγον τεμεῖν.

Proposition 30†

To cut a given finite straight-line in extreme and mean ratio.



Let *AB* be the given finite straight-line. So it is required to cut the straight-line *AB* in extreme and mean

Ἀναγεγράφθω ἀπὸ τῆς AB τετράγωνον τὸ $BΓ$, καὶ παραβεβλήσθω παρὰ τὴν $ΑΓ$ τῷ $BΓ$ ἴσον παραλληλόγραμμον τὸ $ΓΔ$ ὑπερβάλλον εἶδει τῷ $ΑΔ$ ὁμοίῳ τῷ $BΓ$.

Τετράγωνον δὲ ἐστὶ τὸ $BΓ$ · τετράγωνον ἄρα ἐστὶ καὶ τὸ $ΑΔ$. καὶ ἐπεὶ ἴσον ἐστὶ τὸ $BΓ$ τῷ $ΓΔ$, κοινὸν ἀφηρήσθω τὸ $ΓΕ$ · λοιπὸν ἄρα τὸ $BΖ$ λοιπῷ τῷ $ΑΔ$ ἐστὶν ἴσον. ἐστὶ δὲ αὐτῷ καὶ ἰσογώνιον· τῶν $BΖ$, $ΑΔ$ ἄρα ἀντιπεπόνθασιν αἱ πλευραὶ αἱ περὶ τὰς ἴσας γωνίας· ἐστὶν ἄρα ὡς ἡ $ΖΕ$ πρὸς τὴν $ΕΔ$, οὕτως ἡ $ΑΕ$ πρὸς τὴν $ΕΒ$. ἴση δὲ ἡ μὲν $ΖΕ$ τῇ $ΑΒ$, ἡ δὲ $ΕΔ$ τῇ $ΑΕ$. ἐστὶν ἄρα ὡς ἡ $ΒΑ$ πρὸς τὴν $ΑΕ$, οὕτως ἡ $ΑΕ$ πρὸς τὴν $ΕΒ$. μείζων δὲ ἡ $ΑΒ$ τῆς $ΑΕ$ · μείζων ἄρα καὶ ἡ $ΑΕ$ τῆς $ΕΒ$.

Ἡ ἄρα $ΑΒ$ εὐθεῖα ἄκρον καὶ μέσον λόγον τέτμηται κατὰ τὸ $Ε$, καὶ τὸ μείζον αὐτῆς τμημὰ ἐστὶ τὸ $ΑΕ$ · ὅπερ ἔδει ποιῆσαι.

ratio.

Let the square BC have been described on AB [Prop. 1.46], and let the parallelogram CD , equal to BC , have been applied to AC , overshooting by the figure AD (which is) similar to BC [Prop. 6.29].


And BC is a square. Thus, AD is also a square. And since BC is equal to CD , let (rectangle) CE have been subtracted from both. Thus, the remaining (rectangle) BF is equal to the remaining (square) AD . And it is also equiangular to it. Thus, the sides of BF and AD about the equal angles are reciprocally proportional [Prop. 6.14]. Thus, as FE is to ED , so AE (is) to EB . And FE (is) equal to AB , and ED to AE . Thus, as BA is to AE , so AE (is) to EB . And AB (is) greater than AE . Thus, AE (is) also greater than EB [Prop. 5.14].

Thus, the straight-line AB has been cut in extreme and mean ratio at E , and AE is its greater piece. (Which is) the very thing it was required to do.

† This method of cutting a straight-line is sometimes called the “Golden Section”—see Prop. 2.11.

λα'.

Ἐν τοῖς ὀρθογωνίοις τριγώνοις τὸ ἀπὸ τῆς τὴν ὀρθὴν γωνίαν ὑποτείνουσας πλευρᾶς εἶδος ἴσον ἐστὶ τοῖς ἀπὸ τῶν τὴν ὀρθὴν γωνίαν περιεχουσῶν πλευρῶν εἶδεσι τοῖς ὁμοίοις τε καὶ ὁμοίως ἀναγραφόμενοις.




Ἐστω τρίγωνον ὀρθογώνιον τὸ $ΑΒΓ$ ὀρθὴν ἔχον τὴν ὑπὸ $ΒΑΓ$ γωνίαν· λέγω, ὅτι τὸ ἀπὸ τῆς $BΓ$ εἶδος ἴσον ἐστὶ τοῖς ἀπὸ τῶν BA , $ΑΓ$ εἶδεσι τοῖς ὁμοίοις τε καὶ ὁμοίως ἀναγραφόμενοις.

Ἦχθω κάθετος ἡ $ΑΔ$.

Ἐπεὶ οὖν ἐν ὀρθογωνίῳ τριγώνῳ τῷ $ΑΒΓ$ ἀπὸ τῆς πρὸς τῷ $Α$ ὀρθῆς γωνίας ἐπὶ τὴν $BΓ$ βάσιν κάθετος ἤχεται ἡ $ΑΔ$, τὰ $ΑΒΔ$, $ΑΔΓ$ πρὸς τῇ καθετῷ τρίγωνα ὁμοία ἐστὶ τῷ τε ὅλῳ τῷ $ΑΒΓ$ καὶ ἀλλήλοις. καὶ ἐπεὶ ὁμοίον ἐστὶ τὸ $ΑΒΓ$ τῷ $ΑΒΔ$, ἐστὶν ἄρα ὡς ἡ $ΓΒ$ πρὸς τὴν BA , οὕτως ἡ $ΑΒ$ πρὸς τὴν $ΒΔ$. καὶ ἐπεὶ τρεῖς εὐθεῖαι ἀνάλογόν εἰσιν, ἐστὶν ὡς ἡ πρώτη πρὸς τὴν τρίτην, οὕτως τὸ ἀπὸ τῆς πρώτης εἶδος πρὸς

Proposition 31

In right-angled triangles, the figure (drawn) on the side subtending the right-angle is equal to the (sum of the) similar, and similarly described, figures on the sides surrounding the right-angle.



Let ABC be a right-angled triangle having the angle BAC a right-angle. I say that the figure (drawn) on BC is equal to the (sum of the) similar, and similarly described, figures on BA and AC .

Let the perpendicular AD have been drawn [Prop. 1.12].

Therefore, since, in the right-angled triangle ABC , the (straight-line) AD has been drawn from the right-angle at A perpendicular to the base BC , the triangles ABD and ADC about the perpendicular are similar to the whole (triangle) ABC , and to one another [Prop. 6.8]. And since ABC is similar to ABD , thus

τὸ ἀπὸ τῆς δευτέρας τὸ ὁμοιον καὶ ὁμοίως ἀναγραφόμενον. ὡς ἄρα ἡ ΓΒ πρὸς τὴν ΒΔ, οὕτως τὸ ἀπὸ τῆς ΓΒ εἶδος πρὸς τὸ ἀπὸ τῆς ΒΑ τὸ ὁμοιον καὶ ὁμοίως ἀναγραφόμενον. διὰ τὰ αὐτὰ δὴ καὶ ὡς ἡ ΒΓ πρὸς τὴν ΓΔ, οὕτως τὸ ἀπὸ τῆς ΒΓ εἶδος πρὸς τὸ ἀπὸ τῆς ΓΑ. ὥστε καὶ ὡς ἡ ΒΓ πρὸς τὰς ΒΔ, ΔΓ, οὕτως τὸ ἀπὸ τῆς ΒΓ εἶδος πρὸς τὰ ἀπὸ τῶν ΒΑ, ΑΓ τὰ ὁμοια καὶ ὁμοίως ἀναγραφόμενα. ἴση δὲ ἡ ΒΓ ταῖς ΒΔ, ΔΓ· ἴσον ἄρα καὶ τὸ ἀπὸ τῆς ΒΓ εἶδος τοῖς ἀπὸ τῶν ΒΑ, ΑΓ εἶδεσι τοῖς ὁμοίοις τε καὶ ὁμοίως ἀναγραφόμενοις.


Ἐν ἄρα τοῖς ὀρθογωνίοις τριγώνοις τὸ ἀπὸ τῆς τὴν ὀρθὴν γωνίαν ὑποτείνουσας πλευρᾶς εἶδος ἴσον ἐστὶ τοῖς ἀπὸ τῶν τὴν ὀρθὴν γωνίαν περιεχουσῶν πλευρῶν εἶδεσι τοῖς ὁμοίοις τε καὶ ὁμοίως ἀναγραφόμενοις· ὅπερ ἔδει δεῖξαι.

as CB is to BA , so AB (is) to BD [Def. 6.1]. And since three straight-lines are proportional, as the first is to the third, so the figure (drawn) on the first is to the similar, and similarly described, (figure) on the second [Prop. 6.19 corr.]. Thus, as CB (is) to BD , so the figure (drawn) on CB (is) to the similar, and similarly described, (figure) on BA . And so, for the same (reasons), as BC (is) to CD , so the figure (drawn) on BC (is) to the (figure) on CA . Hence, also, as BC (is) to BD and DC , so the figure (drawn) on BC (is) to the (sum of the) similar, and similarly described, (figures) on BA and AC [Prop. 5.24]. And BC is equal to BD and DC . Thus, the figure (drawn) on BC (is) also equal to the (sum of the) similar, and similarly described, figures on BA and AC [Prop. 5.9].

Thus, in right-angled triangles, the figure (drawn) on the side subtending the right-angle is equal to the (sum of the) similar, and similarly described, figures on the sides surrounding the right-angle. (Which is) the very thing it was required to show.

λβ'.

Ἐὰν δύο τρίγωνα συντεθῆ κατὰ μίαν γωνίαν τὰς δύο πλευρὰς ταῖς δυσὶ πλευραῖς ἀνάλογον ἔχοντα ὥστε τὰς ὁμολόγους αὐτῶν πλευρὰς καὶ παραλλήλους εἶναι, αἱ λοιπαὶ τῶν τριγώνων πλευραὶ ἐπ' εὐθείας ἔσσονται.




Ἐστω δύο τρίγωνα τὰ ABG , ΔGE τὰς δύο πλευρὰς τὰς BA , AG ταῖς δυσὶ πλευραῖς ταῖς ΔG , ΔE ἀνάλογον ἔχοντα, ὡς μὲν τὴν AB πρὸς τὴν AG , οὕτως τὴν ΔG πρὸς τὴν ΔE , παράλληλον δὲ τὴν μὲν AB τῇ ΔG , τὴν δὲ AG τῇ ΔE · λέγω, ὅτι ἐπ' εὐθείας ἐστὶν ἡ BG τῇ GE .

Ἐπεὶ γὰρ παράλληλός ἐστιν ἡ AB τῇ ΔG , καὶ εἰς αὐτὰς ἐμπέπτωκεν εὐθεῖα ἡ AG , αἱ ἐναλλάξ γωνίαὶ αἱ ὑπὸ BAG , $AG\Delta$ ἴσαι ἀλλήλαις εἰσίν. διὰ τὰ αὐτὰ δὴ καὶ ἡ ὑπὸ $G\Delta E$ τῇ ὑπὸ $AG\Delta$ ἴση ἐστίν. ὥστε καὶ ἡ ὑπὸ BAG τῇ ὑπὸ $G\Delta E$ ἐστὶν ἴση. καὶ ἐπεὶ δύο τρίγωνα ἐστὶ τὰ ABG , ΔGE μίαν γωνίαν τὴν πρὸς τῷ A μιᾶ γωνίᾳ τῇ πρὸς τῷ Δ ἴσην ἔχοντα, περὶ

Proposition 32

If two triangles, having two sides proportional to two sides, are placed together at a single angle such that the corresponding sides are also parallel, then the remaining sides of the triangles will be straight-on (with respect to one another).



Let ABC and DCE be two triangles having the two sides BA and AC proportional to the two sides DC and DE —so that as AB (is) to AC , so DC (is) to DE —and (having side) AB parallel to DC , and AC to DE . I say that (side) BC is straight-on to CE .


For since AB is parallel to DC , and the straight-line AC has fallen across them, the alternate angles BAC and ACD are equal to one another [Prop. 1.29]. So, for the same (reasons), CDE is also equal to ACD . And, hence, BAC is equal to CDE . And since ABC and DCE are two triangles having the one angle at A equal to the one

δὲ τὰς ἴσας γωνίας τὰς πλευρὰς ἀνάλογον, ὡς τὴν BA πρὸς τὴν AG , οὕτως τὴν GD πρὸς τὴν DE , ἰσογώνιον ἄρα ἐστὶ τὸ ABG τρίγωνον τῷ ΔGE τριγώνῳ· ἴση ἄρα ἡ ὑπὸ ABG γωνία τῇ ὑπὸ ΔGE . ἐδείχθη δὲ καὶ ἡ ὑπὸ AGD τῇ ὑπὸ BAG ἴση· ὅλη ἄρα ἡ ὑπὸ AGE δυσὶ ταῖς ὑπὸ ABG , BAG ἴση ἐστίν. κοινὴ προσκείσθω ἡ ὑπὸ AGB · αἱ ἄρα ὑπὸ AGE , AGB ταῖς ὑπὸ BAG , AGB , GBA ἴσαι εἰσίν. ἀλλ' αἱ ὑπὸ BAG , ABG , AGB δυσὶν ὀρθαῖς ἴσαι εἰσίν· καὶ αἱ ὑπὸ AGE , AGB ἄρα δυσὶν ὀρθαῖς ἴσαι εἰσίν. πρὸς δὲ τινὶ εὐθείᾳ τῇ AG καὶ τῷ πρὸς αὐτῇ σημείῳ τῷ Γ δύο εὐθεῖαι αἱ BG , GE μὴ ἐπὶ τὰ αὐτὰ μέρη κείμεναι τὰς ἐφεξῆς γωνίας τὰς ὑπὸ AGE , AGB δυσὶν ὀρθαῖς ἴσας ποιοῦσιν· ἐπ' εὐθείας ἄρα ἐστὶν ἡ BG τῇ GE .

Ἐὰν ἄρα δύο τρίγωνα συντεθῆ κατὰ μίαν γωνίαν τὰς δύο πλευρὰς ταῖς δυσὶ πλευραῖς ἀνάλογον ἔχοντα ὥστε τὰς ὁμολόγους αὐτῶν πλευρὰς καὶ παραλλήλους εἶναι, αἱ λοιπαὶ τῶν τριγώνων πλευραὶ ἐπ' εὐθείας ἔσσονται· ὅπερ ἔδει δεῖξαι.

λγ'.

Ἐν τοῖς ἴσοις κύκλοις αἱ γωνίαι τὸν αὐτὸν ἔχουσι λόγον ταῖς περιφερείαις, ἐφ' ὧν βεβήκασιν, ἐὰν τε πρὸς τοῖς κέντροις ἐὰν τε πρὸς ταῖς περιφερείαις ὡς βεβηκῆναι.



Ἐστωσαν ἴσοι κύκλοι οἱ ABG , ΔEZ , καὶ πρὸς μὲν τοῖς κέντροις αὐτῶν τοῖς H , Θ γωνίαι ἔστωσαν αἱ ὑπὸ BHG , $E\Theta Z$, πρὸς δὲ ταῖς περιφερείαις αἱ ὑπὸ BAG , $E\Delta Z$ · λέγω, ὅτι ἐστὶν ὡς ἡ BG περιφέρεια πρὸς τὴν EZ περιφέρειαν, οὕτως ἢ τε ὑπὸ BHG γωνία πρὸς τὴν ὑπὸ $E\Theta Z$ καὶ ἡ ὑπὸ BAG πρὸς τὴν ὑπὸ $E\Delta Z$.

Κείσθωσαν γὰρ τῇ μὲν BG περιφέρειᾳ ἴσαι κατὰ τὸ ἐξῆς ὁσαυδηποτοῦν αἱ GK , KL , τῇ δὲ EZ περιφέρειᾳ ἴσαι ὁσαυδηποτοῦν αἱ ZM , MN , καὶ ἐπεζεύχθωσαν αἱ HK , HL , ΘM , ΘN .


Ἐπεὶ οὖν ἴσαι εἰσίν αἱ BG , GK , KL περιφέρειαι ἀλλήλαις, ἴσαι εἰσὶ καὶ αἱ ὑπὸ BHG , GHK , KHL γωνίαι ἀλλήλαις· ὁσαπλασίων ἄρα ἐστὶν ἡ BA περιφέρεια τῆς BG , τοσαυταπλασίων ἐστὶ καὶ ἡ ὑπὸ BHL γωνία τῆς ὑπὸ BHG . διὰ τὰ

angle at D , and the sides about the equal angles proportional, (so that) as BA (is) to AC , so CD (is) to DE , triangle ABC is thus equiangular to triangle DCE [Prop. 6.6]. Thus, angle ABC is equal to DCE . And (angle) ACD was also shown (to be) equal to BAC . Thus, the whole (angle) ACE is equal to the two (angles) ABC and BAC . Let ACB have been added to both. Thus, ACE and ACB are equal to BAC , ACB , and CBA . But, BAC , ABC , and ACB are equal to two right-angles [Prop. 1.32]. Thus, ACE and ACB are also equal to two right-angles. Thus, the two straight-lines BC and CE , not lying on the same side, make adjacent angles ACE and ACB (whose sum is) equal to two right-angles with some straight-line AC , at the point C on it. Thus, BC is straight-on to CE [Prop. 1.14].

Thus, if two triangles, having two sides proportional to two sides, are placed together at a single angle such that the corresponding sides are also parallel, then the remaining sides of the triangles will be straight-on (with respect to one another). (Which is) the very thing it was required to show.

Proposition 33

In equal circles, angles have the same ratio as the (ratio of the) circumferences on which they stand, whether they are standing at the centers (of the circles) or at the circumferences.



Let ABC and DEF be equal circles, and let BGC and EHF be angles at their centers, G and H (respectively), and BAC and EDF (angles) at their circumferences. I say that as circumference BC is to circumference EF , so angle BGC (is) to EHF , and (angle) BAC to EDF .

For let any number whatsoever of consecutive (circumferences), CK and KL , be made equal to circumference BC , and any number whatsoever, FM and MN , to circumference EF . And let GK , GL , HM , and HN have been joined.

Therefore, since circumferences BC , CK , and KL are equal to one another, angles BGC , CGK , and KGL are also equal to one another [Prop. 3.27]. Thus, as many times as circumference BL is (divisible) by BC , so many

αὐτὰ δὴ καὶ ὁσαπλασίων ἐστὶν ἡ NE περιφέρεια τῆς EZ , τοσαυταπλασίων ἐστὶ καὶ ἡ ὑπὸ $N\Theta E$ γωνία τῆς ὑπὸ $E\Theta Z$. εἰ ἄρα ἴση ἐστὶν ἡ BA περιφέρεια τῆς EN περιφέρειᾶς, ἴση ἐστὶ καὶ γωνία ἡ ὑπὸ BHA τῆς ὑπὸ $E\Theta N$, καὶ εἰ μείζων ἐστὶν ἡ BA περιφέρεια τῆς EN περιφέρειᾶς, μείζων ἐστὶ καὶ ἡ ὑπὸ BHA γωνία τῆς ὑπὸ $E\Theta N$, καὶ εἰ ἐλάσσων, ἐλάσσων. τεσσάρων δὴ ὄντων μεγεθῶν, δύο μὲν περιφερειῶν τῶν $B\Gamma$, EZ , δύο δὲ γωνιῶν τῶν ὑπὸ BHG , $E\Theta Z$, εἴληπται τῆς μὲν $B\Gamma$ περιφέρειᾶς καὶ τῆς ὑπὸ BHG γωνίας ἰσάκεις πολλαπλασίων ἢ τε BA περιφέρεια καὶ ἡ ὑπὸ BHA γωνία, τῆς δὲ EZ περιφέρειᾶς καὶ τῆς ὑπὸ $E\Theta Z$ γωνίας ἢ τε EN περιφέρεια καὶ ἡ ὑπὸ $E\Theta N$ γωνία. καὶ δέδεικται, ὅτι εἰ ὑπερέχει ἡ BA περιφέρεια τῆς EN περιφέρειᾶς, ὑπερέχει καὶ ἡ ὑπὸ BHA γωνία τῆς ὑπὸ $E\Theta N$ γωνίας, καὶ εἰ ἴση, ἴση, καὶ εἰ ἐλάσσων, ἐλάσσων. ἔστιν ἄρα, ὡς ἡ $B\Gamma$ περιφέρεια πρὸς τὴν EZ , οὕτως ἡ ὑπὸ BHG γωνία πρὸς τὴν ὑπὸ $E\Theta Z$. ἀλλ' ὡς ἡ ὑπὸ BHG γωνία πρὸς τὴν ὑπὸ $E\Theta Z$, οὕτως ἡ ὑπὸ BAG πρὸς τὴν ὑπὸ $E\Delta Z$. διπλασία γὰρ ἑκατέρα ἑκατέρας. καὶ ὡς ἄρα ἡ $B\Gamma$ περιφέρεια πρὸς τὴν EZ περιφέρειαν, οὕτως ἢ τε ὑπὸ BHG γωνία πρὸς τὴν ὑπὸ $E\Theta Z$ καὶ ἡ ὑπὸ BAG πρὸς τὴν ὑπὸ $E\Delta Z$.

Ἐν ἄρα τοῖς ἴσοις κύκλοις αἱ γωνίαι τὸν αὐτὸν ἔχουσι λόγον ταῖς περιφερειαῖς, ἐφ' ὧν βεβήκασιν, ἐάν τε πρὸς τοῖς κέντροις ἐάν τε πρὸς ταῖς περιφερειαῖς ὡς βεβηκυῖαι· ὅπερ ἔδει δεῖξαι.

times is angle BGL also (divisible) by BGC . And so, for the same (reasons), as many times as circumference NE is (divisible) by EF , so many times is angle NHE also (divisible) by EHF . Thus, if circumference BL is equal to circumference EN then angle BGL is also equal to EHN [Prop. 3.27], and if circumference BL is greater than circumference EN then angle BGL is also greater than EHN ,[†] and if (BL is) less (than EN then BGL is also) less (than EHN). So there are four magnitudes, two circumferences BC and EF , and two angles BGC and EHF . And equal multiples have been taken of circumference BC and angle BGC , (namely) circumference BL and angle BGL , and of circumference EF and angle EHF , (namely) circumference EN and angle EHN . And it has been shown that if circumference BL exceeds circumference EN then angle BGL also exceeds angle EHN , and if (BL is) equal (to EN then BGL is also) equal (to EHN), and if (BL is) less (than EN then BGL is also) less (than EHN). Thus, as circumference BC (is) to EF , so angle BGC (is) to EHF [Def. 5.5]. But as angle BGC (is) to EHF , so (angle) BAC (is) to EDF [Prop. 5.15]. For the former (are) double the latter (respectively) [Prop. 3.20]. Thus, also, as circumference BC (is) to circumference EF , so angle BGC (is) to EHF , and BAC to EDF .

Thus, in equal circles, angles have the same ratio as the (ratio of the) circumferences on which they stand, whether they are standing at the centers (of the circles) or at the circumferences. (Which is) the very thing it was required to show.

[†] This is a straight-forward generalization of Prop. 3.27